

**Table 2.2. HPV genotypes in cytologically normal women and those with low-grade cytological or histological lesions**

		15 613 cytology-normal women (Clifford et al., Lancet 2005)#	5910 women with LSIL/CIN1 (Clifford et al., CEBP, 2005)#++
HPV-negative		90.8	28.9
HPV-positive		9.2	71.1
Ca types	16	1.8	26.6
	58	0.7	8.5
	31	0.7	11.7
	18	0.7	8.6
	56	0.6	9.7
	35	0.5	5.9
	33	0.5	7.6
	45	0.5	4.9
	52	0.5	8.8
	39	0.4	7.8
	51	0.4	10.9
	59	0.3	6.0
	68	0.2	
	73	0.1	
	82	0.1	
Non ca types	42	0.9	
	81	0.6	
	70	0.4	
	66	0.4	8.5
	40	0.3	
	72	0.2	
	43	0.2	
	cand89	0.2	
	54	0.2	
	84	0.1	
	6	0.1	8.1
	11	0.1	
	34	0.1	
	83	0.1	
	53**	0.1	10.1
	44	0.1	
	55	0.1	
	26	<0.1	
	61**	<0.1	
	57	<0.1	
	71	<0.1	

# Any appearance, single or multiple infections

\*\* The very low prevalence of these types among cytologically normal women is partly due to primer choice in the large majority of the summarized studies

++ Only most prevalent 15 types were shown

Compiled by the Working Group from two IARC meta-analyses.