

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments	
Bidel et al. (2010) Finland Enrolment 1972–2002; Mean FU 18 years Cohort	60 041; men and women (age 26–74 years) without history of any cancer at baseline Exposure assessment method: Questionnaire	Colon & rectum	All coffee (cups/day)			Age, study year, education, cigarette smoking, alcohol consumption, leisure time physical activity, history of diabetes, tea consumption, and body mass index	Strengths: homogenous study population; large number of participants randomly selected from the whole population; long follow-up without any loss Limitations: self-reported data; possible misclassification of the exposure; lack of data on caffeine or the other nutrients; not completely excluded the effects of residual confounding	
			Men	267	-			
			0	12	1			
			1–2	32	1.18 (0.6–2.32)			
			3–4	74	1.27 (0.68–2.38)			
			5–6	85	1.06 (0.57–2)			
			7–9	41	1.15 (0.59–2.26)			
			≥ 10	23	0.98 (0.47–2.03)			
		Trend-test p-value: 0.86						
		Colon & rectum	All coffee (cups/day)					
			Women	271	-			
			0	8	1			
			1–2	28	1.37 (0.62–3.05)			
			3–4	88	1.5 (0.71–3.15)			
5–6	97		1.31 (0.62–2.78)					
7–9	38		1.53 (0.69–3.4)					
≥ 10	12	1.24 (0.49–3.14)						
Trend-test p-value: 0.83								

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
		Colon & rectum	All coffee (cups/day)				
			Total	538	-		
			0	20	1		Age, study year, education, cigarette smoking, alcohol consumption, leisure time physical activity, history of diabetes, tea consumption, and body mass index, sex
			1–2	60	1.25 (0.74–2.09)		
			3–4	162	1.32 (0.82–2.14)		
			5–6	182	1.14 (0.71–1.85)		
			7–9	79	1.28 (0.76–2.13)		
			≥ 10	35	1.03 (0.58–1.83)		
			Trend-test p-value: 0.61				
		Colon	All coffee (cups/day)				
			Men	150	-		
			0	11	1		Age, study year, education, cigarette smoking, alcohol consumption, leisure time physical activity, history of diabetes, tea consumption, and body mass index
			1–2	24	1.02 (0.46–2.25)		
			3–4	42	0.8 (0.38–1.71)		
			5–6	49	0.77 (0.37–1.63)		
			7–9	24	0.72 (0.31–1.64)		
			≥ 10	16	0.74 (0.31–1.79)		
			Trend-test p-value: 0.89				

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
		Colon	All coffee (cups/day)				
			Women	154	-		
			0	6	1		Age, study year, education, cigarette smoking, alcohol consumption, leisure time physical activity, history of diabetes, tea consumption, and body mass index
			1–2	16	0.81 (0.3–2.2)		
			3–4	60	1.39 (0.58–1.71)		
			5–6	57	1.06 (0.44–2.59)		
			7–9	24	1.32 (0.51–3.42)		
			≥ 10	4	0.61 (0.16–2.23)		
			Trend-test p-value: 0.31				
		Colon	All coffee (cups/day)				
			Total	304	-		
			0	17	1		Age, study year, education, cigarette smoking, alcohol consumption, leisure time physical activity, history of diabetes, tea consumption, and body mass index, sex
			1–2	40	0.93 (0.5–1.74)		
			3–4	102	1.04 (0.59–1.83)		
			5–6	106	0.87 (0.49–1.54)		
			7–9	48	0.93 (0.5–1.73)		
			≥ 10	20	0.72 (0.35–1.47)		
			Trend-test p-value: 0.76				

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
		Rectum	All coffee (cups/day)				
			Men	117	-		
			0	5	1		Age, study year, education, cigarette smoking, alcohol consumption, leisure time physical activity, history of diabetes, tea consumption, and body mass index
			1–2	10	1.6 (0.43–5.94)		
			3–4	37	2.71 (0.81–9.03)		
			5–6	43	1.95 (0.58–6.55)		
			7–9	24	2.51 (0.72–8.78)		
			≥ 10	10	1.68 (0.43–6.48)		
			Trend-test p-value: 0.32				
		Rectum	All coffee (cups/day)				
			Women	117	-		
			0	2	1		Age, study year, education, cigarette smoking, alcohol consumption, leisure time physical activity, history of diabetes, tea consumption, and body mass index
			1–2	18	2.97 (0.67–13.1)		
			3–4	32	1.89 (0.44–8.13)		
			5–6	47	2.07 (0.48–8.83)		
			7–9	16	2.24 (0.49–10.1)		
			≥ 10	8	2.92 (0.6–14.3)		
			Trend-test p-value: 0.53				

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
		Rectum	All coffee (cups/day)				
			Total	117	-		
			0	7	1		
			1–2	28	2.19 (0.83–5.79)		
			3–4	69	2.19 (0.87–5.53)		
			5–6	90	1.96 (0.77–4.95)		
			7–9	40	2.31 (0.88–6.03)		
			≥ 10	18	1.99 (0.71–5.55)		
			Trend-test p-value: 0.62				
Yamada et al. (2014) Japan Enrolment 1988–1990; Follow up to 2009 Cohort	58 221; 23 607 men & 34 614 women (aged 40–79 years at baseline) without a history of colorectal cancer Exposure assessment method: Questionnaire	Colon	All coffee (cups/day)				
			Men	355	-		
			< 1	240	1		
			1	44	1.06 (0.76–1.47)		
			2–3	58	1.26 (0.93–1.7)		
			≥ 4	13	1.79 (1.01–3.18)		
			Trend-test p-value: 0.03				
						Age, study year, education, cigarette smoking, alcohol consumption, leisure time physical activity, history of diabetes, tea consumption, and body mass index, sex	
						Age, smoking, drinking, family history of colorectal cancer, education, body mass index, walking time, and regular meat consumption, and district	Strengths: large population-based study Limitations: data collected at the baseline survey only; self-reported intake data

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
		Colon	All coffee (cups/day)				
			Women	332	-		Age, smoking, drinking, family history of colorectal cancer, education, body mass index, walking time, and regular meat consumption, and district
			< 1	254	1		
			1	46	1 (0.72–1.37)		
			2–3	27	0.86 (0.57–1.3)		
			≥ 4	5	2.02 (0.81–5.03)		
			Trend-test p-value: 0.96				
		Rectum	All coffee (cups/day)				
			Men	202	-		Age, smoking, drinking, family history of colorectal cancer, education, body mass index, walking time, and regular meat consumption, and district
			< 1	139	1		
			1	28	1.19 (0.79–1.8)		
			2–3	30	1.12 (0.75–1.7)		
			≥ 4	5	1.19 (0.48–2.95)		
			Trend-test p-value: 0.53				
		Rectum	All coffee (cups/day)				
			Women	112	-		Age, smoking, drinking, family history of colorectal cancer, education, body mass index,
			< 1	82	1		

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments	
			1	13	0.88 (0.48–1.59)	walking time, and regular meat consumption, and district		
			2–3	17	1.55 (0.89–2.69)			
			≥ 4	0	0			
			Trend-test p-value: 0.37					
		Colon & rectum	All coffee (cups/day)					
			Men	557	-	Age, smoking, drinking, family history of colorectal cancer, education, body mass index, walking time, and regular meat consumption, and district		
			< 1	379	1			
			1	72	1.11 (0.86–1.43)			
			2–3	88	1.21 (0.95–1.54)			
			≥ 4	18	1.57 (0.97–2.55)			
			Trend-test p-value: 0.03					
		Colon & rectum	All coffee (cups/day)					
			Women	444	-	Age, smoking, drinking, family history of colorectal cancer, education, body mass index, walking time, and regular meat consumption, and district		
			< 1	336	1			
			1	59	0.97 (0.73–1.28)			
			2–3	44	1.04 (0.75–1.44)			
			≥ 4	5	1.42 (0.57–3.5)			
			Trend-test p-value: 0.61					

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
		Colon	All coffee (cups/day)				
			Men	355	-		Age, drinking, family history of colorectal cancer, education, body mass index, walking time, and regular meat consumption
			Current smoker	173	-		
			< 1	109	1		
			1	21	1.08		
			2–3	34	1.21		
			≥ 4	9	1.68		
			Trend-test p-value: 0.13				
		Colon	All coffee (cups/day)				
			Men	355	-		
			Never smoker	61	-		
			< 1	42	1		
			1	7	1.18		
			2–3	10	1.85		
			≥ 4	2	5.58		
			Trend-test p-value: 0.01				



Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
		Colon	All coffee (cups/day)				
			Women	332	-		Age, drinking, family history of colorectal cancer, education, body mass index, walking time, and regular meat consumption
			Current smoker	12	-		
			< 1	4	1		
			1	1	1.03		
			2–3	5	4.71		
			≥ 4	2	6.06		
			Trend-test p-value: 0.02				
		Colon	All coffee (cups/day)				
			Women	332	-		
			Never smoker	279	-		
			< 1	222	1		
			1	37	1.03		
			2–3	17	1.46		
			≥ 4	3	1.69		
			Trend-test p-value: 0.6				

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
		Colon & rectum	All coffee (cups/day)				
			Men	557	-		Age, drinking, family history of colorectal cancer, education, body mass index, walking time, and regular meat consumption
			Current smoker	267	-		
			< 1	168	1		
			1	35	1.17		
			2–3	53	1.21		
			≥ 4	11	1.32		
			Trend-test p-value: 0.2				
		Colon & rectum	All coffee (cups/day)				
			Men	557	-		
			Never smoker	105	-		
			< 1	71	1		
			1	16	1.57		
			2–3	14	1.52		
			≥ 4	4	5.92		
			Trend-test p-value: 0.01				

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
		Colon & rectum	All coffee (cups/day)				
			Women	444	-		Age, drinking, family history of colorectal cancer, education, body mass index, walking time, and regular meat consumption
			Current smoker	16	-		
			< 1	6	1		
			1	1	0.61		
			2–3	7	3.65		
			≥ 4	2	2.87		
			Trend-test p-value: 0.04				
		Colon & rectum	All coffee (cups/day)				
			Women	444	-		
			Never smoker	374	-		
			< 1	292	1		
			1	47	0.98		
			2–3	32	1		
			≥ 4	3	1.46		
			Trend-test p-value: 0.8				

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
Phillips & Snowdon (1985) USA Enrolment, 1960; FU 21 years Cohort	25 493; white men and women California Seventh-Day Adventists aged ≥ 30 years at baseline Exposure assessment method: Questionnaire	Colon & rectum	All coffee (cups/day)			Age, sex	Strengths: unique characteristics of the Adventist population; linkage with registries; FFQ used by the American Cancer Society. Limitations: all the dietary data come from a 1960 self-administered questionnaire; limited accuracy of the underlying cause of death as recorded on death certificates; inferences with regard to risk of CRC; no adjustment for confounders; possibility that subjects may have substantially changed their dietary habits or weigh during long follow-up period Strengths: unique characteristics of the Adventist population. Limitations: all the dietary data come from a 1960 self-administered questionnaire; limited accuracy of the underlying cause of death as recorded on death certificates; inferences with regard to risk of CRC; possibility that subjects may have substantially changed their dietary habits or weigh during long follow-up period
			< 1	110	1		
			1	20	1.5 (0.9–2.3)		
			≥ 2	34	1.5 (1–2.2)		
			Trend-test p-value: 0.02				

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments			
Hartman et al. (1998) Finland Enrolment 1985–1988; Median FU 8 years Cohort	27 111 subjects; male smokers aged 50 and 69 years; men who were alcoholics, who had cirrhosis of the liver, severe angina with exertion, or chronic renal insufficiency, who had been previously diagnosed with cancer, or who had been taking supplements of vitamin E or A or β-carotene in excess of defined amounts or receiving anticoagulant therapy were excluded. Exposure assessment method: Questionnaire	Colon	All coffee (cups/day)			Age, intervention group, calcium, occupational physical activity, BMI	Strengths: all important available measured as confounders Limitations: the distribution and levels of intake of coffee were very different from each other; inability to use nondrinkers of coffee as the referent group; not ruled out the possibility that residual confounding contributed to the results			
			≤ 4	51	1					
			> 4	31	0.73 (0.47–1.16)					
					> 6			24	0.69 (0.42–1.13)	
					Trend-test p-value: 0.11					
		Rectum	All coffee (cups/day)						Age, intervention group, calcium, occupational physical activity, BMI, cholesterol	
			≤ 4	33	1					
> 4	29		1.05 (0.63–1.75)							
> 6	17		0.77 (0.43–1.4)							
			Trend-test p-value: 0.44							
Naganuma et al. (2007) Japan Enrolment from 1990; FU 11.6 years Cohort	38 701; 18 867 men and 19 834 women aged 40–64 years at the baseline with no previous history of cancer Exposure assessment method: Questionnaire	Colon & rectum	All coffee			Age, sex, family history of colorectal cancer, education level, BMI, walking time, smoking, alcohol drinking, tea consumption, consumption of meat, vegetables, fruits, total caloric intake	Strengths: population based design; large number of cases; long follow-up period; high validity and reproducibility of coffee consumption assessment Limitations: collection of coffee consumption data only once before follow-up period; no information on type of coffee consumed (filtered/boiled, caffeinated/decaffeinated)			
			Never	93	1					
			Occasionally	195	1.14 (0.87–1.46)					
			1–2 cups/day	123	0.98 (0.74–1.32)					
			≥ 3 cups/day	46	0.95 (0.65–1.39)					
			Trend-test p-value: 0.55							

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
Domianni et al. (2013) Muticenters (England & USA) 1993–2001, median follow up of 11.4 years Cohort	57 398; men and women aged 55–74 years in the National Cancer Institute-Prostate, Lung, Colorectal and Ovarian (PLCO) cancer screening trial enrolled from 10 centres (England and the United States) Exposure assessment method: Questionnaire	Colon & rectum	All coffee (cups/day)			Age, BMI, smoking, sex, race, family history of colorectal cancer, education, physical activity, NSAID intake, history of diabetes, number of colorectal examinations up to 3 years before the start of study, hormone use (among women), fruit intake (servings per day), vegetable intake (servings per day), meat intake (g per day), alcohol intake (g per day) and study centre	Strengths: collection of diet and demographic information before diagnosis of cancer; collection of extensive baseline and diet intake information allowing for appropriate control of confounders. The majority of the cohort (83.3%) has complete sigmoidoscopy at baseline, so all participants had an equal opportunity to be screened, unlike any screening-related biases. Limitations: Conducted in the screening arm of a randomized controlled trial of CRC screening, results may not be generalizable to all cases; Inherent limitation of dietary questionnaires; Limited power to detect an association in specific subgroups, such as by location (proximal, distal or rectal) or coffee type (caffeinated vs decaffeinated)
			None	73	1		
			< 1	138	0.94 (0.7–1.25)		
			1	147	0.94 (0.7–1.26)		
			2–3	207	1.03 (0.77–1.37)		
			≥ 4	116	1.08 (0.79–1.48)		
			Trend-test p-value: 0.229				

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
Dik et al. (2014) EPIC Enrollment 1992–2000; median FU 11.6 years Cohort	521, 448; men (29.8%) and women (70%) aged between 25 and 70 years without any type of prevalent cancer at enrolment, carcinoma in situ, unknown histology of the tumour, unknown first incidence tumour or a colorectal tumour originating from other organs, within 23 centres in 10 different European countries Exposure assessment method: Questionnaire	Colon & rectum	All coffee			BMI, diabetes, menopausal status, hormone replacement therapy, physical activity, education level, smoking, baseline intake of energy from fat, energy from non-fat, alcohol, fibres, dairy products, red meat, processed meat	Prospective cohort studies have found no significant associations with either proximal colon, distal colon or rectal cancer. Strengths: prospective population-based design; multiple populations; considerable follow up; large number of participants and cases; performing analyses by anatomical subsite studying differences between caffeinated and decaffeinated coffee Limitations: the self-reported consumption; single assessment; absence of data on brewing methods, cup size and levels of caffeine, cafestol, kahweol and antioxidants
			Non/low	914	-		
			Moderately low	761	1.04 (0.94–1.15)		
			Moderate	694	1.06 (0.95–1.19)		
			Moderately high	863	0.99 (0.89–1.1)		
			High	1002	1.06 (0.95–1.18)		
Per 100 ml/day	NR	1.01 (0.99–1.02)					
			Trend-test p-value: 0.58				
Larsson et al. (2006) Sweden Enrolment 1987–1990; Follow up to 2004 Cohort	106 739 (pooled two cohorts); 61 433 women aged 40–76 years (Swedish Mammography Cohort) and 45 306 men aged 45–79 years (Cohort of Swedish Men), excluded subjects who diagnosed with cancer	Colon & rectum	All coffee (cups/day)			Age, education, BMI, family history of colorectal cancer, history of diabetes, smoking, physical activity, aspirin use, multivitamin supplement, daily intake of calories, fruit, vegetables,	Approximately 15% of population consumed boiled coffee only or both boiled and filtered coffee. Decaffeinated coffee is very uncommon in Sweden. Strengths: two large population-based cohorts Prospective design eliminated recall bias
			< 1	103	1		
			1	213	1.26 (0.99–1.6)		
			2–3	701	1.19 (0.96–1.47)		
			≥ 4	262	1.14 (0.9–1.44)		

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
	(except nonmelanoma skin cancer) before baseline Exposure assessment method: Questionnaire		One additional cup/day	1279	1 (0.97–1.04)	milk, and red meat, postmenopausal hormone form women	Large number of cases of CRC Information on many potential risk factors for CRC The use of data from two completely separate cohorts Completeness of follow-up Limitations: self-reported coffee consumption No data on type of coffee
Michels et al. (2005) USA Enrolment from 1980 (NHS), from 1986 (HPFS); Follow up to 1998 Cohort	133 893 (pooled two cohorts); 87 794 women aged 30–55 years at enrolment [Nurses' Health Study (NHS)] and 46 099 men aged 40–75 at enrolment [Health Professionals' Follow-up Study (HPFS)], excluded subjects who diagnosed with Crohn disease, ulcerative colitis, or cancers other than nonmelanoma skin cancer at baseline Exposure assessment method: Questionnaire	Colon & rectum	Caffeinated coffee (cups/day)			Age, family history of cancer, history of sigmoidoscopy, height, BMI, smoking pack-years, physical activity, aspirin use, vitamin supplements, alcohol consumption, red meat intake, total calorie intake, menopause, postmenopausal hormone use	Strengths: large sample size, repeated measurements of diet over time, adjustment for screening history Limitations: –
			Total subjects	1431	-		
			Never	259	1		
			0.5	243	1.05 (0.88–1.26)		
			1	228	0.99 (0.82–1.18)		
			2–3	497	1.02 (0.87–1.19)		
			4–5	164	0.98 (0.8–1.21)		
			> 5	40	0.98 (0.69–1.38)		
			1 additional cup/day	1431	0.99 (0.96–1.03)		
			Trend-test p-value: 0.6				



Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
		Colon & rectum	Caffeinated coffee (cups/day)				
			Non-smoker	1094	-		Age, family history of cancer, history of sigmoidoscopy, height, BMI, smoking pack-years, physical activity, aspirin use, vitamin supplements, alcohol consumption, red meat intake, total calorie intake, menopause, postmenopausal hormone use
			Never	188	1		
			0.5	160	1.02 (0.82–1.27)		
			1	181	0.99 (0.8–1.22)		
			2–3	405	0.99 (0.82–1.19)		
			4–5	130	0.94 (0.75–1.19)		
			> 5	30	0.92 (0.62–1.36)		
			1 additional cup/day	1094	0.99 (0.96–1.03)		
			Trend-test p-value: 0.55				
		Colon & rectum	Decaffeinated coffee (cups/day)				
			Total subjects	1138	-		Age, family history of cancer, history of sigmoidoscopy, height, BMI, smoking pack-years, physical activity, aspirin use, vitamin supplements, alcohol consumption, red meat intake, total calorie intake, menopause,
			Never	463	1		
			0.25	212	0.82 (0.69–0.97)		
			0.5	167	0.7 (0.58–0.85)		
			1–1.9	146	0.74 (0.61–0.9)		
			> 2	150	0.82 (0.67–0.99)		

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
			1 additional cup/day	1138	0.96 (0.91–1.02)	postmenopausal hormone use	
			Trend-test p-value: 0.08				
Nilsson et al. (2010) Sweden Enrolment from 1992; Follow up to 15 years Cohort	64 603; 32 425 men and 32 178 women aged 30–39 at the baseline Exposure assessment method: Questionnaire	Colon & rectum	All coffee (occasions/day)			Age, sex, BMI, smoking, education, and recreational physical activity	Strengths: large sample size; prospective design; long follow-up; data collection before the diagnosis of cases with cancer; population-based data; Limitations: residual confounding by factors not addressed or not sufficiently measured in this study may have been present
			< 1	19	1		
			1–3	206	1.56 (0.96–2.54)		
				≥ 4	96	1.43 (0.86–2.38)	
		Colon & rectum	Filtered coffee (occasions/day)			Age, sex, BMI, smoking, education, and recreational physical activity	
			< 1	87	1		
1–3	196		1.18 (0.91–1.52)				
			≥ 4	38	0.73 (0.5–1.08)		
Sinha et al. (2012) USA Enrolment 1995–1996; FU 10.5 years Cohort	489 706; men and women aged 50–71 years without prevalent cancer or end-stage renal disease at baseline Exposure assessment method: Questionnaire	Colon & rectum	All coffee (cups/day)			Age, race, education, smoking status, time since quitting for former smokers, smoking dose, ever smoke a pipe or cigar, diabetes, colorectal screening, family history of colorectal cancer, regular nonsteroidal anti-inflammatory drug use, marital status, BMI,	Risk estimates were presented stratified by type of coffee (decaffeinated versus caffeinated) and organ sites (colon (proximal, distal), rectal). There was no difference in risk by caffeine content, but colon and proximal colon showed a similar pattern to the CRC results, while distal colon and rectal did not. Strengths: large size; prospective design; wide range of coffee consumption

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
			None	647	1	frequency of vigorous physical activity, calories, fruit and vegetables, red meat, dietary calcium intake, alcohol, and menopausal hormone therapy in women	Limitations: self-reported coffee intake at a single time point; not ruled out reverse causality due to lack of information on inflammatory bowel disease or other conditions that may have caused participants to reduce or eliminate coffee consumption; predominantly college educated non-Hispanic white with healthy lifestyle
			< 1 cup/wk	1083	0.97 (0.88–1.07)		
			1	1206	1 (0.9–1.1)		
			2–3	2999	0.98 (0.9–1.07)		
			4–5	785	0.87 (0.78–0.97)		
			≥ 6	226	0.8 (0.69–0.94)		
			Trend-test p-value: 0.001				
Oba et al. (2006) Japan Enrolment from 1992; Follow up to 2000 Cohort	30 221; 13 894 men and 16 327 women aged ≥ 35 years at the baseline, without a history of cancer other than non-melanoma skin cancer Exposure assessment method: Questionnaire	Colon	All coffee			Age, height, BMI, total pack-years of cigarette smoking, alcohol intake, and physical activity, and tea intake	Strengths: prospective cohort study; diet data collection ahead of the diagnosis of colon cancer; community-based population; high response rate (92.0%) Limitations: small number of cases; possible misclassification for colon cancer
			Men	111	-		
			Never to < 1 cup/month	29	1		
			Once cup/month to < 1 cup/day	52	1.13 (0.71–1.81)		
			One cup/day or more	30	0.81 (0.46–1.42)		
			Trend-test p-value: 0.14				

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
		Colon	All coffee				
			Women	102	-		
			Never to < 1 cup/month	45	1		Age, height, BMI, total pack-years of cigarette smoking, alcohol intake, and physical activity, and tea intake
			Once cup/month to < 1 cup/day	44	0.92 (0.59–1.43)		
			One cup/day or more	13	0.43 (0.22–0.85)		
Peterson et al. (2010) Singapore Enrolment 1993–1998; FU 12 year Cohort	61 321; men and women aged 45–74 years at enrolment, excluded subjects with a history of cancer diagnosis at baseline Exposure assessment method: Questionnaire	Colon	All coffee (cups/day)				
			All subjects	591	-		
			< 1	177	1		
			1	228	1.04 (0.85–1.26)		
			2+	186	0.9 (0.73–1.11)		
			Trend-test p-value: 0.31				
		Colon	All coffee (cups/day)				
			Never smokers	384	-		
			< 1	125	1		Age, sex, BMI, cigarette smoking, alcohol consumption, physical activity, family history of cancer, intake of green tea
			1	147	1 (0.78–1.27)		
			2+	112	1.01 (0.78–1.31)		
			Trend-test p-value: 0.93				
							Strengths: collection of information on coffee consumption and other dietary and lifestyle factors before cancer diagnosis; ruled out the possibility of recall bias and reverse causality; genetically homogenous study population Limitations: –

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
		Colon	All coffee (cups/day)			Age, sex, BMI, cigarette smoking, alcohol consumption, physical activity, family history of cancer, intake of green tea	
			Ever smokers	207	-		
			< 1	52	1		
			1	81	1.13 (0.79–1.6)		
			2+	74	0.77 (0.54–1.11)		
			Trend-test p-value: 0.1				
Simons et al. (2010) Netherlands Enrolment 1986; FU 13.3 years Cohort	120 852; 1832 male and 2045 female subcohort member (aged 55–69 years) without a history of cancers other than nonmelanoma skin cancer at baseline used for case-cohort analyses Exposure assessment method: Questionnaire	Colon & rectum	All coffee (cups/day)			Age, family history of CRC, physical activity at the longest held job, non-occupational physical activity, smoking status, education level, BMI, ethanol intake, meat intake, processed meat intake, folate intake, vitamin B6 intake, fibre intake, fluid intake from other fluids	Strengths: No Limitations: No
			Men	1443	-		
			≤ 2	200	1		
			> 2–4	504	0.98 (0.78–1.23)		
			> 4–6	410	1.06 (0.83–1.35)		
			> 6	146	1 (0.74–1.36)		
			Trend-test p-value: 0.67				
		Rectum	All coffee (cups/day)			Age, family history of CRC, physical activity at the longest held job, non-occupational physical activity, smoking status, education level, BMI, ethanol	
			Men	332	-		
			≤ 2	36	1		
			> 2–4	120	1.32 (0.87–1.99)		

Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
			> 4–6	116	1.5 (0.97–2.31)		
			> 6	50	1.6 (0.96–2.66)		intake, meat intake, processed meat intake, folate intake, vitamin B6 intake, fibre intake, fluid intake from other fluids
			Trend-test p-value: 0.05				
		Colon & rectum	All coffee (cups/day)				Age, family history of CRC, physical activity at the longest held job, non-occupational physical activity, smoking status, education level, BMI, ethanol intake, meat intake, processed meat intake, folate intake, vitamin B6 intake, fibre intake, fluid intake from other fluids
			Women	1040	-		
			≤ 2	206	1		
			> 2–4	425	0.91 (0.74–1.13)		
			> 4–6	245	0.96 (0.75–1.23)		
			> 6	63	1.07 (0.74–1.55)		
			Trend-test p-value: 0.85				
		Rectum	All coffee (cups/day)				Age, family history of CRC, physical activity at the longest held job, non-occupational physical activity, smoking status, education level, BMI, ethanol intake, meat intake, processed meat intake, folate intake, vitamin B6 intake,
			Women	173	-		
			≤ 2	40	1		
			> 2–4	73	0.8 (0.53–1.2)		
			> 4–6	42	0.79 (0.49–1.27)		
			> 6	18	1.41 (0.75–2.63)		



Table 2.22 Cohort studies on cancer of the colorectum and coffee drinking (web only)

Reference, location enrolment/follow-up period, study design	Population size, description, exposure assessment method	Organ site	Exposure category or level	Exposed cases/deaths	Risk estimate (95% CI)	Covariates controlled	Comments
Lukic et al. (2016) Norway Enrolment 1996–2004; FU 6–8 years Cohort	91 767; women aged 30–70 at the baseline excluding subjects with prevalent cancer other than non-melanoma skin cancer at baseline Exposure assessment method: Questionnaire	Colon & rectum	All coffee (cups/day)			Smoking status, age at smoking initiation, number of pack-years smoked, duration of education, physical activity level, use of oral contraceptives, and total energy intake	The interaction between coffee consumption and smoking status for colorectal cancer risk was also tested, but no interaction (data not shown) Strengths: prospective design; relatively large sample size; sampling from general population; validated FFQ; repeated measurements of coffee consumption and smoking exposure Limitations: no information on type of coffee; not excluded the effect of residual confounding; self-reported coffee consumption data with misclassification bias
			Light consumers, ≤ 1	224	1		
			Low moderate consumers, more than 1 up to 3	423	0.95 (0.81–1.11)		
			High moderate consumers, more than 3 up to 7	541	0.83 (0.7–0.98)		
			Heavy consumers, > 7	78	0.98 (0.72–1.32)		
			Trend-test p-value: 0.1				

CI, confidence interval; CRC, colorectal cancer; FU, follow-up; NR, not reported



## References

- Bidel S, Hu G, Jousilahti P, Antikainen R, Pukkala E, Hakulinen T, et al. (2010). Coffee consumption and risk of colorectal cancer. *Eur J Clin Nutr.* 64(9):917–23. <http://dx.doi.org/10.1038/ejcn.2010.103> PMID:20571499
- Dik VK, Bueno-de-Mesquita HB, Van Oijen MG, Siersema PD, Uiterwaal CS, Van Gils CH, et al. (2014). Coffee and tea consumption, genotype-based CYP1A2 and NAT2 activity and colorectal cancer risk-results from the EPIC cohort study. *Int J Cancer.* 135(2):401–12. <http://dx.doi.org/10.1002/ijc.28655> PMID:24318358
- Domianni C, Huang WY, Berndt S, Hayes RB, Ahn J (2013). Prospective study of the relationship between coffee and tea with colorectal cancer risk: the PLCO Cancer Screening Trial. *Br J Cancer.* 109(5):1352–9. <http://dx.doi.org/10.1038/bjc.2013.434> PMID:23907431
- Hartman TJ, Tangrea JA, Pietinen P, Malila N, Virtanen M, Taylor PR, et al. (1998). Tea and coffee consumption and risk of colon and rectal cancer in middle-aged Finnish men. *Nutr Cancer.* 31(1):41–8. <http://dx.doi.org/10.1080/01635589809514676> PMID:9682247
- Larsson SC, Bergkvist L, Giovannucci E, Wolk A (2006). Coffee consumption and incidence of colorectal cancer in two prospective cohort studies of Swedish women and men. *Am J Epidemiol.* 163(7):638–44. <http://dx.doi.org/10.1093/aje/kwj067> PMID:16443798
- Lee KJ, Inoue M, Otani T, Iwasaki M, Sasazuki S, Tsugane S; JPHC Study Group (2007). Coffee consumption and risk of colorectal cancer in a population-based prospective cohort of Japanese men and women. *Int J Cancer.* 121(6):1312–8. <http://dx.doi.org/10.1002/ijc.22778> PMID:17450527
- Lukic M, Licaj I, Lund E, Skeie G, Weiderpass E, Braaten T (2016). Coffee consumption and the risk of cancer in the Norwegian Women and Cancer (NOWAC) Study. *Eur J Epidemiol.* <http://dx.doi.org/10.1007/s10654-016-0142-x> PMID:27010635
- Michels KB, Willett WC, Fuchs CS, Giovannucci E (2005). Coffee, tea, and caffeine consumption and incidence of colon and rectal cancer. *J Natl Cancer Inst.* 97(4):282–92. <http://dx.doi.org/10.1093/jnci/dji039> PMID:15713963
- Naganuma T, Kuriyama S, Akhter M, Kakizaki M, Nakaya N, Matsuda-Ohmori K, et al. (2007). Coffee consumption and the risk of colorectal cancer: a prospective cohort study in Japan. *Int J Cancer.* 120(7):1542–7. <http://dx.doi.org/10.1002/ijc.22505> PMID:17205519
- Nilsson LM, Johansson I, Lenner P, Lindahl B, Van Guelpen B (2010). Consumption of filtered and boiled coffee and the risk of incident cancer: a prospective cohort study. *Cancer Causes Control.* 21(10):1533–44. <http://dx.doi.org/10.1007/s10552-010-9582-x> PMID:20512657
- Oba S, Shimizu N, Nagata C, Shimizu H, Kametani M, Takeyama N, et al. (2006). The relationship between the consumption of meat, fat, and coffee and the risk of colon cancer: a prospective study in Japan. *Cancer Lett.* 244(2):260–7. <http://dx.doi.org/10.1016/j.canlet.2005.12.037> PMID:16519996
- Peterson S, Yuan JM, Koh WP, Sun CL, Wang R, Turesky RJ, et al. (2010). Coffee intake and risk of colorectal cancer among Chinese in Singapore: the Singapore Chinese Health Study. *Nutr Cancer.* 62(1):21–9. <http://dx.doi.org/10.1080/01635580903191528> PMID:20043256

- Phillips RL, Snowdon DA (1985). Dietary relationships with fatal colorectal cancer among Seventh-Day Adventists. *J Natl Cancer Inst.* 74(2):307–17. PMID:3856044
- Simons CC, Leurs LJ, Weijnenberg MP, Schouten LJ, Goldbohm RA, van den Brandt PA (2010). Fluid intake and colorectal cancer risk in the Netherlands Cohort Study. *Nutr Cancer.* 62(3):307–21. <http://dx.doi.org/10.1080/01635580903407098> PMID:20358468
- Sinha R, Cross AJ, Daniel CR, Graubard BI, Wu JW, Hollenbeck AR, et al. (2012). Caffeinated and decaffeinated coffee and tea intakes and risk of colorectal cancer in a large prospective study. *Am J Clin Nutr.* 96(2):374–81. <http://dx.doi.org/10.3945/ajcn.111.031328> PMID:22695871
- Yamada H, Kawado M, Aoyama N, Hashimoto S, Suzuki K, Wakai K, et al.; JACC Study Group (2014). Coffee consumption and risk of colorectal cancer: the Japan Collaborative Cohort Study. *J Epidemiol.* 24(5):370–8. <http://dx.doi.org/10.2188/jea.JE20130168> PMID:24857957