

### 2.2.22 Cancers of the head and neck

Head and neck cancer refers to a group of cancers that develop in (i) the oral cavity; (ii) the pharynx, including the nasopharynx, the oropharynx, and the hypopharynx; (iii) the larynx; (iv) the paranasal sinuses and the nasal cavity; and (v) the salivary glands.

Most head and neck cancers are squamous cell carcinomas. Because of the established associations of head and neck cancer with tobacco use, and because BMI is inversely associated with tobacco use, it is important that associations of BMI with risk of head and neck cancers carefully consider potential confounding and/or effect modification by tobacco use. Notably, only prospective studies with at least 50 cases for any specific site were included in this review.

In 2001, the Working Group of the *IARC Handbook* on weight control and physical activity ([IARC, 2002](#)) concluded that the evidence of an association between avoidance of weight gain and cancers of the head and neck was *inadequate*.

#### (a) Cohort studies

See Table 2.2.22a (web only, available at: <http://publications.iarc.fr/570>).

##### (i) Cancer of the oral cavity

The association between BMI and risk of cancer of the oral cavity has been examined in two individual prospective studies ([Bhaskaran et al., 2014](#); [Etemadi et al., 2014](#)) and in a large pooled analysis of data from 20 prospective studies ([Gaudet et al., 2015](#)). All of these studies adjusted for both tobacco use and alcohol consumption. In the United Kingdom data linkage study of more than 5 million men and women, there was a statistically significant inverse association (RR per 5 kg/m<sup>2</sup> increase in BMI, 0.81; 95% CI, 0.74–0.89;  $P_{\text{trend}} < 0.0001$ ) ([Bhaskaran et al., 2014](#)). No significant association was observed in the NIH-AARP cohort study in the USA ([Etemadi](#)

[et al., 2014](#)) or in the large pooled analysis ([Gaudet et al., 2015](#)).

In contrast, quartiles of waist circumference were positively associated with risk (RR for highest vs lowest quartile, 2.00; 95% CI, 1.24–3.23;  $P_{\text{trend}} < 0.001$ ) in the NIH-AARP study ([Etemadi et al., 2014](#)). Similarly, in the large pooled analysis of 20 prospective studies, there was a 9% increase in risk (95% CI, 1.03–1.16) per 5 cm increase in waist circumference ( $P_{\text{trend}} = 0.006$ ) ([Gaudet et al., 2015](#)).

##### (ii) Cancers of the pharynx (nasopharynx, oropharynx, and/or hypopharynx)

There was no association between BMI and risk of nasopharyngeal cancer in the only study that assessed this relationship ([Samanic et al., 2004](#)). Similarly, there is no evidence that BMI is associated with risk of oropharyngeal cancer incidence ([Gaudet et al., 2012, 2015](#)) or mortality ([Gaudet et al., 2012](#)), or with hypopharyngeal cancer incidence ([Gaudet et al., 2015](#)). In the NIH-AARP cohort, BMI < 18.5 kg/m<sup>2</sup> was associated with a higher risk of oropharyngeal and hypopharyngeal cancer incidence compared with BMI 18.5– < 25 kg/m<sup>2</sup> ([Etemadi et al., 2014](#)). [There were only three cases in the exposed group.]

Waist circumference was not associated with oropharyngeal or hypopharyngeal cancer incidence in the NIH-AARP cohort study ([Etemadi et al., 2014](#)) or in the large pooled analysis ([Gaudet et al., 2015](#)).

##### (iii) Cancer of the larynx

Since 2000, there have been two individual prospective studies ([Samanic et al., 2004](#); [Etemadi et al., 2014](#)) and one large pooled analysis of 20 prospective studies ([Gaudet et al., 2015](#)) of the association between BMI and risk of cancer of the larynx (Table 2.2.22a, web only, available at: <http://publications.iarc.fr/570>). In the large study of more than 4.5 million United States Veterans, there was a statistically significantly lower risk

of laryngeal cancer for obese compared with non-obese White and Black men ([Samanic et al., 2004](#)). [Neither tobacco use nor alcohol consumption was included in the statistical model; therefore, confounding by these factors is likely.] In the NIH-AARP cohort study in the USA, in which both tobacco use and alcohol consumption were adjusted for in the model, BMI was not associated with risk of laryngeal cancer ([Etemadi et al., 2014](#)). Conversely, in the pooled analysis, there was a statistically significant positive association between BMI and risk (RR per 5 kg/m<sup>2</sup> increase, 1.42; 95% CI, 1.19–1.70) ([Gaudet et al., 2015](#)).

In the NIH-AARP study ([Etemadi et al., 2014](#)), there was no evidence of an association between waist circumference and risk of laryngeal cancer, whereas a weak positive association was reported in the pooled analysis (RR per 5 cm increase, 1.10; 95% CI, 0.99–1.22;  $P_{\text{trend}} = 0.08$ ) ([Gaudet et al., 2015](#)).

(iv) *Cancer of the oral cavity, pharynx, and larynx combined*

In two studies, the Asia-Pacific Cohort Studies Collaboration ([Parr et al., 2010](#)) and the Cancer Prevention Study II ([Gaudet et al., 2012](#)), BMI was inversely associated with death from cancer of the oral cavity, pharynx, and larynx combined. In contrast, in the pooled analysis, an incremental increase in BMI of 5 kg/m<sup>2</sup> was associated with a 36% increase in risk ([Gaudet et al., 2015](#)). Results from the Agricultural Health Study ([Andreotti et al., 2010](#)) were inconclusive.

The association between waist circumference and the risk of cancer of the oral cavity, pharynx, and larynx combined was examined in the large pooled analysis of 20 prospective studies, and no evidence of association was observed ([Gaudet et al., 2015](#)).

(v) *Cancer of the salivary glands*

There has been only one study of the association between BMI and incidence of salivary gland cancer ([Samanic et al., 2004](#)). In that study, being

obese was not associated with a higher incidence compared with being non-obese in either White men or Black men.

(vi) *Cancer of the head and neck or upper aerodigestive tract*

For head and neck cancer incidence overall, in the United States Veterans study there was a significantly lower risk for obese compared with non-obese Black men and White men, without adjustment for tobacco use or alcohol consumption ([Samanic et al., 2004](#)). Most other prospective studies found a weak inverse association or no association between BMI at baseline and incidence of head and neck cancer ([Wolk et al., 2001](#); [Gaudet et al., 2012](#); [Hashibe et al., 2013](#); [Etemadi et al., 2014](#)). When the pooled analysis of data from 20 prospective studies was stratified by smoking status, BMI was positively associated with risk in never-smokers but was inversely associated with risk in current smokers ([Gaudet et al., 2015](#)).

BMI was inversely associated with head and neck cancer mortality ( $P_{\text{trend}} = 3 \times 10^{-10}$ ) in the Cancer Prevention Study II in the USA ([Gaudet et al., 2012](#)), and in a smaller cohort study in Switzerland a weaker inverse association was found between BMI and death from cancer of the upper aerodigestive tract ([Meyer et al., 2015](#)).

In the only study that examined the association between BMI at younger ages and risk of head and neck cancer, no association was found with increased BMI at age 20 years or at age 50 years, or with percentage change in BMI from age 20 years or age 50 years to baseline ([Hashibe et al., 2013](#)).

Waist circumference was positively associated with risk of head and neck cancer incidence both in the NIH-AARP cohort study ([Etemadi et al., 2014](#)) and in the pooled analysis of 20 prospective studies, in which a 5 cm increase in waist circumference was associated with a 4% increase in risk (95% CI, 1.03–1.05) ([Gaudet et al., 2015](#)).

(b) *Case-control studies*

Since 2000, a total of seven independent case-control studies, conducted in Australia, China, Cuba, India, Europe, Sudan, and the USA, and one large multicentre case-control study (nine countries) have reported on the association of BMI with various combinations of cancers of the head and neck (Table 2.2.22b, web only, available at: <http://publications.iarc.fr/570>). In addition, [Gaudet et al. \(2010\)](#) and [Lubin et al. \(2010, 2011\)](#) performed pooled reanalyses of 15–17 case-control studies with stratification by smoking status, by alcohol consumption status, and by subsite (Table 2.2.22c, web only, available at: <http://publications.iarc.fr/570>).

In most studies, BMI was assessed on the basis of self-reported height and body weight, referring to either a recent period (mostly 1 or 2 years) before disease diagnosis or to a period in the more distant past (e.g. at age 30 years). All original studies adjusted for potential confounding by smoking or alcohol consumption, in addition to variable adjustments for other potential confounding factors.

Most of the studies found an inverse association of BMI with cancer risk. In several studies, compared with normal-weight individuals ( $18.5 \text{ kg/m}^2 \leq \text{BMI} < 25 \text{ kg/m}^2$ ), those who were overweight or obese had reduced risks of head and neck cancer ([Rajkumar et al., 2003](#); [Rodriguez et al., 2004](#); [Kreimer et al., 2006](#); [Peters et al., 2008](#); [Radoï et al., 2013](#); [Petrick et al., 2014](#) in African Americans only); being underweight ( $\text{BMI} < 18.5 \text{ kg/m}^2$ ) was associated with an approximately 2-fold increase in risk in two large-scale studies (French ICARE study, 689 cases and 3481 controls, [Radoï et al., 2013](#); United States CHANCE study, 1289 cases and 1361 controls, [Petrick et al., 2014](#)). In the one study that additionally reported recalled body weight at age 30 years ([Radoï et al., 2013](#)), this inverse association was also observed for past BMI.

Four case-control studies stratified the analyses by smoking status. In one early study in the USA, the inverse association was more pronounced in current or ever-smokers than in never-smokers ([Kabat et al., 1994](#)). In two more recent studies in the USA, a similar pattern was observed in African Americans but not in Whites ([Petrick et al., 2014](#)) and in both HPV-positive and HPV-negative individuals ([Tan et al., 2015](#)). In contrast, the IARC Multicenter Oral Cancer Study, which included a total of 1670 cases and 1732 controls from nine countries worldwide, found statistically significant inverse associations of BMI (country-specific tertiles) with risk of oral and oropharyngeal squamous cell carcinomas in both tobacco users and never-users, as well as in alcohol consumers and never-drinkers ([Kreimer et al., 2006](#)). Similarly, a pooled reanalysis of the data from 17 case-control studies, which included a total of 12 716 cases and 17 438 controls (INHANCE consortium; [Gaudet et al., 2010](#)) (see Table 2.2.22c, web only, available at: <http://publications.iarc.fr/570>), found inverse relationships of BMI with the risk of cancers of the oral cavity, pharynx, and larynx, in men and women combined, in ever-smokers (for  $\text{BMI} \geq 30 \text{ kg/m}^2$  vs  $18.5- < 25 \text{ kg/m}^2$ : OR, 0.38; 95% CI, 0.30–0.49) but not in never-smokers. Furthermore, the increase in risk in underweight ( $\text{BMI} < 18.5 \text{ kg/m}^2$ ) compared with normal-weight ( $18.5- < 25 \text{ kg/m}^2$ ) individuals was significant only in the smokers (OR, 2.13; 95% CI, 1.75–2.58) ([Gaudet et al., 2010](#)).

## References

- Andreotti G, Hou L, Beane Freeman LE, Mahajan R, Koutros S, Coble J, et al. (2010). Body mass index, agricultural pesticide use, and cancer incidence in the Agricultural Health Study cohort. *Cancer Causes Control*, 21(11):1759–75. doi:[10.1007/s10552-010-9603-9](https://doi.org/10.1007/s10552-010-9603-9) PMID:[20730623](https://pubmed.ncbi.nlm.nih.gov/20730623/)
- Bhaskaran K, Douglas I, Forbes H, dos-Santos-Silva I, Leon DA, Smeeth L (2014). Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5.24 million UK adults. *Lancet*, 384(9945):755–65. doi:[10.1016/S0140-6736\(14\)60892-8](https://doi.org/10.1016/S0140-6736(14)60892-8) PMID:[25129328](https://pubmed.ncbi.nlm.nih.gov/25129328/)
- Emadi A, O'Doherty MG, Freedman ND, Hollenbeck AR, Dawsey SM, Abnet CC (2014). A prospective cohort study of body size and risk of head and neck cancers in the NIH-AARP Diet and Health Study. *Cancer Epidemiol Biomarkers Prev*, 23(11):2422–9. doi:[10.1158/1055-9965.EPI-14-0709-T](https://doi.org/10.1158/1055-9965.EPI-14-0709-T) PMID:[25172872](https://pubmed.ncbi.nlm.nih.gov/25172872/)
- Gaudet MM, Kitahara CM, Newton CC, Bernstein L, Reynolds P, Weiderpass E, et al. (2015). Anthropometry and head and neck cancer: a pooled analysis of cohort data. *Int J Epidemiol*, 44(2):673–81. doi:[10.1093/ije/dyv059](https://doi.org/10.1093/ije/dyv059) PMID:[26050257](https://pubmed.ncbi.nlm.nih.gov/26050257/)
- Gaudet MM, Olshan AF, Chuang SC, Berthiller J, Zhang ZF, Lissowska J, et al. (2010). Body mass index and risk of head and neck cancer in a pooled analysis of case-control studies in the International Head and Neck Cancer Epidemiology (INHANCE) Consortium. *Int J Epidemiol*, 39(4):1091–102. doi:[10.1093/ije/dyp380](https://doi.org/10.1093/ije/dyp380) PMID:[20123951](https://pubmed.ncbi.nlm.nih.gov/20123951/)
- Gaudet MM, Patel AV, Sun J, Hildebrand JS, McCullough ML, Chen AY, et al. (2012). Prospective studies of body mass index with head and neck cancer incidence and mortality. *Cancer Epidemiol Biomarkers Prev*, 21(3):497–503. doi:[10.1158/1055-9965.EPI-11-0935](https://doi.org/10.1158/1055-9965.EPI-11-0935) PMID:[22219317](https://pubmed.ncbi.nlm.nih.gov/22219317/)
- Hashibe M, Hunt J, Wei M, Buys S, Gren L, Lee YC (2013). Tobacco, alcohol, body mass index, physical activity, and the risk of head and neck cancer in the Prostate, Lung, Colorectal, and Ovarian (PLCO) cohort. *Head Neck*, 35(7):914–22. doi:[10.1002/hed.23052](https://doi.org/10.1002/hed.23052) PMID:[22711227](https://pubmed.ncbi.nlm.nih.gov/22711227/)
- IARC (2002). Weight control and physical activity. Lyon, France: IARC Press (IARC Handbooks of Cancer Prevention, Vol. 6). Available from: <http://publications.iarc.fr/376>.
- Kabat GC, Chang CJ, Wynder EL (1994). The role of tobacco, alcohol use, and body mass index in oral and pharyngeal cancer. *Int J Epidemiol*, 23(6):1137–44. doi:[10.1093/ije/23.6.1137](https://doi.org/10.1093/ije/23.6.1137) PMID:[7721514](https://pubmed.ncbi.nlm.nih.gov/7721514/)
- Kreimer AR, Randi G, Herrero R, Castellsagué X, La Vecchia C, Franceschi S; IARC Multicenter Oral Cancer Study Group (2006). Diet and body mass, and oral and oropharyngeal squamous cell carcinomas: analysis from the IARC multinational case-control study. *Int J Cancer*, 118(9):2293–7. doi:[10.1002/ijc.21577](https://doi.org/10.1002/ijc.21577) PMID:[16331628](https://pubmed.ncbi.nlm.nih.gov/16331628/)
- Lubin JH, Gaudet MM, Olshan AF, Kelsey K, Boffetta P, Brennan P, et al. (2010). Body mass index, cigarette smoking, and alcohol consumption and cancers of the oral cavity, pharynx, and larynx: modeling odds ratios in pooled case-control data. *Am J Epidemiol*, 171(12):1250–61. doi:[10.1093/aje/kwq088](https://doi.org/10.1093/aje/kwq088) PMID:[20494999](https://pubmed.ncbi.nlm.nih.gov/20494999/)
- Lubin JH, Muscat J, Gaudet MM, Olshan AF, Curado MP, Dal Maso L, et al. (2011). An examination of male and female odds ratios by BMI, cigarette smoking, and alcohol consumption for cancers of the oral cavity, pharynx, and larynx in pooled data from 15 case-control studies. *Cancer Causes Control*, 22(9):1217–31. doi:[10.1007/s10552-011-9792-x](https://doi.org/10.1007/s10552-011-9792-x) PMID:[21744095](https://pubmed.ncbi.nlm.nih.gov/21744095/)
- Meyer J, Rohrmann S, Bopp M, Faeh D; Swiss National Cohort Study Group (2015). Impact of smoking and excess body weight on overall and site-specific cancer mortality risk. *Cancer Epidemiol Biomarkers Prev*, 24(10):1516–22. doi:[10.1158/1055-9965.EPI-15-0415](https://doi.org/10.1158/1055-9965.EPI-15-0415) PMID:[26215293](https://pubmed.ncbi.nlm.nih.gov/26215293/)
- Parr CL, Batty GD, Lam TH, Barzi F, Fang X, Ho SC, et al.; Asia-Pacific Cohort Studies Collaboration (2010). Body-mass index and cancer mortality in the Asia-Pacific Cohort Studies Collaboration: pooled analyses of 424,519 participants. *Lancet Oncol*, 11(8):741–52. doi:[10.1016/S1470-2045\(10\)70141-8](https://doi.org/10.1016/S1470-2045(10)70141-8) PMID:[20594911](https://pubmed.ncbi.nlm.nih.gov/20594911/)
- Peters ES, Luckett BG, Applebaum KM, Marsit CJ, McClean MD, Kelsey KT (2008). Dairy products, leanness, and head and neck squamous cell carcinoma. *Head Neck*, 30(9):1193–205. doi:[10.1002/hed.20846](https://doi.org/10.1002/hed.20846) PMID:[18642285](https://pubmed.ncbi.nlm.nih.gov/18642285/)
- Petrack JL, Gaudet MM, Weissler MC, Funkhouser WK, Olshan AF (2014). Body mass index and risk of head and neck cancer by race: the Carolina Head and Neck Cancer Epidemiology Study. *Ann Epidemiol*, 24(2):160–164.e1. doi:[10.1016/j.annepidem.2013.11.004](https://doi.org/10.1016/j.annepidem.2013.11.004) PMID:[24342030](https://pubmed.ncbi.nlm.nih.gov/24342030/)
- Radoi L, Paget-Bailly S, Cyr D, Papadopoulos A, Guida F, Tarnaud C, et al. (2013). Body mass index, body mass change, and risk of oral cavity cancer: results of a large population-based case-control study, the ICARE study. *Cancer Causes Control*, 24(7):1437–48. doi:[10.1007/s10552-013-0223-z](https://doi.org/10.1007/s10552-013-0223-z) PMID:[23677332](https://pubmed.ncbi.nlm.nih.gov/23677332/)
- Rajkumar T, Sridhar H, Balaram P, Vaccarella S, Gajalakshmi V, Nandakumar A, et al. (2003). Oral cancer in Southern India: the influence of body size, diet, infections and sexual practices. *Eur J Cancer Prev*, 12(2):135–43. doi:[10.1097/00008469-200304000-00007](https://doi.org/10.1097/00008469-200304000-00007) PMID:[12671537](https://pubmed.ncbi.nlm.nih.gov/12671537/)
- Rodriguez T, Altieri A, Chatenoud L, Gallus S, Bosetti C, Negri E, et al. (2004). Risk factors for oral and pharyngeal cancer in young adults. *Oral Oncol*,

- 40(2):207–13. doi:[10.1016/j.oraloncology.2003.08.014](https://doi.org/10.1016/j.oraloncology.2003.08.014)  
PMID:[14693246](https://pubmed.ncbi.nlm.nih.gov/14693246/)
- Samanic C, Gridley G, Chow WH, Lubin J, Hoover RN, Fraumeni JF Jr (2004). Obesity and cancer risk among white and black United States veterans. *Cancer Causes Control*, 15(1):35–43. doi:[10.1023/B:-CACO.0000016573.79453.ba](https://doi.org/10.1023/B:-CACO.0000016573.79453.ba) PMID:[14970733](https://pubmed.ncbi.nlm.nih.gov/14970733/)
- Tan X, Nelson HH, Langevin SM, McClean M, Marsit CJ, Waterboer T, et al. (2015). Obesity and head and neck cancer risk and survival by human papilloma-virus serology. *Cancer Causes Control*, 26(1):111–9. doi:[10.1007/s10552-014-0490-3](https://doi.org/10.1007/s10552-014-0490-3) PMID:[25398682](https://pubmed.ncbi.nlm.nih.gov/25398682/)
- Wolk A, Gridley G, Svensson M, Nyrén O, McLaughlin JK, Fraumeni JF, et al. (2001). A prospective study of obesity and cancer risk (Sweden). *Cancer Causes Control*, 12(1):13–21. doi:[10.1023/A:1008995217664](https://doi.org/10.1023/A:1008995217664) PMID:[11227921](https://pubmed.ncbi.nlm.nih.gov/11227921/)