

Table 2.2.8a Cohort studies of measures of body fatness and cancer of the lung

Reference Cohort Location Follow-up period	Total number of subjects Sex Incidence/mortality	Exposure categories	Exposed cases	Relative risk (95% CI)	Covariates	Comments	
Olson et al. (2002) Iowa Women's Health Study USA 1986–1998	38 006 Women Incidence	BMI				Age, smoking status (never, former, current), physical activity score, education level, beer consumption	No association with weight at age 18 yr, or with WC; association similar for current, former, and never- smokers
		< 22.8	168	1.0			
		22.9–25.0	126	0.92 (0.73–1.16)			
		25–27.4	87	0.76 (0.58–0.98)			
		27.4–30.7	81	0.69 (0.52–0.90)			
≥ 30.7	70	0.66 (0.50–0.89)					
		[<i>P</i> _{trend}]		< 0.001]			
Calle et al. (2003) Population-based cohort USA	404 576 Men Mortality	BMI				Age, education level, smoking, physical activity, alcohol consumption, marital status, aspirin, fat intake, vegetable intake	
		18.5–24.9	4885	1.00			
		25–29.9	4281	0.78 (0.75–0.82)			
		30–34.9	681	0.79 (0.73–0.86)			
		35–39.9	78	0.67 (0.54–0.84)			
≥ 40							
		[<i>P</i> _{trend}]		< 0.001]			
	495 477 Women Mortality	BMI				Age, education level, smoking, physical activity, alcohol consumption, marital status, aspirin, fat intake, vegetable intake, HRT use	
		18.5–24.9	3693	1.00			
		25–29.9	1278	0.88 (0.83–0.94)			
		30–34.9	305	0.82 (0.72–0.92)			
		35–39.9	54	0.66 (0.50–0.86)			
≥ 40	19	0.81 (0.52–1.28)					
		[<i>P</i> _{trend}]		< 0.001]			
Samanic et al. (2004) United States Veterans cohort USA 1969–1996	4 500 700 Men Incidence	Obesity				Age, calendar year	Obesity defined as discharge diagnosis of obesity: ICD-8: 277; ICD-9: 278.0
		Non-obese	White men:				
		Obese	78 205	1.00			
			4398	0.91 (0.88–0.94)			
		Black men:					
		Non-obese	18 884	1.00			
		Obese	568	0.60 (0.55–0.65)			

Table 2.2.8a Cohort studies of measures of body fatness and cancer of the lung

Reference Cohort Location Follow-up period	Total number of subjects Sex Incidence/mortality	Exposure categories	Exposed cases	Relative risk (95% CI)	Covariates	Comments
Kuriyama et al. (2005) Population-based cohort Japan 1984–1992	12 485 Men Incidence	BMI				Too few lung cancers in women in this cohort (<i>n</i> = 48)
		18.5–24.9	123	1.00		
		25.0–27.4	18	0.77 (0.47–1.27)		
		27.5–29.9	2	0.30 (0.07–1.20)		
		≥ 30	3	0.80 (0.20–3.26)		
		[<i>P</i> _{trend}]		[0.08]		
Rapp et al. (2005) Population-based cohort Austria 1985–2002	67 447 Men Incidence	BMI			Age, smoking, occupation	
		18.5–24.9	209	1.00		
		25–29.9	198	0.80 (0.66–0.97)		
	30–34.9	50	0.88 (0.65–1.20)			
	≥ 35	7	0.88 (0.41–1.86)			
	[<i>P</i> _{trend}]		[0.15]			
78 484 Women Incidence	BMI					
	18.5–24.9	64	1.00			
	25–29.9	45	1.00 (0.68–1.48)			
	30–34.9	17	0.87 (0.50–1.50)			
		[<i>P</i> _{trend}]		[0.67]		
Samanic et al. (2006) Swedish Construction Worker Cohort Sweden 1971–1999	362 552 Men Incidence	BMI			Age, year, smoking status	
		18.5–24.9	1638	1.00		
		25–29.9	1040	0.80 (0.74–0.87)		
	≥ 30	153	0.74 (0.63–0.88)			
	[<i>P</i> _{trend}]		[< 0.001]			
	BMI				Age, year	Never-smokers
18.5–24.9		54	1.00			
25–29.9		63	1.17 (0.81–1.70)			
≥ 30		10	1.11 (0.56–2.20)			
		[<i>P</i> _{trend}]		[0.47]		

Table 2.2.8a Cohort studies of measures of body fatness and cancer of the lung

Reference Cohort Location Follow-up period	Total number of subjects Sex Incidence/mortality	Exposure categories	Exposed cases	Relative risk (95% CI)	Covariates	Comments		
Fujino et al. (2007) Japan Collaborative Cohort Study for Evaluation of Cancer (JACC) Japan NR	Men Incidence	BMI			Age, study area	Weight at age 20 yr not associated with risk		
		< 18.5	85	1.35 (1.07–1.70)				
		18.5–24.9	672	1.00				
		25–29.9	89	0.66 (0.53–0.82)				
		≥ 30	3	0.38 (0.12–1.18)				
	Women Incidence	BMI					Age, study area	Weight at age 20 yr not associated with risk
		< 18.5	25	1.00 (0.40–2.49)				
		18.5–24.9	168	1.00				
25–29.9		53	1.07 (0.78–1.46)					
	≥ 30	2	0.38 (0.09–4.01)					
Reeves et al. (2007) Population-based cohort United Kingdom 1996–2001	1.2 million Women Incidence	BMI			Age, region, SES, reproductive history, smoking, alcohol consumption, physical activity, HRT use	Null association also for never-smokers		
		< 22.5	828	1.17 (1.09–1.25)				
		22.5–24.9	823	1.00				
		25.0–27.4	653	0.91 (0.85–0.99)				
		27.5–29.9	376	0.83 (0.75–0.91)				
		≥ 30	491	0.84 (0.77–0.92)				
Jee et al. (2008) Cohort from National Health Insurance Corporation Republic of Korea 1992–2006	770 556 Men Incidence	BMI			Age, smoking status and dose			
		< 20	1606	1.35 (1.24–1.47)				
		20–22.9	3655	1.13 (1.05–1.21)				
		23–24.9	2172	1.00				
		25.0–29.9	1560	0.92 (0.84–1.00)				
		≥ 30	73	1.29 (0.96–1.73)				
	[<i>P</i> _{trend}]		[< 0.0001]					

Table 2.2.8a Cohort studies of measures of body fatness and cancer of the lung

Reference Cohort Location Follow-up period	Total number of subjects Sex Incidence/mortality	Exposure categories	Exposed cases	Relative risk (95% CI)	Covariates	Comments
Jee et al. (2008) (cont.)	443 273 Women Incidence	BMI < 20 20–22.9 23–24.9 25.0–29.9 ≥ 30 [<i>P</i> _{trend}]	338 740 505 590 58	1.20 (1.00–1.44) 1.18 (1.02–1.37) 1.00 1.10 (0.94–1.29) 0.91 (0.63–1.33) [0.07]		
Kabat et al. (2008) Women’s Health Initiative USA 1998–2006	161 809 Women Incidence	BMI < 23.1 23.1–25.6 25.6–28.3 28.3–32.2 ≥ 32.2 [<i>P</i> _{trend}]	314 311 236 265 227	1.0 1.04 (0.88–1.23) 0.77 (0.64–0.93) 0.81 (0.68–0.98) 0.79 (0.65–0.96) [0.001]	Age, smoking, education level, ethnicity, HRT use, diet, physical activity, study	No association seen for WC, or weight at ages 18, 35, or 50 yr
		BMI < 23.1 23.1–25.6 25.6–28.3 28.3–32.2 ≥ 32.2 [<i>P</i> _{trend}]	197 total	1.0 1.24 (0.80–1.89) 0.94 (0.60–1.50) 0.73 (0.45–1.21) 0.83 (0.50–1.38) [0.15]	Age, education level, ethnicity, HRT use, diet, physical activity, height, study	Never-smokers
Andreotti et al. (2010) Agricultural workers USA 1993–2005	39 628 Men Incidence	BMI < 18.5 18.5–24.9 25–29.9 30–34.9 ≥ 35 [<i>P</i> _{trend}]	2 86 127 40 6	– 1.00 0.85 (0.60–1.20) 0.85 (0.54–1.35) 0.47 (0.15–1.49) [0.15]	Age, race, smoking, vegetable intake, exercise, cancer family history	

Table 2.2.8a Cohort studies of measures of body fatness and cancer of the lung

Reference Cohort Location Follow-up period	Total number of subjects Sex Incidence/mortality	Exposure categories	Exposed cases	Relative risk (95% CI)	Covariates	Comments	
Koh et al. (2010) Population-based cohort Singapore 1993–2006	63 257 Men and women Incidence	BMI				Age, sex	Association not seen in former smokers
		< 20	220	1.34 (0.98–1.83)			
		20–24	609	1.18 (0.88–1.59)			
		24–28	164	0.91 (0.66–1.25)			
		≥ 28	49	1.00			
		[<i>P</i> _{trend}]		[0.0004]			
		BMI					
		< 20	23	0.93 (0.55–1.56)			
20–24	50	1.01 (0.65–1.56)					
24–28	176	0.69 (0.42–1.13)					
≥ 28	38	1.00					
[<i>P</i> _{trend}]		[0.31]					
Parr et al. (2010) Pooled analysis of 39 cohort studies Asia, Australia, and New Zealand 1961–NR	424 519 Men and women Mortality	BMI	1478 total		Age, sex, tobacco use		
		12–18.4		1.11 (0.86–1.44)			
		18.5–24.9		1.0			
		25–29.9		0.68 (0.59–0.79)			
		≥ 30		0.83 (0.64–1.08)			
[<i>P</i> _{trend}]		[0.003]					
Leung et al. (2011) Elderly Health Service clients China 2000–2008	64 574 Men and women Mortality	BMI			Sex, smoking, education level, marital status, alcohol consumption	Similar association for never-smokers and ever-smokers	
		< 18.5	66	1.38 (1.05–1.79)			
		18.5–22.9	325	1.00			
		23–24.9	214	0.92 (0.77–1.09)			
		25–29.9	298	0.87 (0.75–1.02)			
		≥ 30	29	0.55 (0.38–0.80)			
[<i>P</i> _{trend}]							

Table 2.2.8a Cohort studies of measures of body fatness and cancer of the lung

Reference Cohort Location Follow-up period	Total number of subjects Sex Incidence/mortality	Exposure categories	Exposed cases	Relative risk (95% CI)	Covariates	Comments	
Smith et al. (2012) NIH-AARP cohort USA 1995–2006	271 238 Men Incidence	BMI				Age, race, smoking, education level, COPD, physical activity, alcohol consumption	
		< 18.5	38	1.15 (0.83–1.59)			
		18.5–22.5	650	1.12 (1.02–1.23)			
		22.5–24.9	1327	1.00			
		25–29.9	2905	0.92 (0.86–0.98)			
		30–34.9	953	0.87 (0.80–0.95)			
		≥ 35	220	0.81 (0.70–0.94)			
		[<i>P</i> _{trend}]	[< 0.001]				
		BMI				Never-smokers	
	< 18.5	0	–				
	18.5–22.5	16	1.04 (0.58–1.86)				
	22.5–24.9	39	1.00				
	25–29.9	77	1.01 (0.68–1.48)				
	30–34.9	29	1.38 (0.85–2.24)				
≥ 35	5	1.04 (0.41–2.67)					
	[<i>P</i> _{trend}]	[0.44]					
177 494 Women Incidence	BMI						
	< 18.5	81	1.23 (0.97–1.54)				
	18.5–22.5	852	1.15 (1.04–1.26)				
	22.5–24.9	775	1.00				
	25–29.9	1082	0.99 (0.90–1.08)				
	30–34.9	388	0.85 (0.75–0.96)				
	≥ 35	166	0.73 (0.61–0.87)				
	[<i>P</i> _{trend}]	[< 0.001]					

Table 2.2.8a Cohort studies of measures of body fatness and cancer of the lung

Reference Cohort Location Follow-up period	Total number of subjects Sex Incidence/mortality	Exposure categories	Exposed cases	Relative risk (95% CI)	Covariates	Comments
Smith et al. (2012) (cont.)		BMI < 18.5 18.5–22.5 22.5–24.9 25–29.9 30–34.9 ≥ 35 [<i>P</i> _{trend}]	5 54 54 78 41 17	1.81 (0.72–4.52) 1.17 (0.80–1.70) 1.00 1.00 (0.71–1.42) 1.19 (0.79–1.80) 1.00 (0.58–1.74) [< 0.85]	Age, race, education level, COPD, physical activity, alcohol consumption	Never-smokers
Bethea et al. (2013) Black Women’s Health Study USA 1995–2011	56 835 Women Incidence	BMI < 18.5 18.5–24.9 25–29.9 ≥ 30 [<i>P</i> _{trend}]	9 101 122 91	2.70 (1.36–5.42) 1.0 0.85 (0.65–1.11) 0.69 (0.52–0.93) [< 0.01]	Age, education level, physical activity, alcohol consumption, parity, age at first birth, family history, region	WC inversely associated with risk
Lam et al. (2013) NIH-AARP cohort USA 1995–2006	158 415 Men and women Incidence Never-smokers only	BMI < 18.5 18.5–24.9 25–29.9 ≥ 30 [<i>P</i> _{trend}]	23 194 192 123	1.57 (0.77–3.19) 1.00 1.00 (0.81–1.22) 1.21 (0.95–1.53) [0.21]	Age, education level, alcohol consumption, physical activity, energy intake	No association with BMI at ages 18, 35, and 50 yr, or with WC intake
Bhaskaran et al. (2014) Clinical Practice Research Datalink United Kingdom 1987–2012	5 243 978 Men and women Incidence	BMI, per 5 kg/m ² All Never-smokers only	19 339 2674	0.82 (0.81–0.84) 0.99 (0.93–1.05)	Age, sex, year, diabetes, alcohol consumption, SES; for combined analysis, also adjusted for smoking	

BMI, body mass index (in kg/m²); CI, confidence interval; COPD, chronic obstructive pulmonary disease; HRT, hormone replacement therapy; NIH-AARP, National Institutes of Health–AARP Diet and Health Study; NR, not reported; SES, socioeconomic status; WC, waist circumference; yr, year or years

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding
Goodman & Wilkens (1993) USA (Hawaii) 1979–1985	738 (men: 518, women: 230) 1626 (men: 1102, women: 524) Population	BMI 5 yr before diagnosis, quartiles:			
		Men:	NR		
		≤ 21.9		1.0	
		> 21.9–23.9		0.5	
		> 23.9–25.8		0.5	
		> 25.8		0.4	
		[<i>P</i> _{trend}]		< 0.01]	
		Women:	NR		
		≤ 20.2		1.0	
		> 20.2–22.6		0.8	
		> 22.6–25.5		1.0	
		> 25.5		0.6	
[<i>P</i> _{trend}]		[0.05]			
BMI at age 20–29 yr, quartiles					
Men:	NR				
≤ 20.2		1.0			
> 20.2–21.8		0.8			
> 21.8–23.6		0.8			
> 23.6		0.9			
[<i>P</i> _{trend}]		[0.93]			
Women:	NR				
≤ 19.1		1.0			
> 19.1–20.5		1.1			
> 20.5–22.3		1.0			
> 22.3		0.9			
[<i>P</i> _{trend}]		[0.45]			

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding	
Kabat (1996) USA (8 cities) 1981–1990	3607 9681 Hospital	BMI 5 yr before diagnosis				Age, education level, smoking (kg of tar inhaled), race, hospital, time period, alcohol consumption, history of chronic lung disease Additionally adjusted for years since stopping
		Current smokers:	NR			
		Men:				
		≥ 28		1.0		
		25–27.9		1.2 (1.0–1.5)		
		22–24.9		1.5 (1.2–1.9)		
		< 22		2.0 (1.5–2.5)		
		[<i>P</i> _{trend}]		[0.0001]		
		Women:	NR			
		≥ 28		1.0		
		25–27.9		1.2 (0.8–2.0)		
		22–24.9		1.7 (1.2–2.4)		
< 22		2.0 (1.4–2.7)				
[<i>P</i> _{trend}]		[0.0001]				
Former smokers:	NR					
Men:						
≥ 28		1.0				
25–27.9		1.1 (0.9–1.4)				
22–24.9		1.2 (1.0–1.5)				
< 22		1.3 (1.0–1.8)				
[<i>P</i> _{trend}]		[0.02]				
Women:	NR					
≥ 28		1.0				
25–27.9		1.4 (0.9–2.2)				
22–24.9		1.1 (0.7–1.7)				
< 22		1.5 (1.0–2.2)				
[<i>P</i> _{trend}]		[0.2]				

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding
Kabat (1996) (cont.)		Never-smokers: Men: ≥ 28 25–27.9 22–24.9 < 22 [<i>P</i> _{trend}]	NR	1.0 0.8 (0.4–1.5) 0.5 (0.3–1.1) 0.9 (0.4–2.1) [0.4]	
		Women: ≥ 28 25–27.9 22–24.9 < 22 [<i>P</i> _{trend}]	NR	1.0 1.9 (0.9–6.5) 2.4 (1.3–4.2) 2.9 (1.6–5.0) [0.0001]	
Rauscher et al. (2000) USA (New York State) 1982–1985	Pairs of never- smokers and former smokers 412 412 (188 pairs never- smokers; 224 former smokers) Population	BMI before illness (cases) or 1 yr ago (controls), octiles Overall: O1: ≤ 21.26 O2–O7: > 21.26–≤ 30.84 O8: > 30.84 Men: O1: ≤ 21.26 O2–O7: > 21.26–≤ 30.84 O8: > 30.84 Women: O1: ≤ 21.26 O2–O7: > 21.26–≤ 30.84 O8: > 30.84	41 305 66 11 164 31 30 141 35	1.0 1.5 (1.0–2.3) 2.6 (1.5–5.2) 1.0 2.1 (0.9–5.0) 3.4 (1.2–10) 1.0 1.2 (0.7–2.1) 2.9 (1.3–6.5)	Age, years of smoking, number of cigarettes smoked per day, education level

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding
Rauscher et al. (2000) (cont.)		Never-smokers: O1: ≤ 21.26 O2–O7: > 21.26–≤ 30.84 O8: > 30.84	23 137 28	1.0 1.5 (0.9–2.7) 2.4 (1.1–6.0)	
		Former smokers: O1: ≤ 21.26 O2–O7: > 21.26–≤ 30.84 O8: > 30.84	28 168 38	1.0 1.5 (0.8–3.0) 3.2 (1.3–8.1)	
Kubik et al. (2004) Czech Republic 1998–2002	Women: 435 non- smokers; 1710 smokers Population (proxy controls)	Current BMI < 22.9 23.0–25.9 26.0–28.9 > 28.9 [<i>P</i> _{trend}]	128 100 106 101	1.00 0.55 (0.39–0.78) 0.54 (0.39–0.77) 0.34 (0.24–0.49) [< 0.001]	Age, residence, education level, smoking
Pan et al. (2004) Canada (10 provinces), NECSS study 1994–1997	3338 (men: 1736, women: 1602) 5039 Population	BMI 2 yr before interview All: < 25 25– < 30 ≥ 30 [<i>P</i> _{trend}] Men: < 25 25– < 30 ≥ 30 [<i>P</i> _{trend}]	3338 1763	1.00 0.74 (0.65–0.84) 0.77 (0.66–0.91) [< 0.0001] 1.00 0.75 (0.64–0.88) 0.72 (0.57–0.90) [< 0.0005]	5-yr age group, province of residence, education level, pack-years of smoking, alcohol consumption, total energy intake, vegetable intake, dietary fibre intake, recreational physical activity For women, also menopausal status, number of live births, age at menarche, age at end of first pregnancy For all, also sex

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding	
Kanashiki et al. (2005) (cont.)		Women:				
		< 20.8	21	1.3 (0.7–2.7)		
		20.8–22.8	21	1.1 (0.6–2.2)		
		22.9–24.9	20	1.0		
		≥ 25.0	38	1.9 (1.0–3.6)		
		Women, never-smokers:				
		< 20.8	17	1.3 (0.6–2.7)		
		20.8–22.8	20	1.3 (0.6–2.6)		
		22.9–24.9	17	1.0		
		≥ 25.0	36	2.6 (1.2–4.3)		
		BMI at time of diagnosis				
		Men, current smokers:				
		< 20.8	20	0.9 (0.3–2.4)		
		20.8–22.8	13	0.6 (0.2–1.6)		
		22.9–24.9	13	1.0		
		≥ 25.0	6	0.3 (0.1–1.0)		
		Men, former smokers:				
		< 20.8	13	3.1 (0.1–1.0)		
		20.8–22.8	5	1.7 (0.5–6.1)		
		22.9–24.9	9	1.0		
≥ 25.0	12	2.3 (0.8–6.3)				
Men, never-smokers:						
< 20.8	1	1.5 (0.1–28)				
20.8–22.8	3	2.0 (0.2–25)				
22.9–24.9	2	1.0				
≥ 25.0	2	1.5 (0.1–19)				

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding
Kanashiki et al. (2005) (cont.)		Women, never-smokers: < 20.8 20.8–22.8 22.9–24.9 ≥ 25.0	23 14 31 30	0.8 (0.4–1.6) 0.4 (0.2–0.7) 1.0 0.9 (0.5–1.6)	
Brennan et al. (2009) Eastern Europe and Russian Federation (15 centres in 6 countries) 1998–2003	2250 3052 Hospital	BMI 2 yr before interview ≤ 25 26–30 31–35 36–40 ≥ 41 [<i>P</i> _{trend}]	1180 719 231 50 11	1.00 0.60 (0.52–0.69) 0.47 (0.39–0.58) 0.54 (0.36–0.81) 0.30 (0.14–0.66) [5 × 10 ⁻¹⁸]	Age, sex, cumulative tobacco consumption, years of alcohol consumption, country
Heck et al. (2009) USA (10 counties) 2005–2007	223 (men: 100, women: 123) 238 (men: 97, women: 141) Population	BMI 6 mo before interview 17.2–24.9 25–29.9 ≥ 30	89 72 62	1.00 0.80 (0.51–1.24) 0.65 (0.41–1.01)	Crude odds
Tarleton et al. (2012) USA (Los Angeles County) 1999–2004	611 (men: 303, women: 308) 1029 (men: 623, women: 417) Population (neighbourhood controls)	All: BMI at age 21 yr < 20.34 20.34– < 22.15 22.15– < 24.34 ≥ 24.34 [<i>P</i> _{trend}]	187 148 129 134	1.00 0.97 (0.69–1.35) 0.89 (0.63–1.27) 0.93 (0.64–1.34) [0.611]	Age, sex, ethnicity, tobacco smoking status, education level, energy intake

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding
Tarleton et al. (2012) (cont.)		BMI 1 yr before interview			
		< 18.5	18	1.31 (0.55–3.14)	
		18.5– < 25	263	1.00	
		25– < 30	214	0.87 (0.66–1.16)	
		≥ 30	114	0.58 (0.41–0.81)	
		[<i>P</i> _{trend}]		[0.001]	
		BMI change			
		< –5% (loss)	51	1.09 (0.60–1.98)	
		–5%– < +5%	113	1.00	
		5%– < 15%	132	0.76 (0.51–1.13)	
		15%– < 25%	113	0.72 (0.47–1.09)	
		25%– < 35%	67	0.53 (0.33–0.84)	
		> 35%	122	0.53 (0.35–0.80)	
		[<i>P</i> _{trend}]		[0.001]	
		BMI change, by tobacco smoking status			
		Never smokers:			
		< –5% (loss)	4	1.59 (0.36–7.02)	
		–5%– < +5%	10	1.00	
		5%– < 25%	54	1.45 (0.66–3.19)	
		≥ 25%	37	1.11 (0.49–2.54)	
		[<i>P</i> _{trend}]		[0.787]	
		Former smokers:			
		< –5% (loss)	30	1.60 (0.72–3.54)	
		–5%– < +5%	63	1.00	
		5%– < 25%	142	0.67 (0.41–1.08)	
		≥ 25%	120	0.54 (0.33–0.90)	
		[<i>P</i> _{trend}]		[0.017]	

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding
Tarleton et al. (2012) (cont.)		Current smokers			
		< -5% (loss)	17	0.81 (0.31–2.10)	
		-5%–< +5%	40	1.00	
		5%–< 25%	49	0.40 (0.21–0.78)	
		≥ 25%	32	0.28 (0.13–0.57)	
		[<i>P</i> _{trend}]		[0.001]	
Tarnaud et al. (2012) France, ICARE study 2001–2006	2625 (men: 2029, women: 596) 3381 (men: 2641, women: 740) Population	BMI 2 yr before interview			Age, area of residence, tobacco smoking consumption, education level, occupational exposure, previous chronic bronchitis, parental history of lung cancer
		All:			
		Men:			
		< 18.5	28	2.7 (1.2–6.2)	
		≥ 18.5–< 25	712	1.0	
		≥ 25–< 30	660	0.9 (0.7–1.1)	
		≥ 30–< 32.5	152	0.8 (0.6–1.1)	
		≥ 32.5	124	0.8 (0.6–1.0)	
		[<i>P</i> _{trend}]		[0.02]	
		Women:			
		< 18.5	36	1.5 (0.7–2.9)	
		≥ 18.5–< 25	288	1.0	
		≥ 25–< 30	108	0.9 (0.6–1.2)	
		≥ 30	63	0.8 (0.6–1.3)	
		[<i>P</i> _{trend}]		[0.20]	
		Current smokers:			
		Men:			
		< 18.5	24	4.1 (1.1–15.1)	
		≥ 18.5–< 25	502	1.0	
		≥ 25–< 30	317	0.6 (0.5–0.8)	
		≥ 30–< 32.5	66	0.7 (0.4–1.2)	
		≥ 32.5	47	0.5 (0.3–0.8)	
		[<i>P</i> _{trend}]		[< 0.001]	

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding	
Tarnaud et al. (2012) (cont.)	2625 (men: 2029, women: 596) 3381 (men: 2641, women: 740) Population	Women:				
		< 18.5	29	2.0 (0.6–6.5)		
		≥ 18.5– < 25	170	1.0		
		≥ 25– < 30	41	0.9 (0.4–2.1)		
		≥ 30	20	0.3 (0.1–0.7)		
		[<i>P</i> _{trend}]		[0.007]		
		Former smokers:				
		Men:				
		< 18.5	4	2.9 (0.6–14.5)		
		≥ 18.5– < 25	192	1.0		
		≥ 25– < 30	319	1.1 (0.8–1.4)		
		≥ 30– < 32.5	83	0.9 (0.6–1.3)		
		≥ 32.5	74	1.0 (0.7–1.5)		
		[<i>P</i> _{trend}]		[0.68]		
		Women:				
		< 18.5	3	0.5 (0.1–3.0)		
≥ 18.5– < 25	50	1.0				
≥ 25– < 30	25	0.7 (0.3–1.5)				
≥ 30	9	0.6 (0.2–1.7)				
[<i>P</i> _{trend}]		[0.42]				
Never-smokers:						
Men:						
< 18.5	–	–				
≥ 18.5– < 25	18	1.0				
≥ 25– < 30	24	1.4 (0.7–2.7)				
≥ 30– < 32.5	3	0.7 (0.2–2.7)				
≥ 32.5	3	1.3 (0.4–4.9)				
[<i>P</i> _{trend}]		[0.71]				

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding
Tarnaud et al. (2012) (cont.)	2625 (men: 2029, women: 596) 3381 (men: 2641, women: 740) Population	Women: < 18.5 ≥ 18.5– < 25 ≥ 25– < 30 ≥ 30 [<i>P</i> _{trend}]	4 68 42 34	1.5 (0.4–5.3) 1.0 1.0 (0.6–1.5) 1.2 (0.7–2.0) [0.75]	
		BMI at age 30 yr			
		All:			
		Men:			
		< 18.5	26	1.5 (0.8–3.0)	
		≥ 18.5– < 25	1090	1.0	
		≥ 25– < 30	401	1.1 (0.9–1.4)	
		≥ 30– < 32.5	32	0.6 (0.3–1.1)	
		≥ 32.5	9	0.3 (0.1–0.7)	
		[<i>P</i> _{trend}]		[0.07]	
		Women:			
		< 18.5	66	1.0 (0.7–1.6)	
		≥ 18.5– < 25	344	1.0	
		≥ 25– < 30	35	0.9 (0.6–1.5)	
		≥ 30	10	0.7 (0.3–1.7)	
		[<i>P</i> _{trend}]		[0.44]	
		Current smokers:			
		Men:			
		< 18.5	17	2.1 (0.6–7.1)	
		≥ 18.5– < 25	638	1.0	
		≥ 25– < 30	210	0.9 (0.7–1.3)	
		≥ 30– < 32.5	15	0.6 (0.2–1.7)	
		≥ 32.5	3	0.2 (0.5–0.9)	
		[<i>P</i> _{trend}]		[0.04]	

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding
Tarnaud et al. (2012) (cont.)	2625 (men: 2029, women: 596) 3381 (men: 2641, women: 740) Population	Women: < 18.5 ≥ 18.5– < 25 ≥ 25– < 30 ≥ 30 [<i>P</i> _{trend}]	50 176 10 2	1.4 (0.6–3.0) 1.0 0.6 (0.2–1.9) 0.1 (0.0–0.8) [0.014]	
		Former smokers: Men: < 18.5 ≥ 18.5– < 25 ≥ 25– < 30 ≥ 30– < 32.5 ≥ 32.5 [<i>P</i> _{trend}]	9 425 177 16 6	1.5 (0.6–14.5) 1.0 1.2 (0.8–1.4) 0.5 (0.6–1.3) [sic] 0.4 (0.7–1.5) [sic] [0.17]	
		Women: < 18.5 ≥ 18.5– < 25 ≥ 25– < 30 ≥ 30 [<i>P</i> _{trend}]	9 65 7 2	1.3 (0.4–4.4) 1.0 0.7 (0.2–2.3) 1.2 (0.1–11.1) [0.68]	
		Never-smokers: Men: < 18.5 ≥ 18.5– < 25 ≥ 25– < 30 ≥ 30– < 32.5 ≥ 32.5 [<i>P</i> _{trend}]	– 27 14 1 –	– 1.0 1.7 (0.7–2.7) 1.3 (0.2–2.7) – (0.4–4.9) [sic] [0.35]	

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding
Tarnaud et al. (2012) (cont.)		Women: < 18.5 ≥ 18.5– < 25 ≥ 25– < 30 ≥ 30 [<i>P</i> _{trend}]	8 103 18 6	0.6 (0.2–1.3) 1.0 1.0 (0.5–1.9) 1.1 (0.4–3.1) [0.39]	
El-Zein et al. (2013) Canada (greater Montreal area) 1996–2002	1076 (men: 628, women: 443) 1439 (men: 849, women: 582) Population	BMI 2 yr before diagnosis (cases) or interview (controls) All: < 18.5 18.5–24.9 25–29.9 ≥ 30 Never-smokers and light smokers: < 18.5 18.5–24.9 25–29.9 ≥ 30 Heavy smokers: < 18.5 18.5–24.9 25–29.9 ≥ 30 BMI at age 20 yr All: < 18.5 18.5–24.9 25–29.9 ≥ 30	63 521 346 141 7 60 47 20 56 461 299 121 114 799 145 13	2.30 (1.30–4.10) 1.00 0.83 (0.67–1.04) 0.96 (0.71–1.31) 2.28 (0.85–6.09) 1.00 0.70 (0.45–1.08) 1.01 (0.56–1.83) 2.33 (1.11–4.90) 1.00 0.89 (0.69–1.16) 0.93 (0.64–1.34) 0.69 (0.50–0.95) 1.00 1.18 (0.87–1.60) 0.58 (0.24–1.42)	Age, sex, education level, respondent status, ancestry, consumption of fruits and vegetables, occupational exposure, recreational physical activity, alcohol consumption, Comprehensive Smoking Index (CSI)

Table 2.2.8b Case-control studies of measures of body fatness and cancer of the lung

Reference Study location Period	Total number of cases Total number of controls Source of controls	Exposure categories	Exposed cases	Relative risk (95% CI)	Adjustment for confounding
El-Zein et al. (2013) (cont.)		Never-smokers and light smokers:			
		< 18.5	12	0.59 (0.30–1.14)	
		18.5–24.9	104	1.00	
		25–29.9	15	0.70 (0.37–1.32)	
		≥ 30	3	1.63 (0.40–6.56)	
		Heavy smokers:			
		< 18.5	102	0.69 (0.47–1.01)	
		18.5–24.9	695	1.00	
25–29.9	130	1.38 (0.96–1.98)			
≥ 30	10	0.37 (0.13–1.03)			

BMI, body mass index (in kg/m²); CI, confidence interval; mo, month or months; NECSS, National Enhanced Cancer Surveillance System; NR, not reported; SD, standard deviation; yr, year or years

Table 2.2.8c Mendelian randomization studies of measures of body fatness and cancer of the lung

Reference Study	Characteristics of study population	Sample size	Exposure (unit)	Outcome	Odds ratio (95% CI); <i>P</i> value (with each unit increase in exposure) of the association between the exposure and outcome(s)
Brennan et al. (2009)	Men and women from 15 centres in 6 countries in central and eastern Europe (Czech Republic, Hungary, Poland, Romania, Russian Federation, and Slovakia)	7067 (4015 cases and 3052 controls)	Increase of 1 kg/m ² in BMI	All tumours	0.85 (0.72–0.99); <i>P</i> = 0.04
				Adenocarcinoma	0.51 (0.33–0.82); <i>P</i> = 0.004
				Squamous cell carcinoma	0.72 (0.57–0.90); <i>P</i> = 0.01
				Never-smokers	0.57 (0.35–0.94); <i>P</i> = 0.03
				Former smokers	0.76 (0.49–1.17); <i>P</i> = 0.22
				Current smokers	0.89 (0.67–1.17); <i>P</i> = 0.40
Men and women from 6 studies of individuals of European ancestry	28 998 (12 160 cases and 16 838 controls)	Increase of 1 unit in genetically predicted adult BMI	Adult BMI:		
			All tumours	1.05 (1.02–1.09); <i>P</i> = 2.9 × 10 ⁻³	
			Adenocarcinoma	0.98 (0.93–1.10); <i>P</i> = 0.59	
			Squamous cell carcinoma	1.10 (1.04–1.16); <i>P</i> = 6.6 × 10 ⁻⁴	
			Childhood BMI:		
			All tumours	1.01 (0.85–1.20); <i>P</i> = 0.90	
Adenocarcinoma	0.90 (0.69–1.19); <i>P</i> = 0.47				
Squamous cell carcinoma	1.08 (0.82–1.43); <i>P</i> = 0.57				
			Increase of 1 SD (~0.073 kg/m ²) in genetically predicted childhood BMI		

BMI, body mass index (in kg/m²); CI, confidence interval; SD, standard deviation

References

- Andreotti G, Hou L, Beane Freeman LE, Mahajan R, Koutros S, Coble J, et al. (2010). Body mass index, agricultural pesticide use, and cancer incidence in the Agricultural Health Study cohort. *Cancer Causes Control*. 21(11):1759–75. <http://dx.doi.org/10.1007/s10552-010-9603-9> PMID:20730623
- Bethea TN, Rosenberg L, Charlot M, O'Connor GT, Adams-Campbell LL, Palmer JR (2013). Obesity in relation to lung cancer incidence in African American women. *Cancer Causes Control*. 24(9):1695–703. <http://dx.doi.org/10.1007/s10552-013-0245-6> PMID:23744044
- Bhaskaran K, Douglas I, Forbes H, dos-Santos-Silva I, Leon DA, Smeeth L (2014). Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5.24 million UK adults. *Lancet*. 384(9945):755–65. [http://dx.doi.org/10.1016/S0140-6736\(14\)60892-8](http://dx.doi.org/10.1016/S0140-6736(14)60892-8) PMID:25129328
- Brennan P, McKay J, Moore L, Zaridze D, Mukeria A, Szeszenia-Dabrowska N, et al. (2009). Obesity and cancer: Mendelian randomization approach utilizing the *FTO* genotype. *Int J Epidemiol*. 38(4):971–5. <http://dx.doi.org/10.1093/ije/dyp162> PMID:19542184
- Calle EE, Rodriguez C, Walker-Thurmond K, Thun MJ (2003). Overweight, obesity, and mortality from cancer in a prospectively studied cohort of U.S. adults. *N Engl J Med*. 348(17):1625–38. <http://dx.doi.org/10.1056/NEJMoa021423> PMID:12711737
- El-Zein M, Parent ME, Nicolau B, Koushik A, Siemiatycki J, Rousseau MC (2013). Body mass index, lifetime smoking intensity and lung cancer risk. *Int J Cancer*. 133(7):1721–31. <http://dx.doi.org/10.1002/ijc.28185> PMID:23553144
- Fujino Y; Japan Collaborative Cohort Study for Evaluation of Cancer (2007). Anthropometry, development history and mortality in the Japan Collaborative Cohort Study for Evaluation of Cancer (JACC). *Asian Pac J Cancer Prev*. 8(Suppl):105–12. PMID:18260709
- Gao C, Patel CJ, Michailidou K, Peters U, Gong J, Schildkraut J, et al.; Colorectal Transdisciplinary Study (CORECT); Discovery, Biology and Risk of Inherited Variants in Breast Cancer (DRIVE); Elucidating Loci Involved in Prostate Cancer Susceptibility (ELLIPSE); Follow-up of Ovarian Cancer Genetic Association and Interaction Studies (FOCI); and Transdisciplinary Research in Cancer of the Lung (TRICL) (2016). Mendelian randomization study of adiposity-related traits and risk of breast, ovarian, prostate, lung and colorectal cancer. *Int J Epidemiol*. 45(3):896–908. <http://dx.doi.org/10.1093/ije/dyw129> PMID:27427428
- Goodman MT, Wilkens LR (1993). Relation of body size and the risk of lung cancer. *Nutr Cancer*. 20(2):179–86. <http://dx.doi.org/10.1080/01635589309514284> PMID:8233983
- Heck JE, Andrew AS, Onega T, Rigas JR, Jackson BP, Karagas MR, et al. (2009). Lung cancer in a U.S. population with low to moderate arsenic exposure. *Environ Health Perspect*. 117(11):1718–23. PMID:20049123
- Jee SH, Yun JE, Park EJ, Cho ER, Park IS, Sull JW, et al. (2008). Body mass index and cancer risk in Korean men and women. *Int J Cancer*. 123(8):1892–6. <http://dx.doi.org/10.1002/ijc.23719> PMID:18651571
- Kabat GC (1996). Aspects of the epidemiology of lung cancer in smokers and nonsmokers in the United States. *Lung Cancer*. 15(1):1–20. [http://dx.doi.org/10.1016/0169-5002\(95\)00566-8](http://dx.doi.org/10.1016/0169-5002(95)00566-8) PMID:8865119

- Kabat GC, Kim M, Hunt JR, Chlebowski RT, Rohan TE (2008). Body mass index and waist circumference in relation to lung cancer risk in the Women's Health Initiative. *Am J Epidemiol.* 168(2):158–69. <http://dx.doi.org/10.1093/aje/kwn109> PMID:18483121
- Kanashiki M, Sairenchi T, Saito Y, Ishikawa H, Satoh H, Sekizawa K (2005). Body mass index and lung cancer: a case-control study of subjects participating in a mass-screening program. *Chest.* 128(3):1490–6. <http://dx.doi.org/10.1378/chest.128.3.1490> PMID:16162748
- Koh W-P, Yuan JM, Wang R, Lee HP, Yu MC (2010). Body mass index and smoking-related lung cancer risk in the Singapore Chinese Health Study. *Br J Cancer.* 102(3):610–4. <http://dx.doi.org/10.1038/sj.bjc.6605496> PMID:20010947
- Kubík AK, Zatloukal P, Tomásek L, Pauk N, Havel L, Krepela E, et al. (2004). Dietary habits and lung cancer risk among non-smoking women. *Eur J Cancer Prev.* 13(6):471–80. <http://dx.doi.org/10.1097/00008469-200412000-00002> PMID:15548939
- Kuriyama S, Tsubono Y, Hozawa A, Shimazu T, Suzuki Y, Koizumi Y, et al. (2005). Obesity and risk of cancer in Japan. *Int J Cancer.* 113(1):148–57. <http://dx.doi.org/10.1002/ijc.20529> PMID:15386435
- Lam TK, Moore SC, Brinton LA, Smith L, Hollenbeck AR, Gierach GL, et al. (2013). Anthropometric measures and physical activity and the risk of lung cancer in never-smokers: a prospective cohort study. *PLoS One.* 8(8):e70672. <http://dx.doi.org/10.1371/journal.pone.0070672> PMID:23940620
- Leung CC, Lam TH, Yew WW, Chan WM, Law WS, Tam CM (2011). Lower lung cancer mortality in obesity. *Int J Epidemiol.* 40(1):174–82. <http://dx.doi.org/10.1093/ije/dyq134> PMID:20709687
- Olson JE, Yang P, Schmitz K, Vierkant RA, Cerhan JR, Sellers TA (2002). Differential association of body mass index and fat distribution with three major histologic types of lung cancer: evidence from a cohort of older women. *Am J Epidemiol.* 156(7):606–15. <http://dx.doi.org/10.1093/aje/kwf084> PMID:12244029
- Pan SY, Johnson KC, Ugnat AM, Wen SW, Mao Y; Canadian Cancer Registries Epidemiology Research Group (2004). Association of obesity and cancer risk in Canada. *Am J Epidemiol.* 159(3):259–68. <http://dx.doi.org/10.1093/aje/kwh041> PMID:14742286
- Parr CL, Batty GD, Lam TH, Barzi F, Fang X, Ho SC, et al.; Asia-Pacific Cohort Studies Collaboration (2010). Body-mass index and cancer mortality in the Asia-Pacific Cohort Studies Collaboration: pooled analyses of 424,519 participants. *Lancet Oncol.* 11(8):741–52. [http://dx.doi.org/10.1016/S1470-2045\(10\)70141-8](http://dx.doi.org/10.1016/S1470-2045(10)70141-8) PMID:20594911
- Rapp K, Schroeder J, Klenk J, Stoehr S, Ulmer H, Concin H, et al. (2005). Obesity and incidence of cancer: a large cohort study of over 145,000 adults in Austria. *Br J Cancer.* 93(9):1062–7. <http://dx.doi.org/10.1038/sj.bjc.6602819> PMID:16234822
- Rauscher GH, Mayne ST, Janerich DT (2000). Relation between body mass index and lung cancer risk in men and women never and former smokers. *Am J Epidemiol.* 152(6):506–13. <http://dx.doi.org/10.1093/aje/152.6.506> PMID:10997540
- Reeves GK, Pirie K, Beral V, Green J, Spencer E, Bull D; Million Women Study Collaboration (2007). Cancer incidence and mortality in relation to body mass index in the Million Women Study: cohort study. *BMJ.* 335(7630):1134. <http://dx.doi.org/10.1136/bmj.39367.495995.AE> PMID:17986716
- Samanic C, Chow WH, Gridley G, Jarvholm B, Fraumeni JF Jr (2006). Relation of body mass index to cancer risk in 362,552 Swedish men. *Cancer Causes Control.* 17(7):901–9. <http://dx.doi.org/10.1007/s10552-006-0023-9> PMID:16841257

- Samanic C, Gridley G, Chow WH, Lubin J, Hoover RN, Fraumeni JF Jr (2004). Obesity and cancer risk among white and black United States veterans. *Cancer Causes Control*. 15(1):35–43. <http://dx.doi.org/10.1023/B:CACO.0000016573.79453.ba> PMID:14970733
- Smith L, Brinton LA, Spitz MR, Lam TK, Park Y, Hollenbeck AR, et al. (2012). Body mass index and risk of lung cancer among never, former, and current smokers. *J Natl Cancer Inst*. 104(10):778–89. <http://dx.doi.org/10.1093/jnci/djs179> PMID:22457475
- Tarleton HP, Park SL, Zhu WM, Lee YC, Hashibe M, Morgenstern H, et al. (2012). Body mass index change in adulthood and lung and upper aerodigestive tract cancers. *Int J Cancer*. 131(6):1407–16. <http://dx.doi.org/10.1002/ijc.27383> PMID:22131048
- Tarnaud C, Guida F, Papadopoulos A, Cénéé S, Cyr D, Schmaus A, et al. (2012). Body mass index and lung cancer risk: results from the ICARE study, a large, population-based case-control study. *Cancer Causes Control*. 23(7):1113–26. <http://dx.doi.org/10.1007/s10552-012-9980-3> PMID:22610667