

# Sulindac

## 1. Chemical and Physical Characteristics

### 1.1 Name

Chemical Abstracts Services Registry Number  
38194-50-2

Chemical Abstracts Primary Name  
Sulindac


### IUPAC Systematic Name

1*H*-Indene-3-acetic acid, 5-fluoro-2-methyl-1-  
{[4-(methylsulfinyl)phenyl]methylene}

### Synonyms

(*Z*)-5-Fluoro-2-methyl-1-[[4-(methylsulfinyl)-  
phenyl]methylene]-1*H*-indene-3-acetic acid;  
*cis*-5-fluoro-2-methyl-1-[*p*-(methylsulfinyl)-  
benzylidene]indene-3-acetic acid; MK-231

### 1.2 Structural and molecular formula and relative molecular mass



$C_{20}H_{17}FO_3S$

Relative molecular mass: 356.42

### 1.3 Physical and chemical properties

The data presented are for the pure substance and are taken from Budavari (1989) and Reynolds (1993), unless otherwise specified.

#### Description

Yellow, odourless crystals

#### Melting-point

182–185 °C

#### Solubility

A weak acid with a  $pK_a$  of 4.7 at 25 °C. Sparingly soluble in methanol and ethanol; slightly soluble in ethyl acetate; soluble in chloroform; practically insoluble in water at pH below 4.5. Solubility increases with rising pH to about 3.0 mg/ml at pH 7.

#### Spectroscopy data

Ultraviolet, infrared, nuclear magnetic resonance and mass spectra have been reported.

#### Stability

Stable in aqueous solutions of acids and bases. In a solid state, stable for at least three days in air at 100 °C.

#### Stereoisomers

Sulindac is the *Z* isomer of 1*H*-indeno-3-acetic acid, 5-fluoro-2-methyl-1-[[4-(methylsulfinyl)phenyl]methylene].

### 1.4 Technical products

#### Trade names

Aflodac, Algocetil, Apo-Sulin, Arthrobid, Arthrocin, Artribid, Citireuma, Clinoril, Clisundac, Imbaral, Lyndak, Novosundac, Reumofil, Reumyl, Sudac, Sulartrene, Sulen, Sulic, Sulindal, Sulinol, Sulreuma.

## 2. Occurrence, Production, Use, Analysis and Human Exposure

### 2.1 Occurrence

Sulindac is not known to occur as a natural product.

### 2.2 Production

Sulindac is a synthetic product; it is manufactured in several countries. Accepted standard procedures for the synthesis of sulindac are described by Shen and Winter (1977). Technical details regarding its current commercial production were not available to the Working Group.

## 2.3 Use

Sulindac was introduced in the 1970s (Shen & Winter, 1977). It has analgesic, anti-inflammatory and antipyretic properties and is used in musculoskeletal and joint disorders, such as ankylosing spondylitis, osteoarthritis and rheumatoid arthritis, and also in the short-term management of conditions such as bursitis and tendinitis and acute gouty arthritis. The usual dose by mouth is 100–200 mg twice daily, taken with food (Reynolds, 1993).

Sulindac is a pro-drug, a pharmacologically inactive precursor that is converted *in vivo* to an active drug by metabolism or other physiological processes. Its sulfide metabolite inhibits cyclo-oxygenases. Some of its clinical properties may be attributable to the sulfone metabolite (Singh *et al.*, 1994).

## 2.4 Analysis

Sulindac has been measured in human plasma by high-performance liquid chromatography (Shimek *et al.*, 1981; Swanson & Boppana, 1981; Stubbs *et al.*, 1987), by combined isotope dilution radioimmunoassay (Hare *et al.*, 1977) and by differential pulse polarographic analysis (Zamboni *et al.*, 1983). Fluorescence detection combined with reversed-phase high-performance liquid chromatography has also been used to determine sulindac and its sulfone and sulfide metabolites in human serum (Siluveru & Stewart, 1995).

## 2.5 Human exposure

Estimates of the prevalence of sulindac use vary, but it is limited. Jones and Tait (1995) reported that sulindac accounted for less than 7% of NSAID prescriptions in 1014 cases identified in general practice in the United Kingdom.

Sulindac accounts for a minor proportion of NSAID use in the USA, as seen in the Tennessee Medicaid Programme (Griffin *et al.*, 1991).

# 3. Metabolism, Kinetics and Genetic Variation

## 3.1 Human studies

### 3.1.1 Metabolism

Sulindac is an anti-inflammatory pro-drug which, after absorption, undergoes two major

biotransformations in humans: reversible reduction to the sulfide metabolite, the most potent inhibitor of prostaglandin production, and irreversible oxidation to the sulfone metabolite, which is inactive as an anti-inflammatory agent. Figure 1 shows the proposed metabolic scheme for sulindac. Approximately 88% of sulindac is absorbed in humans after oral administration of a 200-mg dose (Duggan *et al.*, 1977). In normal subjects and patients with surgical ileostomies, as much as 50% of the total sulfide was formed by gut bacteria (Strong *et al.*, 1985), probably from sulindac excreted in bile. It appears to undergo extensive reabsorption and enterohepatic recycling. Incubation of sulindac and its derivatives with over 200 strains of bacteria isolated from human faeces *in vitro* showed extensive reduction by both aerobes and anaerobes (Strong *et al.*, 1987).

### 3.1.2 Pharmacokinetics

After a single 200-mg oral dose of sulindac, the peak plasma concentration was 4.7 µg/ml after 1.6 h in one study (Strong *et al.*, 1985) and 5.4 µg/ml after 1 h in another (Duggan *et al.*, 1977). Sulindac binds tightly to human serum albumin (Russeva *et al.*, 1994) and has a mean half-life of 97 h (Strong *et al.*, 1985). The peak plasma concentration of the sulfide metabolite was about 2.7 mg/ml and was reached after 3.1 h. The half-life was 14 h in normal subjects and 2.6 h in patients with an ileostomy (Strong *et al.*, 1985). Sustained plasma levels of the sulfide metabolite are consistent with a prolonged anti-inflammatory action. The peak plasma concentration of the sulfone metabolite was about 1.5 mg/ml and was reached after 2.9 h. The half-life was 20 h in normal subjects and 5.4 h in patients with an ileostomy (Strong *et al.*, 1985).

The primary route of excretion in humans is via the urine, as both sulindac and its sulfone metabolite (free and as glucuronide conjugates). About 50% of an administered dose is excreted in the urine, the conjugated sulfone metabolite accounting for the major portion. No significant level of free or conjugated sulfide was detected in urine. About 25% is found in the faeces, primarily as the sulfone and sulfide

metabolites (Duggan *et al.*, 1977). The concentrations of these metabolites in the gut lumen and mucosa were not known to the Working Group.

Because sulindac is excreted in the urine primarily as biologically inactive forms, it may affect renal function to a lesser extent than other NSAIDs (Miller *et al.*, 1984); however, adverse renal effects have been reported (see section 7.1.1.(c)). Patients with end-stage renal failure had substantially lower total and free plasma concentrations of the active sulfide metabolite of sulindac (Ravis *et al.*, 1993). The apparent half-lives of sulindac and sulindac sulfide were similar in the two groups, but the half-life of the sulfone metabolite was longer in patients with renal failure.

Since many patients who take NSAIDs have rheumatoid arthritis and are also given methotrexate for their joint inflammation, a study was conducted to evaluate the potential interactions between methotrexate and sulindac. Sulindac had little effect on the disposition


of methotrexate but a minor effect on the 7-OH-methotrexate metabolite (Furst *et al.*, 1990).

Sulindac and its active sulfide metabolite undergo placental transfer; however, the reduction of sulindac to the metabolite is decreased in the human fetus, and the process seems to be independent of gestational age (Kramer *et al.*, 1995).

### 3.2 Experimental models

The half-life of sulindac in plasma was 10 h in Sprague-Dawley rats, 3 h in beagle dogs and 0.5 h in rhesus monkeys (Hucker *et al.*, 1973). Rats had by far the highest plasma concentration (44.0 µg/ml) 1 h after an oral dose of 10 mg/kg bw; the concentrations were 0.8 µg/ml in dogs at 1 h, not detectable in rhesus monkeys 1 h after a dose of 3 mg/kg bw and 1.3 µg/ml in humans 1 h after a 50-mg oral dose. Rats and dogs eliminated the drug almost exclusively in faeces, whereas urinary excretion is favoured in monkeys and humans. The tissue distribution in rats indicated that the drug was

Figure 1. Sulindac and its sulfide and sulfone metabolites



present at levels in the following order: plasma > liver > stomach > kidney > small intestine. The values increased after 4 h in stomach (37 µg/g), small intestine (17 µg/g) and large intestine (7.2 µg/g) but declined in all other tissues. In rats, 86% of the dose was recovered in bile 24 h after an intravenous dose of 10 mg/kg, whereas 53% was recovered after an equivalent oral dose. Bile collected from a dog given 10 mg/kg intravenously contained 93% of the dose. Administration of a large oral dose (100 mg/kg bw) to rats did not appreciably alter the excretion pattern. In rats, placental transfer is quite low. The concentrations of sulindac and the sulfide and sulfone metabolites found in rat milk were 10–20% of plasma levels.

Sulindac can be reduced to the sulfide by rat liver or kidney homogenates and by rat faecal contents (Lee & Renwick, 1995a,b).

### 3.3 Genetic variation

No data were available to the Working Group.

## 4. Cancer-preventive Effects

### 4.1 Human studies

#### 4.1.1 *Studies of adenomatous polyps in patients with familial adenomatous polyposis*

The relationship between familial adenomatous polyposis (FAP) and colorectal cancer is discussed in the General Remarks.

##### (a) *Non-randomized intervention studies*

Thirteen non-randomized case series comprising a total of 128 patients with FAP have been studied to examine the efficacy of sulindac in preventing or inhibiting adenomatous polyps (Waddell & Loughry, 1983; Gonzaga *et al.*, 1985; Waddell *et al.*, 1989; Charneau *et al.*, 1990; Friend, 1990; Rigau *et al.*, 1991; Tonelli & Valanzano, 1993; Winde *et al.*, 1993; Mäkelä & Laitinen, 1994; Spagnesi *et al.*, 1994; Cerdán *et al.*, 1995; Kadmon *et al.*, 1995; Winde *et al.*, 1995). Sulindac at doses of 100–400 mg daily, maintained for up to four years, results in a reduction in the number and size of colorectal polyps in comparison with those before treatment, regression of some adenomatous polyps on endoscopy and regression of some extra-

colonic desmoid tumours. (Desmoid tumours are histologically benign, extracolonic, connective tissue tumours that affect FAP patients.) When treatment with sulindac was stopped, the polyps reappeared; when treatment was resumed, the polyps regressed again (Charneau *et al.*, 1990; Labayle *et al.*, 1991; Rigau *et al.*, 1991; Mäkelä & Laitinen, 1994). In studies to establish optimal doses, in which sulindac was administered in rectal suppositories, sustained treatment with doses as low as 50 mg/day maintained polyp suppression in many patients. The maintenance of treatment with sulindac appeared to be more important for polyp inhibition than did the amount taken daily (Winde *et al.*, 1993, 1995, 1997). Carcinoma of the rectum was, however, reported in three patients with FAP during sulindac therapy in whom regression of polyps had been observed (Niv & Fraser, 1994; Thorson *et al.*, 1994; Lynch *et al.*, 1995).

##### (b) *Randomized trials*

Three small randomized clinical trials have confirmed that sulindac reduces the number and size of colorectal polyps in patients with FAP (Table 1).

Labayle *et al.* (1991) conducted a randomized, placebo-controlled, double-blind crossover study of 10 patients with FAP and rectal polyps, previously treated by colectomy and ileorectal anastomosis. One patient was excluded from treatment because of noncompliance. Nine patients received either sulindac, 300 mg/day (100 mg three times per day), or placebo during two four-month periods separated by one month. During sulindac treatment, the rectal polyps regressed completely in six patients and almost completely in three. During placebo treatment, the polyps increased in size in five patients, remained unchanged in two and decreased in two. The difference between the groups given sulindac and placebo was statistically significant ( $p < 0.01$ ).

Giardiello *et al.* (1993) conducted a randomized, double-blind, placebo-controlled trial of 22 patients with FAP, including 18 who had not undergone colectomy. The patients received either sulindac at 300 mg/day (150 mg twice a day) or placebo for nine months and were

evaluated by flexible sigmoidoscopy for the number and size of polyps every three months for one year. When sulindac treatment was stopped at nine months, the mean number of polyps had decreased to 44% of the original number ( $p = 0.014$ ) and the diameter of the polyps to 35% of the original value ( $p < 0.001$ ). No patient had complete resolution of polyps. Three months after treatment with sulindac was stopped, both the number and the size of the polyps in the sulindac-treated patients had increased, but the values remained significantly lower than the original ones.

Nugent *et al.* (1993) described a randomized, placebo-controlled trial of 24 patients with FAP, all of whom had duodenal polyps, and 14 with rectal polyps. Patients were randomized to sulindac at 400 mg/day (200 mg twice daily) or placebo for six months. Treatment was associated with a reduction in epithelial-cell proliferation, as measured by 5-bromo-2-deoxyuridine in the duodenum (median labelling index, 15.8 or 14.4%;  $p = 0.003$ ) and some regression of duodenal polyps ( $p = 0.12$ ). In the rectum, a larger reduction in cell proliferation (median labelling index, 8.5 vs. 7.4;  $p = 0.018$ ) and a greater reduction in rectal polyps were seen. Debinski *et al.* (1995) later clarified that this trial showed a significant reduction in small ( $< 2$  mm) duodenal polyps in the group given sulindac (9/11) versus the placebo group (4/12) but no reduction in

polyps  $\geq 3$  mm. [The Working Group noted that duodenal polyps in FAP patients may not be relevant to the situation of sporadic polyps and colon cancer, that cell proliferation in the duodenum may be less relevant to cancer than is proliferation in the colon or rectum and that the number of patients was not clearly stated for all end-points.]

### (c) Studies of biomarkers

Spagnesi *et al.* (1994) studied 20 FAP patients (six with ileorectal anastomoses and 14 with an intact colon) and examined three end-points: cell proliferation (by thymidine labelling), labelled cell distribution along the crypts and number of polyps. While administration of sulindac (100 mg twice a day for two months) induced a significant reduction in the number of polyps, there were no changes in the proliferation indices.

Winde *et al.* (1997) measured genetic biomarkers of epithelial proliferation in a non-randomized, dose-finding, intervention study of 38 patients with FAP. All of the patients had had a colectomy and ileoanastomosis. Twenty-eight were given sulindac by intrarectal suppository at an initial dose of 150 mg twice daily, which was reduced to a mean dose of 64 mg/day for up to four years. Ten control patients with FAP received no sulindac. The treated patients had a marked reduction in the number of rectal polyps (386 polyps in

**Table 1. Published randomized clinical trials of sulindac and adenomatous polyps**

Authors	Patients	Dosage	Design	Results
Labayle <i>et al.</i> (1991)	10 patients with FAP, colectomy and rectal polyps, France	300 mg/day or placebo, 4-month regimens, alternating	Randomized double-blind, cross-over	Polyps regressed completely in 6 patients, partly in 3 during sulindac treatment
Giardiello <i>et al.</i> (1993)	22 patients with FAP (18 without colectomy), USA	300 mg/day or placebo x 9 months	Randomized double-blind	Sulindac decreased number of polyps by 56% ( $p = 0.014$ ) and size by 65% ( $p < 0.001$ ), in comparison with baseline
Nugent <i>et al.</i> (1993); Debinski <i>et al.</i> (1995)	24 patients with FAP and duodenal polyps, USA	400 mg/day or placebo x 6 months	Randomized double-blind	Duodenal polyps $< 2$ mm regressed in 9 of 11 patients on sulindac vs 4 of 12 on placebo. No change in larger ( $\geq 3$ mm) polyps

28 people originally; no polyps in 24 people treated for 11 months or longer). Mutant *ras* oncogene expression became undetectable after sulindac treatment for  $\geq 6$  months. The proliferative index was significantly (20%) lower in patients treated with sulindac for one year than at baseline. A significant decrease in the frequency of wild and mutant *p53* was seen only when antibodies with broad specificity were used but not with those more specific to mutant *p53*. Immunohistochemical overexpression of the apoptosis-blocking protein BCL-2 was correlated inversely to mutant *p53* staining in adenomatous tissue.

#### 4.1.2 Randomized clinical trial of sporadic adenomatous polyps

Ladenheim *et al.* (1995) randomized 44 adults to either sulindac (150 mg twice a day for four months) or placebo to assess the effect on regression of sporadic polyps that were originally  $< 1$  cm. The patients were selected from 162 asymptomatic people aged  $\geq 50$ , who were screened by flexible sigmoidoscopy at three hospitals. People with adenomatous polyps  $> 1$  cm or contraindications to taking NSAIDs were excluded. Treatment of four of the 22 patients given sulindac was stopped because of side-effects (anaemia in two patients, heartburn in one) or complications (urogenital sepsis in one patient), thought to be unrelated to treatment. No significant difference in the number or the mean size of polyps was seen among the sulindac-treated patients. Five polyps in patients given sulindac had disappeared in comparison with three in the group given placebo; nine adenomas remained in the group given sulindac and 12 in controls. [The Working Group noted the low statistical power of the study.]

#### 4.1.3 Case studies of treatment for desmoid tumours

Regression of desmoid tumours during treatment with sulindac has been reported in some studies (Belliveau & Graham, 1984; Waddell & Kirsch, 1991; Kadmon *et al.*, 1995; Izes *et al.*, 1996) but not in all (Klein *et al.*, 1987).

## 4.2 Experimental models

### 4.2.1 Experimental animals

#### (a) Colon

*Short-term studies.* Groups of 14 male Fischer 344 rats, seven weeks of age, were fed AIN-76A diets containing sulindac at concentrations of 0, 200 or 400  $\mu\text{g/g}$  diet [0, 200 or 400 ppm]. After seven days of feeding sulindac, all rats were given weekly subcutaneous injections of 15 mg/kg bw azoxymethane for two weeks. Formation of aberrant colonic foci was determined eight weeks after the second weekly treatment. The numbers of foci per colon were 181 in controls, 118 in rats at 200 ppm (7.7% inhibition) and 81 in those at 400 ppm (37% inhibition;  $p < 0.05$ ) (Pereira *et al.*, 1994). In a similar study, dietary administration of sulindac at 320 ppm in the diet suppressed the number of foci per colon by 59% ( $p < 0.05$ ) (Reddy *et al.*, 1996).

*Long-term studies.* These studies are summarized in Table 2.

Two groups of 48 female Balb/c mice, six to eight weeks of age, were injected subcutaneously with 25 mg/kg bw 1,2-dimethylhydrazine dissolved in 0.4% ethylenediamine tetraacetic acid in normal saline, pH 6.5, once weekly with or without sulindac for up to 25 weeks. Sulindac [purity unspecified] was administered in drinking-water at a concentration [unspecified] such that each animal received about 5 mg/kg bw per day. In a second experiment, 48 female Balb/c mice were treated with 1,2-dimethylhydrazine for 17 weeks and then randomized to control (22 rats) or sulindac (23 rats). The control group received water. Those given sulindac received a solution in drinking-water for up to 11 weeks, as described above. Body weights were not given.

In the first study, which was designed to investigate the efficacy of sulindac administered simultaneously with 1,2-dimethylhydrazine, 38 animals in each group survived until the end of the study. Of the tumours observed macroscopically, 96% were adenomas and the rest were adenocarcinomas. Microadenomas developed in 92% of control animals and 47% of those given sulindac

( $p < 0.0001$ ), and colon tumours developed in 89% (34/38) of control animals and 42% (16/38) treated with sulindac (odds ratio, 12; 95% CI, 3.5–40). The overall colon tumour burden, the number of tumours per mouse and the number of tumours per tumour-bearing mouse were significantly reduced in the group given sulindac as compared with the control group ( $p < 0.022$ – $0.0001$ ). In the second study, in which sulindac was administered after the carcinogen, it did not inhibit colon carcinogenesis (numbers of animals with tumours, 17/18 and 18/18) (Moorghen *et al.*, 1988).

A group of 18 male Sprague-Dawley rats, weighing 400–500 g (age unspecified), was given 1,2-dimethylhydrazine dihydrochloride subcutaneously at a dose of 20 mg/kg bw once weekly (equivalent to 10 mg/kg bw 1,2-dimethylhydrazine) for 20 weeks. The site and diameter of each colon tumour were determined by laparotomy and colonoscopy. The rats were then randomized into two groups, receiving either 10 mg/kg bw sulindac (eight rats) or the vehicle, 0.5% methylcellulose (10 rats), twice daily by intragastric administration for four weeks. The animals given sulindac developed no additional tumours, whereas the controls had 13 additional colon tumours. Furthermore, the mean size of the 14 tumours in animals treated with sulindac was significantly smaller (9.3 mm) than that of the 39 tumours in the controls (57 mm;  $p = 0.026$ ). In this model system, predominantly carcinomas were induced (Skinner *et al.*, 1991).

Groups of 30 male Fischer 344 rats (age unspecified) were given subcutaneous injections of 15 mg/kg bw azoxymethane once weekly for two weeks. Two weeks after the second injection, 30 animals were fed AIN-76A diet containing 0.04% [400 ppm] sulindac for 31 weeks. These animals showed a 27% reduction in the number of colon tumours per rat in comparison with controls (Alberts *et al.*, 1995). [The Working Group noted that the actual numbers of colon tumours were not given and statistical significance was not reported.]

In a preliminary study, the MTD of sulindac in male Fischer 344 rats was shown to be about 400 ppm. In a preclinical efficacy study, groups of 36 male Fischer 344 rats, five weeks of age,

were fed AIN-76A diet containing 0, 160 or 320 ppm sulindac (purity, > 98%). At seven weeks of age, all animals were given two weekly subcutaneous injections of 15 mg/kg bw azoxymethane per week. Animals intended for studying the post-initiation effects of sulindac were given the compound in the diet at 320 ppm, starting 14 weeks after the second azoxymethane injection. Treatment was continued in all groups until termination of the study 52 weeks after azoxymethane treatment. All of the colonic tumours were evaluated histopathologically. No differences in body weight were seen among the groups. Inhibition of the incidence (percentage of animals with tumours) and multiplicity (tumours per animal) of invasive and non-invasive adenocarcinomas of the colon in rats during the initiation and post-initiation phases was seen to be dose-dependent ( $p < 0.001$ – $0.0001$ ). Furthermore, administration of sulindac post-initiation significantly suppressed the incidence ( $p < 0.0001$ ) and multiplicity ( $p < 0.0001$ ) of colon adenocarcinomas (see Table 2). Oral administration of sulindac also reduced the colon tumour volume by more than 52% ( $p < 0.01$ ) (Rao *et al.*, 1995).

*Genetically altered mice.* Sulindac has been tested in the Min/+ mouse model, which is described in the General Remarks. These studies are summarized in Table 2.

Groups of 10 female C57Bl/6J-Min/+ mice, five to six weeks of age, were fed AIN-76A diet and given sulindac at 160 mg/ml [160 ppm] in the drinking-water (estimated to provide 0.5 mg/animal per day). Control animals were given the diet alone. No differences in body weights or food intake were seen among the study groups. All animals were killed at 110 days of age. Multiple intestinal tumours were seen in all control animals, whereas only one mouse treated with sulindac had an adenoma of the colon. The control Min/+ mice had 12 intestinal tumours per mouse, and those given sulindac had 0.1 tumour/mouse ( $p < 0.001$ ). Ninety percent of the tumours were located in the ileum and jejunum (Boolbol *et al.*, 1996).

In another study, groups of 32–36 male and female Min/+ mice, 28 days of age, were given

**Table 2. Chemopreventive activity of sulindac against tumourigenesis in the colon and small intestine of experimental animals**

Site	Strain, species, sex and basal diet	Carcinogen, doses, route of administration	Sulindac dose, route and duration of administration	Tumour incidence (% animals with tumours)		Tumour multiplicity (tumours/animals)		Reference
				Control	Sulindac	Control	Sulindac	
Colon	Mice, Balb/c, female, standard chow	DMH, s.c., 25 mg/kg bw once weekly for 25 weeks	5 mg/kg bw, daily in drinking-water during and after carcinogen treatment	89	42	NR	NR	Moorghen <i>et al.</i> (1988)
					$p < 0.05$			
Colon	Mice, Balb/c, female, standard chow	DMH, s.c., 25 mg/kg bw, once weekly for 17 weeks	5 mg/kg bw, daily in drinking-water during and after carcinogen treatment	94	100	NR	NR	Moorghen <i>et al.</i> (1988)
					$p > 0.05$			
Colon	Rats, Sprague-Dawley, male, standard chow	DMH, s.c., 10 mg/kg bw, once weekly for 20 weeks	10 mg/kg bw, daily by gavage for 4 weeks after carcinogen treatment	100	100	39	14	Skinner <i>et al.</i> (1991)
					$p > 0.05$		$p < 0.012$	
Colon	Rats, F344, male, AIN-76A diet	AOM, 15 mg/kg bw, once weekly for 2 weeks	0.4 mg/g diet, daily two weeks after carcinogen treatment	Reduced to 39% of control		Reduced to 27% of control		Alberts <i>et al.</i> (1995)
Colon	Rats, F344, male, AIN-76A diet	AOM, 15 mg/kg bw, once weekly for 2 weeks	0.16 mg/g diet, daily one week before, during and after carcinogen treatment	81	42	1.5	0.68	Rao <i>et al.</i> (1995)
					$p < 0.01$		$p < 0.001$	
Colon	Rats, F344, male, AIN-76A diet	AOM, 15 mg/kg bw, once weekly for 2 weeks	0.16 mg/g diet, daily one week before, during and after carcinogen treatment	81	55	1.5	0.44	Rao <i>et al.</i> (1995)
					$p < 0.001$		$p < 0.001$	
Colon	Rats, F344, male, AIN-76A diet	AOM, 15 mg/kg bw, once weekly for 2 weeks	0.32 mg/g diet, beginning 14 weeks after carcinogen treatment	81	68	1.5	0.29	Rao <i>et al.</i> (1995)
					$p < 0.0001$		$p < 0.0001$	



**Table 2 (contd)**

Site	Strain, species, sex, and basal diet	Carcinogen, doses, route of administration	Sulindac dose, route and duration of administration	Tumour incidence (% animals with tumours)		Tumour multiplicity (tumours/animals)		Reference		
				Control	Sulindac	Control	Sulindac			
Small intestine	Mice, C57Bl/6J-Min, female, AIN-76A diet	No treatment	0.5 mg/mouse per day in drinking-water daily	100	NR	10	12	$p < 0.001$	0.1	Boolbol <i>et al.</i> (1996)
Small intestine	Mice, C57Bl/6J-Min, male, AIN-76A diet	No treatment	21 mg/l in drinking-water daily	100		100	49	$p < 0.05$	52	Beazer-Barclay <i>et al.</i> (1996)
Small intestine	Mice, C57Bl/6J-Min, male, AIN-76A diet	No treatment	84 mg/l in drinking-water daily	100		100	49	$p < 0.001$	30	Beazer-Barclay <i>et al.</i> (1996)
Small intestine	Mice, C57Bl/6J-Min, male, AIN-76A diet	No treatment	0.17 mg/g diet daily	100		100	33	$p < 0.0001$	12	Beazer-Barclay <i>et al.</i> (1996)
Small intestine	Mice, C57Bl/6J-Min, male, AIN-76A diet	No treatment	0.33 mg/g diet daily	100		100	33	$p < 0.00001$	9.8	Beazer-Barclay <i>et al.</i> (1996)

DMH, 1,2-dimethylhydrazine; s.c., subcutaneous; bw, body weight; NR, not reported; AOM, azoxymethane

sulindac in the drinking-water at 21 or 84 mg/litre [5 and 20 mg/kg bw per day] or in AIN-76A diet at 167 or 334 µg/g [40 and 80 mg/kg bw per day] for about 50 days. Administration of sulindac had no effect on body-weight gains, but the numbers of intestinal tumours were significantly suppressed by treatment with sulindac in drinking-water at 84 mg/litre or in the diet at either dose (Beazer-Barclay *et al.*, 1996).

(b) *Mammary gland*

These studies are summarized in Table 3.

Groups of 30 female Sprague-Dawley rats, 50 days old, were given a single intraperitoneal injection of *N*-methyl-*N*-nitrosourea at 50 mg/kg bw and fed AIN-76A diet containing 0 or 0.06% [600 ppm] sulindac [purity unspecified] beginning seven days after the carcinogen treatment. Sulindac had no effect on body-weight gain. After 24 weeks of treatment with sulindac, all mammary tumours were examined histopathologically. The tumour incidence was not significantly different in the controls (93%; 28/30) and those treated with sulindac (87%; 26/30) groups; however, the tumour multiplicity and burden were significantly inhibited in animals fed sulindac as compared with those fed the control diet ( $p < 0.05$ ). The mean numbers of mammary tumours ( $\pm$  SE) were  $4.2 \pm 0.5$  in the control group and  $2.7 \pm 0.4$  in the sulindac-treated group; the tumour burdens in grams  $\pm$  SE were  $15.8 \pm 2.5$  in the control group and  $5.7 \pm 1.8$  in the sulindac-treated group. Mammary tumour latency was longer in the rats given sulindac than in controls ( $p < 0.02$ ) (Thompson *et al.*, 1995).

Female Sprague-Dawley rats, 50 days of age, were injected with either 12.5 or 37.5 mg/kg bw *N*-methyl-*N*-nitrosourea to induce mammary tumours. Seven days later, the metabolite of sulindac, sulindac sulfone, was incorporated into AIN-76A diet at a concentration of 0.03 or 0.06% (w/w) [300 or 600 ppm]. Thirty rats were assigned to each dietary group treated with the high dose of carcinogen and 44 rats to each group treated with the low dose. Sulindac sulfone significantly reduced the incidence and the number of tumours per rat, irrespective of the dose of carcinogen injected. Its preven-

tive activity was comparable to that of sulindac. Tumour latency was prolonged significantly by sulindac sulfone, particularly at the low dose of carcinogen, when it was prolonged by more than eight weeks (Thompson *et al.*, 1997).

(c) *Oesophagus*

This study is summarized in Table 3. Groups of 15 male Fischer 344 rats [age unspecified] received three subcutaneous injections of 1 mg/kg bw *N*-nitrosomethylbenzylamine per week for five weeks. Two weeks before the start of this treatment, 15 rats received sulindac (purity, > 99%) in the diet at a concentration of 125 mg/g [125 ppm]; another group received sulindac one week after completion of carcinogen treatment, up to 25 weeks. There were no differences in mean body weights among the groups. The oesophageal tumours observed in this study were primarily exophytic, pedunculated lesions which proved to be squamous-cell papillomas. Neither the tumour incidence (100, 93 and 93%) nor the tumour multiplicity (2.9, 2.3 and 2.3 tumours/rat) was significantly inhibited when sulindac was administered before and during or after carcinogen treatment, respectively (Siglin *et al.*, 1995). [The Working Group noted that the tumour yield induced by *N*-nitrosomethylbenzylamine was high and that the dose of sulindac may have been insufficient to inhibit these oesophageal tumours.]

(d) *Urinary bladder*

This study is summarized in Table 3. Groups of 75 male B6D2F1 mice, five to six weeks of age, received sulindac in the diet a doses of 200 or 400 mg/g [200 and 400 ppm] one week before the first of eight weekly oral doses of 7.5 mg *N*-nitroso(4-hydroxybutyl)amine in ethanol:water. The study was terminated 24 weeks after administration of the carcinogen. The survival of animals at termination ranged from 93 to 100% in both groups, and administration of sulindac had a minimal effect on terminal body weights. Sulindac induced a dose-related decrease in the incidence of nitrosamine-induced urinary bladder cancer: control, 20/74 (27%); 200 ppm sulindac, 6/73 (8.2%); and 400 ppm sulindac, 2/75 (2.7%). There was no significant difference

**Table 3. Chemopreventive activity of sulindac against carcinogenesis in other organs of experimental animals**

Site	Strain, species, sex and basal diet	Carcinogen, doses, route of administration	Sulindac, dose, route and duration of administration	Tumour incidence (% animals with tumours)		Tumour multiplicity (tumours/animal)		Reference
				Control	Sulindac	Control	Sulindac	
Mammary gland	Rats, Sprague-Dawley, female, AIN076A diet	MNU, i.p., 50 mg/kg bw, single dose	600 ppm in diet for 24 weeks	93	87	4.2	2.7 <i>p</i> < 0.05	Thompson <i>et al.</i> (1995)
Mammary gland	Rats, Sprague-Dawley, female, AIN 76A diet	MNU, i.p., 12.5 mg/kg bw, single dose	Sulindac sulfone, 300 ppm diet 1 week after carcinogen treatment	100	39 <i>p</i> < 0.05	16	9 <i>p</i> < 0.05	Thompson <i>et al.</i> (1997)
			600 ppm in diet 1 week after carcinogen treatment	100	55 <i>p</i> < 0.05	16	8 <i>p</i> < 0.05	Thompson <i>et al.</i> (1997)
Mammary gland	Rats, Sprague-Dawley, female, AIN 76A diet	MNU, i.p., 37.5 mg/kg bw, single dose	300 ppm diet, 1 week after carcinogen treatment	100	100	126	103	Thompson <i>et al.</i> (1997)
			600 ppm in diet, 1 week after carcinogen treatment	100	100	126	68 <i>p</i> < 0.005	
Oesophagus	Rats, F344, male, AIN-76A diet	NMBzA, s.c., 1.0 mg/kg bw, 3 doses/week for 5 weeks	125 ppm diet before, during and after carcinogen for 25 weeks	100	93	2.9	2.3	Siglin <i>et al.</i> (1995)
Urinary bladder	Mice, B6D2F <sub>1</sub> , AIN-76A diet	OH-BBN, oral, 7.5 mg/week for 8 weeks	200 ppm diet for 24 weeks	27	8.2 <i>p</i> < 0.01	NR	NR	Rao <i>et al.</i> (1996)
			400 ppm diet for 24 weeks	27	2.7 <i>p</i> < 0.01	NR	NR	
Lung	Mice, A/J AIN-76A diet	NNK, 9.1 mg/mouse, in drinking-water for 7 weeks	130 ppm diet for 2 weeks before carcinogen and until termination at 16 weeks	100	92	15.7	7.4 <i>p</i> < 0.01	Castonguay <i>et al.</i> (1991); Pepin <i>et al.</i> (1992)
Lung	Mice, A/J, AIN-76A diet	NNK, 9.1 mg/animal in drinking-water for 7 weeks	-2 → 23 weeks	100	100	8.44	4.12 <i>p</i> < 0.001	Jalbert & Castonguay (1992)
			-2 → 7 weeks	100	100	8.44	6.04 <i>p</i> < 0.05	
			+7 → 23 weeks	100	100	8.44	7.41 <i>p</i> > 0.05	

MNU, *N*-methyl-*N*-nitrosourea; i.p., intraperitoneal; bw, body weight; NMBzA, *N*-nitrosomethylbenzylamine; s.c., subcutaneously; OH-BBN, *N*-nitrosobutyl(4-hydroxybutyl)amine; NR, not reported; NNK, 4-*N*-nitrosomethylamino)-1-(3-pyridyl)-1-butanone

in the histopathology of the lesions, including the degree of malignancy (Rao *et al.*, 1996).

#### (e) Lung

These studies are summarized in Table 3.

Groups of 23–28 female A/J mice, six to seven weeks of age, were fed sulindac (purity, 99%) at 0 or 130 mg/g diet [130 ppm]. Two weeks later, 4-(*N*-nitrosomethylamino)-1-(3-pyridyl)-1-butanone (NNK) (purity, 98.5%), which is present in tobacco smoke, was administered in the drinking-water for seven weeks (total dose, 9.0–9.2 mg/mouse). All animals were necropsied 16 weeks after the end of carcinogen treatment. Dietary administration of sulindac had no effect on body-weight gain or food consumption. The incidence of lung adenomas in the carcinogen-treated group was 90%, and the multiplicity was 15.7 tumours per mouse. Administration of sulindac reduced the tumour multiplicity to 7.4 tumours per mouse (53% inhibition;  $p < 0.001$ ), without affecting tumour incidence (Castonguay *et al.*, 1991; Pepin *et al.*, 1992).

In a further study, NNK was administered by the same protocol as described above, but three groups of mice were given sulindac (123 ppm) on weeks –2 to +23, –2 to +7 or +7 to +23. Treatment with NNK alone induced 8.4 lung adenomas per mouse; the treatments with sulindac reduced the multiplicity ( $\pm$  SE) of lung adenomas to  $4.12 \pm 0.44$  ( $p < 0.001$ ),  $6.04 \pm 0.43$  ( $p < 0.05$ ) and  $7.41 \pm 0.82$  ( $p > 0.05$ ), respectively. The authors concluded that the efficacy of sulindac given during and after initiation was additive (Jalbert & Castonguay, 1992).

#### 4.2.2 *In-vitro* models

No data were available to the Working Group.

#### 4.3 Mechanisms of chemoprevention

Sulindac was the first NSAID reported to be useful for the treatment of FAP (Waddell & Loughry, 1983). Administration of sulindac to affected individuals induced polyp regression, which was reversed on cessation of treatment. These findings have been confirmed in several subsequent studies (Belliveau & Graham, 1984;

Waddell *et al.*, 1989; Labayle *et al.*, 1991; Rigau *et al.*, 1991; Giardiello *et al.*, 1993). The efficacy of sulindac may be related to its unique pharmacokinetic properties, which lead to high concentrations of its active metabolite, sulindac sulfide, in the colonic lumen (Hucker *et al.*, 1973; Duggan *et al.*, 1977; Shen & Winter, 1977).

#### 4.3.1 Effects on cell proliferation and apoptosis

Numerous studies of the human colon cancer line HT-29 have shown that sulindac sulfide and sulfone can inhibit cell growth and stimulate apoptosis (Piazza *et al.*, 1995; Shiff *et al.*, 1995; Hanif *et al.*, 1996; Shiff *et al.*, 1996). In these studies, the cultured HT-29 cells were exposed to various concentrations of the sulindac metabolites, and several indices of growth regulation and/or apoptosis were evaluated. Both compounds inhibited cell growth and stimulated apoptosis, but in many of the studies only at concentrations in the range 0.5–1.2 mmol/litre. Therefore, the biological significance of these findings is uncertain.

Sulindac sulfone, when administered at pharmacological doses rather than the doses that result from the metabolism of sulindac, has no effect on cyclooxygenase (COX)<sup>1</sup> activity but inhibits cell proliferation and stimulates apoptosis (Thompson *et al.*, 1995), indicating that these effects are not related to COX activity (Shiff *et al.*, 1995; Hanif *et al.*, 1996; Schnitzler *et al.*, 1996). Another non-prostaglandin-related effect of sulindac sulfide and sulfone is to reduce the levels of cell-cycle regulatory proteins such as cdc2 (Shiff *et al.*, 1996), which may be related to the ability of these metabolites to inhibit cell proliferation.

A study of another colon cancer cell line, LIM 1215, with very sensitive polymerase chain reaction assay techniques, demonstrated that both sulindac sulfide and sulfone at doses of about 10 mmol/litre can induce the expression of APC mRNA (Schnitzler *et al.*, 1996). Since the APC gene has been shown to play a key role in colorectal carcinogenesis in both humans and

<sup>1</sup> Used as a synonym for prostaglandin endoperoxide synthase (PGH synthase)

animal models, this could have some significance; however, the investigators did not determine whether the increase in *APC* mRNA resulted in an increase in the amount of the functional APC protein. A question raised by this study is that, if the *APC* gene has already been inactivated by a mutation leading to premature chain termination, which occurs in 50% of spontaneous human colorectal adenocarcinomas, what benefit would there be in increasing the mRNA levels of this detective tumour suppressor gene?

Some NSAIDs cause apoptosis when applied to *v-src*-transformed chicken embryo fibroblasts (Lu *et al.*, 1995); however, sulindac had no effect on morphological inhibition of transformation. This study was the first to indicate that cells which overexpress *COX-2* may be somewhat resistant to programmed cell death, and that this can be reversed by addition of an NSAID.

Few nontransformed intestinal epithelial cell lines are available for in-vitro studies. Several groups have evaluated the mechanisms of growth control of a rat intestinal epithelial-1 (RIE-1) cell line in culture (DuBois *et al.*, 1995). Sulindac sulfide can inhibit mitogenesis of these cells (DuBois *et al.*, 1994a). Additionally, the cells express the inducible *COX-2* enzyme after treatment with cytokines and growth factors (DuBois *et al.*, 1994b), and the increased expression of *COX-2* in these cells makes them resistant to apoptosis (Tsuji & DuBois, 1995). Treatment of the cells with relatively low doses of sulindac sulfide (5–10 mmol/litre), however, reversed these phenotypic changes, indicating that a potential mechanism for the chemopreventive effect of sulindac may be its ability to inhibit *COX-2*.

#### **4.3.2 Effects on oncogene expression**

See the General Remarks for a discussion of the role of the *ras* oncogene in colon cancer.

Groups of six male Fischer 344 rats were fed a diet containing 320 ppm sulindac and were given azoxymethane dissolved in normal saline subcutaneously at a dose of 15 mg/kg bw per week for two weeks. Vehicle control groups received an equal volume of normal saline. The animals were sacrificed 52 weeks after

treatment and their colonic mucosa and tumours were analysed for mutations in codons 12 and 13 of *K-ras* and for expression of *ras p21*. Azoxymethane-induced G to A transitions were observed at the second nucleotide of codon 12 of *K-ras*, in which the amino acid aspartic acid replaced wild-type glycine. Sulindac not only suppressed the selective amplification of initiated cells with azoxymethane-induced mutated *K-ras* codon 12, but significantly inhibited the azoxymethane-induced expression of total and mutant *ras p21* (Singh & Reddy, 1995). The important finding of this study is that sulindac, which has been reported to induce polyp regression in patients with FAP and to inhibit carcinogen-induced colon carcinogenesis in experimental animals, significantly suppressed *ras* activation at both the DNA and protein level.

## **5. Other Beneficial Effects**

No data were available to the Working Group.

## **6. Carcinogenicity**

### **6.1 Humans**

The Working Group was not aware of any epidemiological studies of the carcinogenicity of sulindac or its metabolites

### **6.2 Experimental animals**

No study of adequate duration to evaluate carcinogenicity was available to the Working Group.

## **7. Other Toxic Effects**

### **7.1 Adverse effects**

#### **7.1.1 Humans**

See also the General Remarks and the chapter on sulindac. Relatively few studies refer specifically to the toxicity of sulindac. In many studies of the possible toxicity of NSAIDs, distinction was made only between aspirin and NSAIDs of other types. In common with other non-aspirin NSAIDs, sulindac is used at anti-inflammatory doses and inhibits *COX-1* and *COX-2*; therefore, data on the toxicity of non-aspirin NSAIDs are relevant to the assessment of the toxicity of sulindac.

*(a) Gastrointestinal tract toxicity*

NSAIDs can damage the oesophagus and can exacerbate gastro-oesophageal reflux disease (Heller *et al.*, 1982; Kikendale, 1991). Gastrointestinal bleeding and ulceration are potentially serious side-effects of all NSAIDs, as discussed in the chapter on aspirin. In the meta-analysis of Henry *et al.* (1996), described in the General Remarks, the risks for serious complications in the upper gastrointestinal tract after use of sulindac were similar to those of most other NSAIDs. In a ranking analysis, sulindac ranked 7 (1 being the most and 12 the least toxic) of 12 NSAIDs analysed. The pooled estimated RRs for NSAID users in comparison with ibuprofen users were between 1.6 and 2.7. Given the baseline risk for ulcer disease, which ranges from 0.5 per 1000 per year in young adults (Garcia Rodriguez *et al.*, 1992) to 4 per 1000 in older adults (Smalley *et al.*, 1995), the rates of serious ulcer complications among sulindac users can be estimated to be about 2 per 1000 in young adults and 15 per 1000 in people over the age of 65.

NSAIDs other than aspirin have also been associated with other deleterious effects on the small intestine, including inflammation resulting in blood and protein loss (Bjarnasson *et al.*, 1987), stricture (Matsushashi *et al.*, 1992), perforation and diarrhoea (Kwo & Tremaine, 1995). Large-bowel perforation and haemorrhage are also associated with use of non-aspirin NSAIDs (Langman *et al.*, 1985).

*(b) Hepatotoxicity*

Acute hepatotoxicity due to NSAIDs other than aspirin is an uncommon complication (Friis & Andreasen, 1992; Tarazi *et al.*, 1993; Rodriguez *et al.*, 1994). The incidence of acute liver injury was 3.7 per 100 000 NSAID users, or 1.1 per 100 000 NSAID prescriptions, among 625 307 persons who were prescribed any of 12 NSAIDs by a defined group of general practitioners in England between 1987 and 1991 (Rodriguez *et al.*, 1994).

*(c) Nephrotoxicity*

Acute renal toxicity due to NSAIDs can result from allergic interstitial nephritis, concurrent exposure to phenacetin, urate obstruction or

haemodynamic alterations in patients with underlying kidney disease (Murray & Brater, 1993; Murray *et al.*, 1995).

In a study of patients with chronic glomerular disease who were treated with therapeutic doses of sulindac, no effect was found on renal blood flow, glomerular filtration rate or urinary excretion of prostaglandins (Eriksson *et al.*, 1991); however, in a study of healthy volunteers and patients with liver disease (Laffi *et al.*, 1986), sulindac was found to blunt the renal responses to intravenous furosemide (Patrono, 1986).

Many experimental studies have shown that a large proportion of patients with conditions such as congestive heart failure, dehydration and cirrhosis, which result in a dependence on prostaglandins to maintain renal perfusion, suffer a decline in renal function when exposed to specific NSAIDs (Murray & Brater, 1990; Whelton *et al.*, 1990). There is some debate over whether all NSAIDs have a similar deleterious effect and, specifically, whether sulindac is less likely to cause a deterioration in renal function (Whelton & Hamilton, 1991). It is clear, however, that all such drugs, including sulindac, can both decrease renal prostaglandin production and cause a deterioration in renal function under conditions of decreased effective circulating volume (Brater *et al.*, 1985, 1986; Kleinknecht *et al.*, 1986; Stillman & Schlesinger, 1990).

The decline in renal function is usually reversible with discontinuation of the drugs. The frequency of the effect varies with the population studied, but may reach 13% in frail, elderly patients in nursing homes (Gurwitz *et al.*, 1990) and is much less frequent in healthier populations. In the general population, hospitalizations for acute renal failure are rare. Use of non-aspirin NSAIDs increases the risk by about fourfold, from a rate among non-users of about 2 per 100 000 yearly. The increase in risk is dose-dependent and highest during the first month of use (Perez-Gutthann *et al.*, 1996).

The data on NSAIDs and chronic renal failure are sparse. Typical analgesic nephropathy is widely accepted to be caused by phenacetin (Dubach *et al.*, 1991), but there is

also evidence that acetaminophen (Sandler *et al.*, 1989; Pernerger *et al.*, 1994) and NSAIDs (Adams *et al.*, 1986; Pernerger *et al.*, 1994) produce a similar type of kidney damage and/or other types of chronic renal failure.

#### (d) Blood pressure

Non-aspirin NSAIDs as a group interfere with the efficacy of antihypertensive drugs (Wong *et al.*, 1986; Radack & Deck, 1987; Chrischilles & Wallace, 1993; Johnson *et al.*, 1994), raise blood pressure in hypertensive subjects (Radack & Deck, 1987; Pope *et al.*, 1993) and may result in the initiation of antihypersensitive treatment in older persons (Gurwitz *et al.*, 1994).

In a meta-analysis of 50 randomized, placebo-controlled trials, use of NSAIDs for periods of weeks raised supine mean blood pressure by 5 mm Hg (range 1.2–8.7) (Johnson *et al.*, 1994). All NSAIDs appeared to have this effect, but the most marked increases in blood pressure were observed with piroxicam, indomethacin and ibuprofen; however, the numbers were too small to demonstrate statistically significant differences.

#### (e) Reproductive and developmental effects

The reported adverse effects of NSAID treatment during pregnancy and labour include: (i) prolongation of pregnancy and labour, (ii) increased maternal blood loss associated with delivery and (iii) anaemia. Use of sulindac during the third trimester of pregnancy is contraindicated because of potential premature closure of the ductus arteriosus, pulmonary hypertension and haemostatic abnormalities causing bleeding. Reductions in fetal urine output and in the volume of amniotic fluid are other possible adverse effects of NSAID therapy (Ostensen, 1994). At least one case report has indicated that ingestion of sulindac during pregnancy can lead to toxic epidermal necrolysis in newborns (Roupe *et al.*, 1986).

#### 7.1.2 Experimental animals

The values for the short-term  $TD_{50}$  (a comparative measure of toxicity) for intestinal ulceration and perforation in rats are 27 and 71 mg/kg bw (0.08 and 0.2 mmol/kg bw), respectively (Shen & Winter, 1977). Rats

given split daily doses of 0.28 mmol/kg bw per day sulindac intragastrically for four days (17.5 times the maximum human daily dose) developed medium and severe gastrointestinal ulceration with some evidence of perforation (Venuti *et al.*, 1989). In 90-day studies, rats given a dose of 40 mg/kg bw per day had ulcerative enteritis (reviewed by Shen & Winter, 1977). Dogs showed hepatic changes at 20 mg/kg bw per day, without concomitant gastrointestinal ulceration. Microscopic examination revealed portal fibrosis, bile-duct proliferation and inflammatory cell infiltration. Monkeys also showed no gastrointestinal ulceration, but they had hepatic effects similar to those seen in dogs.

Sulindac increases intestinal permeability in rats about five times less than indomethacin (Davies *et al.*, 1994), indicating that it may cause less small intestinal ulceration than indomethacin. Although the clinical literature suggests that the sulfone metabolite does not cause gastric ulcer formation, it exacerbated ulcer formation due to stress in rodents (Glavin & Sitar, 1986).

Long-term administration of sulindac at a dose of 40 mg/kg bw per day resulted in one death related to ulcer in 30 animals. No occult blood was detected in faeces at any dose, but at autopsy, on termination of the experiment, ulcerative enteritis was detected in about one-third of the rats at the highest dose. Only minor effects on liver and kidney function have been reported in rats, mice, dogs and monkeys, and only at a maximum dose of 40 mg/kg bw per day (Shen & Winter, 1977).

#### 7.2 Genetic and related effects

No data were available to the Working Group.

### 8. Summary of data

#### 8.1 Chemistry, occurrence and human exposure

Sulindac, a non-steroidal anti-inflammatory drug introduced in the 1970s, has analgesic, anti-inflammatory and anti-pyretic properties. It has limited use. The usual dose for musculoskeletal and joint disorders is 100–200 mg twice daily.

## 8.2 Metabolism and kinetics

Therapeutic doses of sulindac are almost totally absorbed by humans after oral administration. The peak plasma concentrations of sulindac and its sulfide metabolite are achieved within a few hours. Sulindac undergoes extensive enterohepatic recycling. The primary route of excretion in humans is via the urine as both sulindac and its sulfone metabolite.

## 8.3 Cancer-preventive effects

### 8.3.1 Humans

Randomized trials have shown conclusively that daily doses of 300–400 mg sulindac reduce the number and size of adenomatous colorectal polyps in patients with familial adenomatous polyposis. One observational study suggested that suppression of adenomas can be maintained with a daily intrarectal dose of 60 mg. Tumour inhibition may be incomplete: Three patients in whom polyps were suppressed by treatment with sulindac developed carcinoma of the rectum despite continuing therapy.

One small randomized trial of sulindac with low statistical power showed no regression of small (< 1 cm) adenomatous polyps in patients without familial adenomatous polyposis. Thus, the efficacy of sulindac against sporadic polyps remains to be established.

No adequate data were available to address the effect of sulindac on the risk for colorectal cancer.

### 8.3.2 Experimental animals

In seven studies in mouse and rat models, sulindac administered with or subsequent to carcinogens inhibited the development of pre-neoplastic and neoplastic lesions in the colon. In mice predisposed to intestinal malignancy by a germ-line mutation in the *Apc* gene, sulindac reduced the incidence of spontaneous intestinal tumours.

In single studies on the effects of sulindac on urinary bladder and lung carcinogenesis in mouse models, sulindac administered during and after the carcinogen inhibited tumour development. In two studies on mammary carcinogenesis in rats, sulindac and its metabolite sulindac sulfone, given one week after

carcinogen treatment, inhibited tumour development.

### 8.3.3 Mechanism of action

The precise mechanism for the chemopreventive action of sulindac is currently unknown. In human transformed intestinal epithelial cells in culture, doses of sulindac sulfide and sulindac sulfone higher than those that could be achieved pharmacologically inhibited cell growth and stimulated programmed cell death. In cells that overexpress the *Cox-2* gene, sulindac can reverse the tumorigenic phenotype by inhibiting *Cox-2*. Sulindac suppresses *ras* activation in colon mucosal cells of azoxymethane-treated rats.

## 8.4 Other beneficial effects

No data were available to the Working Group.

## 8.5 Carcinogenicity

No data were available to the Working Group.

## 8.6 Toxic effects

### 8.6.1 Humans

In common with other NSAIDs, sulindac increases the risk for gastrointestinal toxicity, including complications of ulcer, in a dose-dependent manner and increases the risks for toxic effects on the liver and kidney.

### 8.6.2 Experimental animals

Sulindac is well tolerated at high doses in rats, mice, dogs and rhesus monkeys. Minor effects occur in liver and kidney. Ulcerative enteritis is seen at very high doses in rats. The toxicity of the major human metabolites in these experimental models has not been reported.

## 9. Recommendations for research

Research to define the relative importance of the two metabolites, sulindac sulfide and sulindac sulfone, in the prevention of tumorigenesis in given human organs is of high priority. Better understanding of the chemopreventive action of sulindac in humans would require long-term studies of patients with familial adenomatous polyposis or large epidemiological studies in the general population.



## 10. Evaluation<sup>1</sup>

### 10.1 Cancer-preventive activity

#### 10.1.1 Humans

There is *limited evidence* that sulindac has cancer-preventive activity in patients with familial adenomatous polyposis. This evaluation is based on clear demonstration of the inhibition and regression of adenomatous polyps.

There is *inadequate evidence* that sulindac has cancer-preventive activity in people without familial adenomatous polyposis.

#### 10.1.2 Experimental animals

There is *sufficient evidence* that sulindac has cancer-preventive activity in experimental animals. This evaluation is based on models of cancers of the colon, urinary bladder, lung and mammary gland.

### 10.2 Overall evaluation

Randomized controlled trials in humans provide *limited evidence* that sulindac prevents colorectal cancer by suppressing adenomatous polyps in patients with familial adenomatous polyposis. There is no evidence for the prevention of cancers at other sites. Experimental animal models provide *sufficient evidence* that sulindac prevents cancers of the colon, mammary gland, lung and urinary bladder. These findings indicate the need for further evaluation of the cancer-preventive activity of sulindac against colorectal cancer in persons at high risk for the disease but are not applicable to the general population. The adverse effects of sulindac in humans comprise dose-dependent upper gastrointestinal bleeding and ulceration and hepatic and renal toxicity. If sulindac were to be used as a chemopreventive agent in large populations, the evidence of benefit would have to be clear and the benefits themselves significant.

## 11. References

- Adams, D.H., Howie, A.J., Michael, J., McConkey, B., Bacon, P.A. & Adu, D. (1986) Non-steroidal anti-inflammatory drugs and renal failure. *Lancet*, **i**, 57-60
- Alberts, D.S., Hixson, L., Ahnen, D., Bogert, C., Einspahr, J., Paranka, N., Brendel, K., Gross, P.H., Pamukcu, R. & Burt, R.W. (1995) Do NSAIDs exert their colon cancer chemoprevention activities through the inhibition of mucosal prostaglandin synthetase? *J. cell. Biochem.*, **22**, 18-23
- Beazer-Barclay, Y., Levy, D.B., Moser, A.R., Dove, W.F., Hamilton, S.R., Vogelstein, B. & Kinzler, K.W. (1996) Short communication: Sulindac suppresses tumorigenesis in the min mouse. *Carcinogenesis*, **17**, 1757-1760
- Belliveau, P. & Graham, A.M. (1984) Mesenteric desmoid tumor in Gardner's syndrome treated by sulindac. *Dis. Colon Rectum*, **27**, 53-54
- Bjarnasson, I., Zanelli, G., Prouse, P., Smethurst, P., Smith, T., Levi, S., Gumpel, M.J. & Levi, A.J. (1987) Blood and protein loss via small-intestinal inflammation induced by non-steroidal anti-inflammatory drugs. *Lancet*, **ii**, 711-714
- Boolbol, S.K., Dannenberg, A.J., Chadburn, A., Martucci, C., Guo, X.-J., Ramonetti, J.T., Abreu-Goris, M., Newmark, H.L., Lipkin, M.L., DeCosse, J.J. & Bertagnolli, M.M. (1996) Cyclooxygenase-2 overexpression and tumor formation are blocked by sulindac in a murine model of familial adenomatous polyposis. *Cancer Res.*, **56**, 2556-2560
- Brater, D.C., Anderson, S., Baird, B. & Campbell, W.B. (1985) Effects of ibuprofen, naproxen and sulindac on prostaglandins in men. *Kidney Int.*, **27**, 66-73
- Brater, D.C., Anderson, S.A., Brown Cartwright, D. & Toto, R.D. (1986) Effects of nonsteroidal anti-inflammatory drugs on renal function in patients with renal insufficiency and in cirrhotics. *Am. J. Kidney Dis.*, **8**, 351-355
- Budavari, S., ed. (1989) *The Merck Index*, Rahway, NJ, Merck & Co., p. 8964
- Castonguay, A., Pepin, P. & Stoner, G.D. (1991) Lung tumorigenicity of NNK given orally to A/J mice: Its application to chemopreventive efficacy studies. *Exp. Lung Res.*, **17**, 485-499

<sup>1</sup> For definitions of the italicized terms, see the Preamble, pp. 12-13

- Cerdán, F.J., Torres-Melero, J., Martínez, S., Gutiérrez del Olmo, A. & Balibrea, J.L. (1995) [Treatment with sulindac of adenomatous polyps in familial polyposis.] *Rev. esp. Enferm. Dig.*, **87**, 574–576 (in Spanish)
- Charneau, J., D'Aubigny, N., Burtin, P., Person, B. & Boyer, J. (1990) [Rectal micropolyps after total colectomy for familial polyposis: Efficacy of sulindac.] *Gastroenterol. clin. biol.*, **14**, 153–157 (in French)
- Chrischilles, E.A. & Wallace, R.B. (1993) Non-steroidal anti-inflammatory drugs and blood pressure in an elderly population. *J. Gerontol.*, **48**, 91–6
- Davies, N.M., Wright, M.R. & Jamali, F. (1994) Antiinflammatory drug-induced small intestinal permeability: The rat is a suitable model. *Pharm. Res.*, **11**, 1652–1656
- Debinski, H.S., Trojan, J., Nugent, K.P., Spigelman, A.D. & Phillips, R.K.S. (1995) Effect of sulindac on small polyps in familial adenomatous polyposis. *Lancet*, **345**, 855–856
- Dubach, U.C., Rosner, B. & Sturmer, T. (1991) An epidemiological study of abuse of analgesic drugs: Effects of phenacetin and salicylate on mortality and cardiovascular morbidity. *New Engl J. Med.*, **324**, 155–160
- Dubois, R.N., Awad, J., Morrow, J., Roberts, L.J. & Bishop, P.R. (1994a) Regulation of eicosanoid production and mitogenesis in rat intestinal epithelial cells by transforming growth factor  $\alpha$  and phorbol ester. *J. clin. Invest.*, **93**, 493–498
- DuBois, R.N., Bishop, P., Awad, J.A., Makita, K. & Lanhan, A. (1994b) Cloning and characterization of a growth factor-inducible cyclooxygenase gene from rat intestinal epithelial cells. *Am. J. Physiol.*, **266**, G8922–G8927
- Duggan, D.E., Hare, L.E., Ditzler, C.A., Lei, B.W. & Kwan, K.C. (1977) The disposition of sulindac. *Clin. Pharmacol. Ther.*, **21**, 326–335
- Eriksson, L.O., Sturfelt, G., Thysell, H. & Wollheim, F.A. (1991) Effects of sulindac and naproxen on prostaglandin excretion in patients with impaired renal function and rheumatoid arthritis. *Am. J. Med.*, **89**, 313–321
- Friend, W.G. (1990) Sulindac suppression of colorectal polyps in Gardner's syndrome. *Am. Family Phys.*, **41**, 891–894
- Friis, H. & Andreasen, P.B. (1992) Drug-induced hepatic injury: An analysis of 1100 cases reported to the Danish Committee on Adverse Drug Reactions between 1978 and 1987. *J. intern. Med.*, **232**, 133–138
- Furst, D.E., Herman, R.A., Koehnke, R., Ericksen, N., Hash, L., Riggs, C.E., Porras, A. & Veng-Pedersen, P. (1990) Effect of aspirin and sulindac on methotrexate clearance. *J. pharm. Sci.*, **79**, 782–786
- Garcia Rodriguez, L.A., Walker, A.M. & Perez Gutthann, S. (1992) Nonsteroidal antiinflammatory drugs and gastrointestinal hospitalizations in Saskatchewan: A cohort study. *Epidemiology*, **3**, 337–342
- Giardiello, F.M., Hamilton, S.R., Krush, A.J., Piantadosi, S., Hyland, L.M., Celano, P., Booker, S.V., Robinson, C.R. & Offerhaus, G.J.A. (1993) Treatment of colonic and rectal adenomas with sulindac in familial adenomatous polyposis. *New Engl. J. Med.*, **328**, 1313–1316
- Glavin, G.B. & Sitar, D.S. (1986) The effects of sulindac and its metabolites on acute stress-induced gastric ulcers in rats. *Toxicol. appl. Pharmacol.*, **83**, 386–389
- Gonzaga, R.A.F., Lima, F.R., Carneiro, S., Maciel, J. & Aramante, M., Jr (1985) Sulindac treatment for familial polyposis coli (Letter to the Editor). *Lancet*, **i**, 751
- Griffin, M.R., Piper, J.M., Dougherty, J.R., Snowden, W. & Ray, W.A. (1991) Non-steroidal anti-inflammatory drug use and increased risk for peptic ulcer disease in elderly persons. *Ann. intern. Med.*, **114**, 257–263
- Gurwitz, J.H., Avorn, J., Ross Degnan, D. & Lipsitz, L.A. (1990) Nonsteroidal anti-inflammatory drug associated azotemia in the very old. *J. Am. med. Assoc.*, **264**, 471–475
- Gurwitz, J.H., Avorn, J., Bohn, R.L., Glynn, R.J., Monane, M. & Mogun, H. (1994) Initiation of antihypertensive treatment during nonsteroidal anti-inflammatory drug therapy. *J. Am. med. Assoc.*, **272**, 781–786

- Hanif, R., Pittas, A., Feng, Y., Koutsos, M.I., Quao, L., Staiano-Coico, L., Shiff, S.I. & Rigas, B. (1996) Effects of nonsteroidal anti-inflammatory drugs on proliferation and on induction of apoptosis in colon cancer cells by a prostaglandin-independent pathway. *Biochem. Pharmacol.*, **52**, 237–245
- Hare, L.E., Ditzler, C.A., Hichens, M., Rosegay, A. & Duggan D.E. (1977) Analysis of sulindac and metabolites by combined isotope dilution—radioimmunoassay. *J. pharm. Sci.*, **66**, 414–417
- Heller, S.R., Fellow, I.W & Ogilvie, A.L. (1982) Non-steroidal anti-inflammatory drugs and benign oesophageal stricture. *Br. med. J.*, **285**, 167–168
- Henry, D., Lim, L.L., Garcia Rodriguez, L.A., Perez Gutthann, S., Carson, J.L., Griffin, M., Savage, R., Logan, R., Moride, Y., Hawkey, C., Hill, S.L. & Fries, J.T. (1996) Variability in risk of gastrointestinal complications with individual non-steroidal anti-inflammatory drugs. Results of a collaborative meta-analysis. *Br. med. J.*, **312**, 1563–1566
- Hucker, H.B., Stauffer, S.C., White, S.D., Rhodes, R.E., Arison, B.H., Umbenhauer, E.R., Bower, R.J. & McMahon, F.G. (1973) Physiologic disposition and metabolic fate of a new anti-inflammatory agent, cis-5-fluoro-2-methyl-1-[p-(methysulfinyl)-benzylidenyl]indeno-3-acetic acid in the rat, dog, rhesus monkey, and man. *Drug Metab. Disposition*, **1**, 721–736
- Izes, J.K., Zinman, L.N. & Larsen, C.R. (1996) Regression of large pelvic desmoid tumor by tamoxifen and sulindac. *Urology*, **47**, 756–759
- Jalbert, G. & Castonguay, A. (1992) Effects of NSAIDs on NNK-induced pulmonary and gastric tumorigenesis in A/J mice. *Cancer Lett.*, **66**, 21–28
- Johnson, A.G., Nguyen, T.V. & Day, R.O. (1994) Do nonsteroidal anti-inflammatory drugs affect blood pressure? A meta-analysis. *Ann. intern. Med.*, **121**, 289–300
- Jones, R.H. & Tait, C.L. (1995) Gastrointestinal side-effects of NSAIDs in the community. *Br J. clin. Pract.*, **49**, 67–70
- Kadmon, M., Möslein, G., Buhr, H.J. & Herfarth, C. (1995) [Desmoids in patients with familial polyposis (FAP).] *Klin. ther. Beobachtung. Heidelberger Polyposis-Reg. Chirurg.*, **66**, 997–1005 (in German)
- Kikendale, J.W. (1991) Pill-induced oesophageal injury. *Gastroenterol. Clin. North Am.*, **20**, 835–846
- Klein, W.A., Miller, H.H., Anderson, M. & DeCosse, J.J. (1987) The use of indomethacin, sulindac, and tamoxifen for the treatment of desmoid tumors associated with familial polyposis. *Cancer*, **60**, 2863–2868
- Kleinknecht, D., Landais, P. & Goldfarb, B. (1986) Analgesic and non-steroidal anti-inflammatory drug associated acute renal failure: A prospective collaborative study. *Clin. Nephrol.*, **25**, 275–281
- Kramer, W.B., Saade, G., Ou, C.N., Rognerud, C., Dorman, K., Mayes, M. & Moise, K.J. (1995) Placental transfer of sulindac and its active sulfide metabolite in humans. *Am. J. Obstet. Gynecol.*, **172**, 886–890
- Kwo, P.Y. & Tremaine, W.J. (1995) Nonsteroidal anti-inflammatory drug-induced enteropathy: Case discussion and review of the literature. *Mayo Clin. Proc.*, **70**, 55–61
- Labayle, D., Fischer, D., Vielh, P., Drouhin, F., Pariente, A., Bories, C., Duhamel, O., Troussset, M. & Attali, P. (1991) Sulindac causes regression of rectal polyps in familial adenomatous polyposis. *Gastroenterology*, **101**, 635–639
- Ladenheim, J., Garcia, G., Titzer, D., Herzenberg, H., Lavori, P., Edson, R. & Omary, M.B. (1995) Effect of sulindac on sporadic colonic polyps. *Gastroenterology*, **108**, 1083–1087
- Laffi, G., Daskalopoulos, G., Kronborg, I., Hsueh, W., Gentilini, P. & Zipser, R.D. (1986) Effects of sulindac and ibuprofen in patients with cirrhosis and ascites. An explanation for the renal-sparing effect of sulindac. *Gastroenterology*, **90**, 182–187
- Langman, M.J., Morgan, L. & Worrall, A. (1985) Use of anti-inflammatory drugs by patients admitted with small or large bowel perforations and haemorrhage. *Br med. J.*, **290**, 347–349
- Lee, S.C. & Renwick, A.G. (1995a) Sulphoxide reduction by rat intestinal flora and by *Escherichia coli* in vitro. *Biochem. Pharmacol.*, **49**, 1567–1576
- Lee, S.C. & Renwick, A.G. (1995b) Sulphoxide reduction by rat and rabbit tissues in vitro. *Biochem. Pharmacol.*, **49**, 1557–1565

- Lu, X., Xie, W., Reed, D., Bradshaw, W.S. & Simmons, D.L. (1995) Nonsteroidal anti-inflammatory drugs cause apoptosis and induce cyclooxygenase in chicken embryo fibroblasts. *Proc. natl Acad. Sci. USA*, **92**, 7961–7965
- Lynch, H.T., Thorson, A.G. & Smyrk, T. (1995) Rectal cancer after prolonged sulindac chemoprevention. A case report. *Cancer*, **75**, 936–938
- Mäkelä, J.T. & Laitinen, S. (1994) Sulindac therapy for familial adenomatous polyposis after colectomy and ileorectal anastomosis. *Ann. chir. gynaecol.*, **83**, 265–267
- Matsushashi, N., Yamada, A., Hiraishi, M., Konishi, T., Minota, S., Saito, T., Sugano, K., Yazaki, Y., Mori, M. & Shiga, J. (1992) Multiple strictures of the small intestine after long-term nonsteroidal anti-inflammatory drug therapy. *Am. J. Gastroenterol.*, **87**, 1183–1186
- McManus, P., Primrose, J.G., Henry, D.A., Birkett, D.J., Lindner, J. & Day, R.O. (1996) Pattern of nonsteroidal anti-inflammatory drug use in Australia 1990–1994. A report from the Drug Utilization Sub-committee of Pharmaceutical Benefits Advisory Committee. *Med. J. Aust.*, **164**, 589–592
- Miller, M.J., Bednar, M.M. & McGiff, J.C. (1984) Renal metabolism of sulindac: Functional implications. *J. Pharmacol. exp. Ther.*, **231**, 449–456
- Moorghen, M., Ince, P., Finney, K.J., Sunter, J.P., Appleton, D.R. & Watson, A.J. (1988) A protective effect of sulindac against chemically-induced primary colonic tumours in mice. *J. Pathol.*, **156**, 341–347
- Murray, M.D. & Brater, D.C. (1990) Adverse effects of nonsteroidal anti-inflammatory drugs on renal function. *Ann. intern. Med.*, **112**, 559–560
- Murray, M.D. & Brater, D.C. (1993) Renal toxicity of the nonsteroidal anti-inflammatory drugs. *Ann. Rev. Pharmacol. Toxicol.*, **33**, 435–465
- Murray, M.D., Black, P.K., Kuzmik, D.D., Haag, K.M., Manatunga, A.K., Mullin, M.A., Hall, S.D. & Brater, D.C. (1995) Acute and chronic effects of nonsteroidal antiinflammatory drugs on glomerular filtration rate in elderly patients. *Am. J. med. Sci.*, **310**, 188–197
- Niv, Y. & Fraser, G.M. (1994) Adenocarcinoma in the rectal segment in familial polyposis coli is not prevented by sulindac therapy. *Gastroenterology*, **107**, 854–857
- Nugent, K.P., Farmer, K.C.R., Spigelman, A.D., Williams, C.B. & Phillips, R.K.S. (1993) Randomized controlled trial of the effect of sulindac on duodenal and rectal polyposis and cell proliferation in patients with familial adenomatous polyposis. *Br. J. Surg.*, **80**, 1618–1619
- Ostensen, M. (1994) Optimisation of anti-rheumatic drug treatment in pregnancy. *Clin. Pharmacokinet.*, **27**, 486–503
- Patrono, C. (1986) Inhibition of renal prostaglandin synthesis in man: Methodological and clinical implications. *Scand. J. Rheumatol.*, **Suppl. 62**, 14–25
- Pepin, P., Bouchard, L., Nicole, P. & Castonguay, A. (1992) Effects of sulindac and oltipraz on the tumorigenicity of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in A/J mouse lung. *Carcinogenesis*, **13**, 341–348
- Pereira, M.A., Barnes, L.H., Rassman, V.L., Kelloff, G.V. & Steele, V.E. (1994) Use of azoxymethane-induced foci of aberrant crypts in rat colon to identify potential cancer chemopreventive agents. *Carcinogenesis*, **15**, 1049–1054
- Perez-Gutthann, S., Garcia Rodriguez, L.A., Raiford, D.S., Duque-Oliart, A. & Ris-Romeu, J. (1996) Nonsteroidal anti-inflammatory drugs and the risk of hospitalization for acute renal failure. *Arch. intern. Med.*, **156**, 2433–2439
- Pernerger, T.V., Whelton, P.K. & Klag, M.J. (1994) Risk of kidney failure associated with the use of acetaminophen, aspirin and nonsteroidal antiinflammatory drugs. *New Engl. J. Med.*, **331**, 1675–1679
- Piazza, G.A., Rahm, A.L., Krutzsch, M., Sperl, G., Paranka, N.S., Gross, P.H., Brendel, K., Burt, R.W., Alberts, D.S., Pamukcu, R. & Ahnen, D.J. (1995) Antineoplastic drugs sulindac sulfide and sulfone inhibit cell growth by inducing apoptosis. *Cancer Res.*, **55**, 3110–3116
- Pope, J.E., Anderson, J.J. & Felson, D.T. (1993) A meta-analysis of the effects of nonsteroidal anti-inflammatory drugs on blood pressure. *Arch. intern. Med.*, **153**, 477–484
- Radack, K. & Deck, C. (1987) Do nonsteroidal anti-inflammatory drugs interfere with blood pressure control in hypotensive patients? *J. gen. intern. Med.*, **2**, 108–112

- Rao, C.V., Rivenson, A., Simi, B., Zang, E., Kelloff, G., Steele, V. & Reddy, B.S. (1995) Chemo-prevention of colon carcinogenesis by sulindac, a non-steroidal anti-inflammatory agent. *Cancer Res.*, **55**, 1464–1472
- Rao, K.V.N., Detrisac, C.J., Steele, V.E., Hawk, E.T., Kelloff, G.J. & McCormick, D.L. (1996) Differential activity of aspirin, ketoprofen and sulindac as cancer chemopreventive agents in the mouse urinary bladder. *Carcinogenesis*, **17**, 1435–1438
- Ravis, W.R., Diskin, C.J., Campagna, K.D., Clark, C.R. & McMillian, C.L. (1993) Pharmacokinetics and dialyzability of sulindac and metabolites in patients with end-stage renal failure. *J. clin. Pharmacol.*, **33**, 527–534
- Reddy, B.S., Rao, C.V., & Seibert, K. (1996) Evaluation of cyclooxygenase-2 inhibitor for potential chemopreventive properties in colon carcinogenesis. *Cancer Res.*, **56**, 4566–4569.
- Reynolds, J.E.F., ed. (1993) *Martindale, The Extra Pharmacopoeia*, 13th Ed., London, The Pharmaceutical Press, pp. 34–35
- Rigau, J., Pigué, J.M., Rubio, E., Planas, R., Tarrech, J.M. & Bordas, J.M. (1991) Effects of long-term sulindac therapy on colonic polyposis. *Ann. intern. Med.*, **115**, 952–954
- Rodriguez, L.A.G., Williams, R., Derby, L.E., Dean, A.D. & Jick, H. (1994) Acute liver injury associated with nonsteroidal anti-inflammatory drugs and the role of risk factors. *Arch. intern. Med.*, **154**, 311–316
- Roupe, G., Ahlmen, M., Fagerberg, B. & Suurkula, M. (1986) Toxic epidermal necrolysis with extensive mucosal erosions of the gastrointestinal and respiratory tracts. *Int. Arch. Allergy appl. Immunol.*, **80**, 145–151
- Russeva, V., Stavreva, N., Rakovska, R. & Michailova, D. (1994) Binding of sulindac to human serum albumin studied by circular dichroism. *Arzneimittel.-forsch./Drug Res.*, **44**, 159–162
- Sandler, D.P., Smith, J.C., Weinberg, C.R., Buckalew, V.M., Dennis, V.W., Blythe, W.B. & Burgess, W.P. (1989) Analgesic use and chronic renal disease. *New Engl. J. Med.*, **320**, 1238–1243
- Schnitzler, M., Dwight, T. & Robinson, B.G. (1996) Sulindac increases the expression of APC mRNA in malignant colonic epithelial cells: An in vitro study. *Gut*, **38**, 707–713
- Shen, T.Y. & Winter, C.A. (1977) Chemical and biological studies on indomethacin, sulindac and their analogs. *Adv. Drug Res.*, **12**, 89–295
- Shiff, S.J., Tsai, L.-L., Qiao, L. & Rigas, B. (1995) Sulindac sulfide inhibits proliferation, causes cell cycle quiescence, and induces apoptosis in HT29 human colon adenocarcinoma cells. *J. clin. Invest.*, **96**, 491–503
- Shiff, S.J., Koutsos, M.I., Qiao, L. & Rigas, B. (1996) Nonsteroidal antiinflammatory drugs inhibit the proliferation of colon adenocarcinoma cells: Effects on cell cycle and apoptosis. *Exp. Cell. Res.*, **222**, 179–188
- Shimek, J.L., Rao, N.G.S. & Khalil, S.K.W. (1981) High-performance liquid-chromatographic analysis of tolmetin, indomethacin and sulindac in plasma. *J. liq. Chromatogr.*, **4**, 1987–2013
- Siglin, J.C., Barch, D.H. & Stoner, G.D. (1995) Effects of dietary phenethyl isothiocyanate, ellagic acid, sulindac and calcium on the induction and progression of *N*-nitrosomethylbenzylamine-induced esophageal carcinogenesis in rats. *Carcinogenesis*, **16**, 1101–1106
- Siluveru, M. & Stewart, J.T. (1995) Determination of sulindac and its metabolites in human serum by reversed-phase high-performance liquid chromatography using on-line post-column ultraviolet irradiation and fluorescence detection. *J. Chromatogr.*, **673**, 91–96
- Singh, J. & Reddy, B.S. (1995) Molecular markers in chemoprevention of colon cancer. Inhibition of expression of ras-p21 and p53 by sulindac during azoxymethane induced colon carcinogenesis. *Ann. N.Y. Acad. Sci.*, **768**, 205–209
- Singh, J., Rao, C.V., Kulkarni, N., Simi, B. & Reddy, B.S. (1994) Molecular markers as intermediate end-points in chemoprevention of colon cancer: Modulation of *ras* activation by sulindac and phenyl hexylisothiocyanate during colon carcinogenesis. *Int. J. Oncol.*, **5**, 1009–1018
- Skinner, S.A., Penney, A.G. & O'Brien, P.E. (1991) Sulindac inhibits the rate of growth and appearance of colon tumors in the rat. *Arch. Surg.*, **126**, 1094–1096
- Smalley, W.E., Ray, W.A., Daugherty, J.R. & Griffin, M.R. (1995) Nonsteroidal anti-inflammatory drugs and the incidence of hospitalizations for peptic ulcer disease in elderly persons. *Am J. Epidemiol.*, **141**, 539–545

- Spagnesi, M.T., Tonelli, F., Dolara, P., Caderni, G., Valanzano, R., Anastasi, A. & Bianchini, F. (1994) Rectal proliferation and polyp occurrence in patients with familial adenomatous polyposis after sulindac treatment. *Gastroenterology*, **106**, 362–366
- Stillman, M.T. & Schlesinger, P.A. (1990) Non-steroidal anti-inflammatory drug nephrotoxicity. Should we be concerned? *Arch. intern. Med.*, **150**, 268–270
- Strong, H.A., Warner, N.J., Renwick, A.G. & George, C.F. (1985) Sulindac metabolism: The importance of an intact colon. *Clin. Pharmacol. Ther.*, **38**, 387–393
- Strong, H.A., Renwick, A.G., George, C.F., Liu, Y.F. & Hill, M.J. (1987) The reduction of sulphinpyrazone and sulindac by intestinal bacteria. *Xenobiotica*, **17**, 685–696
- Stubbs, R.J., Ng, L.L., Entwistle, L.A. & Bayne W.F. (1987) Analysis of sulindac and metabolites in plasma and urine by high-performance liquid chromatography. *J. Chromatogr.*, **413**, 171–180
- Swanson, B.N. & Boppana, V.K. (1981) Measurement of sulindac and its metabolites in human plasma and urine by high-performance liquid chromatography. *J. Chromatogr.*, **225**, 123–130
- Tarazi, E.M., Harter, J.G., Zimmerman, H.J., Oshak, K.G. & Eaton, R.A. (1993) Sulindac-associated hepatic injury: Analysis of 91 cases reported to the Food and Drug Administration. *Gastroenterology*, **104**, 569–574
- Thompson, H.J., Briggs, S., Paranak, N.S., Piazza, G.A., Brendel, K., Gross, P.H., Sperl, G.J., Pamukcu, R. & Ahnen, D.J. (1995) Brief communications: Inhibition of mammary carcinogenesis in rats by sulfone metabolite of sulindac. *J. natl Cancer Inst.*, **87**, 1259–1260
- Thompson, H.J., Jiang, C., Lu, J., Mehta, R.G., Piazza, G.A., Paranka, N.S., Pamukcu, R. & Ahnen, D.J. (1997) Sulfone metabolite of sulindac inhibits mammary carcinogenesis. *Cancer Res.*, **57**, 267–271
- Thorson, A.G., Lynch, H.T. & Smyrk, T.C. (1994) Rectal cancer in FAP patients after sulindac. *Lancet*, **343**, 180
- Tonelli, F. & Valanzano, R. (1993) Sulindac in familial adenomatous polyposis (Letter to the Editor). *Lancet*, **342**, 1120
- Tsuji, M. & DuBois, R.N. (1995) Alterations in cellular adhesion and apoptosis in epithelial cells overexpressing prostaglandin endoperoxide synthase-2. *Cell*, **83**, 493–501
- Venuti, M.C., Young, J.M., Maloney, P.J., Johnson, D. & McGreevy, K. (1989) Synthesis and biological evaluation of omega-(N,N,N-trialkylammonium) alkyl esters and thioesters of carboxylic acid nonsteroidal antiinflammatory agents. *Pharm. Res.*, **6**, 867–873
- Waddell, W.R. & Kirsch, W.M. (1991) Testolactone, sulindac, warfarin, and vitamin K<sub>1</sub> for unresectable desmoid tumors. *Am. J. Surg.*, **161**, 416–421
- Waddell, W.R. & Loughry, R.W. (1983) Sulindac for polyposis of the colon. *J. surg. Oncol.*, **24**, 83–87
- Waddell, W.R., Ganser, G.F., Cerise, E.J. & Loughry, R.W. (1989) Sulindac for polyposis of the colon. *Am. J. Surg.*, **157**, 175–179
- Whelton, A. & Hamilton, C.W. (1991) Non-steroidal anti-inflammatory drugs: Effects on kidney function. *J. clin. Pharmacol.*, **31**, 588–598
- Whelton, A., Stout, R.L., Spilman, P.S. & Klassen, D.K. (1990) Renal effects of ibuprofen, piroxicam and sulindac in patients with asymptomatic renal failure. A prospective, randomized, crossover comparison. *Ann. intern. Med.*, **112**, 568–576
- Winde, G., Gumbinger, H.G., Osswald, H., Kemper, F. & Bünte, H. (1993) The NSAID sulindac reverses rectal adenomas in colectomized patients with familial adenomatous polyposis: Clinical results of a dose-finding study on rectal sulindac administration. *Int. J. colorectal Dis.*, **8**, 13–17
- Winde, G., Schmid, K.W., Schlegel, W., Fischer, R., Osswald, H. & Bünte, H. (1995) Complete reversion and prevention of rectal adenomas in colectomized patients with familial adenomatous polyposis by rectal low-dose sulindac maintenance treatment. Advantages of a low-dose nonsteroidal anti-inflammatory drug regimen in reversing adenomas exceeding 33 months. *Dis. Colon Rectum*, **38**, 813–830
- Winde, G., Schmid, K.W., Brandt, B., Müller, C. & Osswald, H. (1997) Clinical and genomic influence of sulindac on rectal mucosa in familial adenomatous polyposis (FAP). *Dis. Colon Rectum* (in press)

Wong, D.G., Spence, J.D., Lamki, L., Freeman, D. & McDonald, J.W. (1986) Effect of non-steroidal anti-inflammatory drugs on control of hypotension by beta-blockers and diuretics. *Lancet*, **i**, 997-1001

Zamboni, R., Corti, P. & Nucci, L. (1983) Polarographic analysis of sulindac in pharmaceutical products and differential pulse polarographic analysis of sulindac and its metabolites in plasma. *Boll. Chim. Farm.*, **122**, 505-511