References

Adams S, Cotti C (2008). Drunk driving after the passage of smoking bans in bars. *J Public Econ*, 92(5-6):1288-1305.

Adda J, Cornaglia F (2005). The effects of taxes and bans on passive smoking. University College London, Institute for Fiscal Studies. Department of Economics. Centre for microdata and methods and practice (CEMMAP) Working Paper 20/05 (http://cemmap.ifs.org.uk/wps/cwp2005.pdf).

Akhtar PC, Currie DB, Currie CE, et al. (2007). Changes in child exposure to environmental tobacco smoke (CHETS) study after implementation of smoke-free legislation in Scotland: national cross sectional survey. Br Med J, 335(7619):545-549.

Al-Delaimy W, Fraser T, Woodward A (2001a). Nicotine in hair of bar and restaurant workers. *N Z Med J*, 114(1127):80-83.

Al-Delaimy WK (2002). Hair as a biomarker for exposure to tobacco smoke. *Tob Control*, 11(3):176-182.

Al-Delaimy WK, Crane J, Woodward A (2001b). Passive smoking in children: effect of avoidance strategies, at home as measured by hair nicotine levels. *Arch Environ Health*, 56(2):117-122.

Al-Delaimy WK, Mahoney GN, Speizer FE, et al. (2002a). Toenail nicotine levels as a biomarker of tobacco smoke exposure. Cancer Epidemiol Biomarkers Prev, 11(11):1400-1404.

Al-Delaimy WK, Crane J, Woodward A (2002b). Is the hair nicotine level a more accurate biomarker of environmental tobacco smoke exposure than urine cotinine? *J Epidemiol Community Health*, 56(1):66-71.

Al-Delaimy WK, Cho E, Chen WY, et al. (2004). A prospective study of smoking and risk of breast cancer in young adult women. Cancer Epidemiol Biomarkers Prev, 13(3):398-404.

Al-Delaimy WK, Pierce J, Messer K, et al. (2007). The California Tobacco Control Program's effect on adult smokers: (2) Daily cigarette consumption levels. *Tob Control*, 16(2):91-95.

Al-Delaimy WK, White MM, Trinidad DR, et al. (2008). The California Tobacco Program: can we maintain the progress? Results from the California Tobacco Survey, 1990-2005. Sacramento, CA, California Department of Public Health (http://ssdc.ucsd.edu/tobacco/).

Alamar B, Glantz S (2004). Smoke-free ordinances increase restaurant profit and value. *Contemp Econ Policy*, 22(4):520-525.

Alamar B, Glantz S (2007). Effect of smokefree bar laws on bar value and profits. *Am J Public Health*, 97(8):1401-1402.

Alberg AJ, Kouzis A, Genkinger JM, et al. (2007). A prospective cohort study of bladder cancer risk in relation to active cigarette smoking and household exposure to secondhand cigarette smoke. Am J Epidemiol, 165(6):660-666.

Albers AB, Siegel M, Cheng DM, et al. (2007). Effect of smoking regulations in local restaurants on smokers' anti-smoking attitudes and quitting behaviours. *Tob Control*, 16(2):101-106.

Albers AB, Biener L, Siegel M, et al. (2008). Household smoking bans and adolescent antismoking attitudes and smoking initiation: findings from a longitudinal study of a Massachusetts youth cohort. Am J Public Health, 98(10):1886-1893.

Allred EN, Bleecker E, Chaitman B, et al. (1989). Short-term effects of carbon monoxide exposure on the exercise performance of subjects with coronary artery disease. N Engl J Med, 321(21):1426-1432.

Allwright S, Paul G, Greiner B, et al. (2005). Legislation for smoke-free workplaces and health of bar workers in Ireland: before and after study. *Br Med J*, 331(7525):1117-1120.

Alpert HR, Carpenter CM, Travers MJ, et al. (2007). Environmental and economic evaluation of the Massachusetts smoke-free workplace law. J Community Health, 32(4):269-281.

American Society of Heating, Refrigerating, and Air-Conditioning Engineers (2005). Environmental tobacco smoke. Atlanta, GA, American Society of Heating Refrigerating and Air-Conditioning Engineers (http://www.ashrae.org/doclib/20058211239 347.pdf).

American Nonsmokers' Rights Foundation (2009). 100% smokefree correctional facilities. Berkeley, CA (http://no-smoke.org/pdf/ 100smokefreeprisons.pdf).

Andersen MR, Leroux B, Bricker J, et al. (2004). Antismoking parenting practices are associated with reduced rates of adolescent smoking. *Arch Pediatr Adolesc Med*, 158(4):348-352.

Anderson HR, Cook DG (1997). Passive smoking and sudden infant death syndrome: review of the epidemiological evidence. *Thorax*, 52(11):1003-1009.

Andreeva TI, Krasovsky KS, Semenova DS (2007). Correlates of smoking initiation among young adults in Ukraine: a cross-sectional study. *BMC Public Health*, 7:106.

Arcus J, Boey S, Bradshaw R, et al. (2007). SMOKES 2007 - Smoking outdoors in a kids environment: a pilot study: evaluating the Upper Hutt City Council smokefree parks policy. Wellington, New Zealand, University of Otago https://www.smokefreecouncils.org.nz/fileadmin/clients/cancer-society/pdf/The_SMOKE_study_2007_report_FINAL.pdf).

Arheart KL, Lee D, Dietz N, et al. (2008). Declining trends in serum cotinine levels in US worker groups: the power of policy. *J Occup Environ Med*, 50(1):57-63.

Ashley MJ, Ferrence R (1998). Reducing children's exposure to environmental tobacco smoke in homes: issues and strategies. *Tob Control*, 7(1):61-65.

Ashley MJ, Cohen J, Ferrence R, et al. (1998). Smoking in the home: changing attitudes and current practices. Am J Public Health, 88(5):797-800.

Auerbach O, Petrik TG, Stout AP, et al. (1956). The anatomical approach to the study of smoking and bronchogenic carcinoma; a preliminary report of forty-one cases. *Cancer*, 9(1):76-83.

Auerbach O, Forman JB, Gere JB, *et al.* (1957). Changes in the bronchial epithelium in relation to smoking and cancer of the lung; a report of progress. *N Engl J Med*, 256(3):97-104.

Australian National Research Council (NRC) (1986). Environmental tobacco smoke: measuring exposures and assessing health effects. Washington, DC, National Academy Press (http://books.nap.edu/catalog.php?record_id=943).

Author Unknown (1961). The great debate [editorial]. *N Engl J Med*, 264:1266.

Aveyard P, Markham W, Cheng K (2004). A methodological and substantive review of the evidence that schools cause pupils to smoke. *Soc Sci Med*, 58(11):2253-2265.

Baker JA, Odunuga OO, Rodabaugh KJ, *et al.* (2006). Active and passive smoking and risk of ovarian cancer. *Int J Gynecol Cancer*, 16(Suppl 1):211-218.

Bakoula CG, Kafritsa Y, Kavadias G, *et al.* (1997). Factors modifying exposure to environmental tobacco smoke in children (Athens, Greece). *Cancer Causes Control*, 8(1):73-76.

Barnett TA, Gauvin L, Lambert M, et al. (2007). The influence of school smoking policies on student tobacco use. Arch Pediatr Adolesc Med, 161(9):842-848.

Barnoya J, Glantz S (2005). Cardiovascular effects of secondhand smoke: nearly as large as smoking. *Circulation*, 111(20):2684-2698.

Barnoya J, Mendoza-Montano C, Navas-Acien A (2007). Secondhand smoke exposure in public places in Guatemala: comparison with other Latin American countries. *Cancer Epidemiol Biomarkers Prev*, 16(12):2730-2735.

Barone-Adesi F, Vizzini L, Merletti F, et al. (2006). Short-term effects of Italian smoking regulation on rates of hospital admission for acute myocardial infarction. Eur Heart J, 27(20):2468-2472.

Barrientos-Gutierrez T, Reynales-Shigematsu L, Vila-Tang E, et al. (2007a). [Environmental tobacco smoke exposure in homes of Mexico City: analysis of environmental samples and children and women hair]. Salud Publica Mex, 49(Suppl 2):S224-S232.

Barrientos-Gutierrez T, Valdes-Salgado R, Reynales-Shigematsu L, *et al.* (2007b). [Involuntary exposure to tobacco smoke in public places in Mexico City] *Salud Publica Mex*, 49(Suppl 2):S205-S212.

Bartecchi C, Alsever R, Nevin-Woods C, et al. (2006). Reduction in the incidence of acute myocardial infarction associated with a citywide smoking ordinance. *Circulation*, 114(14):1490-1496.

Baska T, Sovinova H, Nemeth A, et al. (2007). Environmental tobacco smoke of youngsters in Czech Republic, Hungary, Poland and Slovakia--findings from the Global Youth Tobacco Survey (GYTS). Int J Public Health, 52(1):62-66.

Bates MN, Fawcett J, Dickson S, et al. (2002). Exposure of hospitality workers to environmental tobacco smoke. *Tob Control*, 11(2):125-129.

Bauer JE,, Hyland A, et al. (2005). A longitudinal assessment of the impact of smoke-free worksite policies on tobacco use. Am J Public Health, 95(6):1024-1029.

Behan DF, Eriksen M, Lin Y (2005). Economic effects of environmental tobacco smoke. Society of Actuaries (http://www.soa.org/research/life/research-economic-effect.aspx).

Bello S, Soto M, Michalland S, et al. (2004). Encuesta nacional de tabaquismo en funcionarios de salud. Rev Med Chile, 132(2):223-232.

Bennett WP, Alavanja MC, Blomeke B, et al. (1999). Environmental tobacco smoke, genetic susceptibility, and risk of lung cancer in never-smoking women. J Natl Cancer Inst, 91(23):2009-2014.

Benowitz NL (1999). Biomarkers of environmental tobacco smoke exposure. Environ Health Perspect, 107(Suppl 2):349-355

Berg CJ, Cox L, Nazir N, et al. (2006). Correlates of home smoking restrictions among rural smokers. *Nicotine Tob Res*, 8(3):353-360.

Berman S (2003). Caring for parents vs caring for children: is there a difference? *Arch Pediatr Adolesc Med*, 157(3):221-222.

Berman BA, Wong GC, Bastani R, et al. (2003). Household smoking behavior and ETS exposure among children with asthma in low-income, minority households. Addict Behav, 28(1):111-128.

Best JA, Hakstian AR (1978). A situationspecific model for smoking behavior. *Addict Behav*, 3(2):79-92. Bewley BR, Johnson M, Banks M (1979). Teachers' smoking. *J Epidemiol Community Health*, 33(3):219-222.

Biener L, Nyman AL (1999). Effect of workplace smoking policies on smoking cessation: results of a longitudinal study. *J Occup Environ Med*, 41(12):1121-1127.

Biener L, Cullen D, Di Z, et al. (1997). Household smoking restrictions and adolescents' exposure to environmental tobacco smoke. *Prev Med*, 26(3):358-363.

Biener L, Harris J, Hamilton W (2000). Impact of the Massachusetts tobacco control programme: population based trend analysis. *Br Med J*, 321(7257):351-354.

Biener L, Garrett C, Skeer M, et al. (2007). The effects on smokers of Boston's smoke-free bar ordinance: a longitudinal analysis of changes in compliance, patronage, policy support, and smoking at home. J Public Health Manag Pract, 13(6):630-636.

Bird Y, Moraros J, Olsen L, et al. (2007). Smoking practices, risk perception of smoking, and environmental tobacco smoke exposure among 6th-grade students in Ciudad Juarez, Mexico. Nicotine Tob Res, 9(2):195-203.

Bjerregaard BK, Raaschou-Nielsen O, Sorensen M, et al. (2006). Tobacco smoke and bladder cancer--in the European prospective investigation into cancer and nutrition. *Int J Cancer*, 119(10):2412-2416.

Blackburn C, Spencer N, Bonas S, et al. (2003). Effect of strategies to reduce exposure of infants to environmental tobacco smoke in the home: cross sectional survey. *Br Med J*, 327(7409):257.

Blecher EH (2006). The effects of the tobacco products control amendment Act of 1999 on restaurant revenues in South Africa: A panel data approach. S Afr J Econ, 74(1):123-130.

Bloch M, Shopland D (2000). Outdoor smoking bans: more than meets the eye. *Tob Control*, 9(1):99.

Bolte G, Heitmann D, Kiranoglu M, et al. (2008). Exposure to environmental tobacco smoke in German restaurants, pubs and discotheques. *J Expo Sci Environ Epidemiol*, 18(3):262-271.

Bonita R, Duncan J, Truelsen T, et al. (1999). Passive smoking as well as active smoking increases the risk of acute stroke. *Tob Control*, 8(2):156-160.

Borland R, Pierce JP, Burns DM, et al. (1992). Protection from environmental tobacco smoke in California. The case for a smoke-free workplace [see comment]. *JAMA*, 268(6):749-752.

Borland R, Cappiello M, Owen N (1997). Leaving work to smoke. *Addiction*, 92(10):1361-1368.

Borland R, Mullins R, Trotter L, *et al.* (1999). Trends in environmental tobacco smoke restrictions in the home in Victoria, Australia. *Tob Control.* 8(3):266-271.

Borland R, Yong H, Cummings K, et al. (2006a). Determinants and consequences of smoke-free homes: findings from the International Tobacco Control (ITC) four country survey. *Tob Control*, 15(Suppl 3):iii42-iii50.

Borland R, Yong H, Siahpush M, et al. (2006b). Support for and reported compliance with smoke-free restaurants and bars by smokers in four countries: findings from the International Tobacco Control (ITC) four country survey. *Tob Control*, 15(Suppl 3):iii34-iii41.

Bourns B, Malcomson A (2001). Economic impact analysis of the non-smoking by law on the hospitality industry in Ottawa. KPMG (http://www.tobaccoscam.ucsf.edu/pdf/087-OttawaKPMG.pdf).

Boyle P, Levin B (2008). *World Cancer Report* 2008. Lyon, International Agency for Research on Cancer.

Brandt AM (2007). The cigarette century: the rise, fall, and deadly persistence of the product that defined America. New York, NY, Basic Rooks

Braverman MT, Aarø LE, Hetland J (2008). Changes in smoking among restaurant and bar employees following Norway's comprehensive smoking ban. *Health Promot Int*, 23(1):5-15.

Brenner H, Mielck A (1992). Smoking prohibition in the workplace and smoking cessation in the Federal Republic of Germany. *Prev Med*, 21(2):252-261.

Brown DC, Pereira L, Garner JB (1982). Cancer of the cervix and the smoking husband. *Can Fam Physician*, 28:499-502.

Brownell KD, Glynn TJ, Glasgow R, et al. (1986). Interventions to prevent relapse. *Health Psychol.* 5 (Suppl):53-68.

Brownson RC, Eriksen MP, Davis R, et al. (1997). Environmental tobacco smoke: health effects and policies to reduce exposure. *Annu Rev Public Health*, 18:163-185.

Buckley JD, Harris RW, Doll R, et al. (1981). Case-control study of the husbands of women with dysplasia or carcinoma of the cervix uteri. *Lancet*, 2(8254):1010-1015.

Burch JD, Rohan TE, Howe GR, et al. (1989). Risk of bladder cancer by source and type of tobacco exposure: a case-control study. Int J Cancer, 44(4):622-628.

Burns DM, Lee L, Gilpin E, et al. (1996). Cigarette smoking behaviour in the United States. In: Changes in cigarette-related disease risks and their implication for prevention and control. Smoking and Tobacco Control Monograph 8. Bethesda, MD, U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute. NIH Pub. No. 974213; 13-112.

Burns EK, Levinson A, Lezotte D, et al. (2007). Differences in smoking duration between Latinos and Anglos. Nicotine *Tob Res*, 9(7):731-737.

Butler T, Richmond R, Belcher J, et al. (2007). Should smoking be banned in prisons? *Tob Control*, 16(5):291-293.

California Department of Health Services (2006). California tobacco control update 2006: the social norm change approach. Sacramento, California, California Department of Health Services (http://www.dhs.ca.gov/tobacco/).

California Department of Public Health (2007). Tobacco control section (http://www.cdph.ca.gov/programs/Tobacco/Pages/default.aspx).

California Environmental Protection Agency (Cal EPA) (1997). Health effects of exposure to environmental tobacco smoke, California Environmental Protection Agency, Office of Environmental Health Hazard Assessment (http://www.oehha.org/air/environmental_tobacco/finalets.html).

California Environmental Protection Agency: Air Resources Board (2005). Proposed identification of environmental tobacco smoke as a toxic air contaminant. Tobacco control. Surveys and program evaluations from outside UCSF. Sacramento, CA, California Environmental Protection Agency. (http://repositories.cdlib.org/context/tc/article/1194/type/pdf/viewcontent/)

Cameron L, Williams J (2001). Cannabis, alcohol and cigarettes: substitutes or complements? *Econ Rec*, 77(236):19-34.

Cameron M, Wakefield M, Trotter L, et al. (2003). Exposure to secondhand smoke at work: a survey of members of the Australian Liquor, Hospitality and Miscellaneous Workers Union. Aust N Z J Public Health, 27(5):496-501

Carmella SG, Han S, Fristad A, et al. (2003). Analysis of total 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) in human urine. Cancer Epidemiol Biomarkers Prev, 12(11 Pt 1):1257-1261.

Celermajer DS, Adams M, Clarkson P, et al. (1996). Passive smoking and impaired endothelium-dependent arterial dilatation in healthy young adults. N Engl J Med, 334(3):150-154.

Centers for Disease Control and Prevention (1999). Decline in cigarette consumption following implementation of a comprehensive tobacco prevention and education program-Oregon, 1996-1998. MMWR Morb Mortal Wkly Rep, 48(7):140-143.

Centers for Disease Control and Prevention (2005a). Annual smoking-attributable mortality, years of potential life lost, and productivity losses--United States, 1997-2001. MMWR Morb Mortal Wkly Rep, 54(25):628.

Centers for Disease Control and Prevention (2005b). Tobacco use among students aged 13-15 years--Philippines, 2000 and 2003. MMWR Morb Mortal Wkly Rep, 54(4):94-97.

Centers for Disease Control and Prevention (2006a). State-specific prevalence of current cigarette smoking among adults and secondhand smoke rules and policies in homes and workplaces--United States, 2005. MMWR Morb Mortal Wkly Rep, 55(42):1148-1151.

Centers for Disease Control and Prevention (2006b). Tobacco use among students aged 13-15 years--Kurdistan Region, Iraq, 2005. MMWR Morb Mortal Wkly Rep, 55(20):556-559

Centers for Disease Control and Prevention (2007a). State-specific prevalence of smoke-free home rules-United States, 1992-2003. *MMWR Morb Mortal Wkly Rep*, 56(20):501-504.

Centers for Disease Control and Prevention (2007b). Reduced secondhand smoke exposure after implementation of a comprehensive statewide smoking ban-New York, June 26, 2003-June 30, 2004. MMWR Morb Mortal Wkly Rep. 56(28):705-708.

Centers for Disease Control and Prevention (2007c). Decline in smoking prevalence -New York City, 2002-2006. MMWR Morb Mortal Wkly Rep, 56(24):604-608.

Centers for Disease Control and Prevention (2008). Global Youth Tobacco Surveillance, 2000-2007. *MMWR Morb Mortal Wkly Rep*, 57(SS-1):1-32.

Cesaroni G, Forastiere F, Agabiti N, *et al.* (2008). Effect of the Italian smoking ban on population rates of acute coronary events. *Circulation*, 117(9):1183-1188.

Chaloupka FJ (1992). Clean indoor air laws, addiction and cigarette smoking. *Appl Econ*, 24:193-205.

Chaloupka FJ, Saffer H (1992). Clean indoor air laws and the demand for cigarettes. *Contemporary Policy Issues*, 72-83.

Chaloupka FJ, Grossman M (1996). Price, tobacco control policies and youth smoking. National Bureau of Economic Research. NBER Working Paper Series. Working Paper #5740 (http://www.nber.org/).

Chaloupka FJ, Wechsler H (1997). Price, tobacco control policies and smoking among young adults. *J Health Econ*, 16(3):359-373.

Chaloupka F, Warner K (2000). The economics of smoking. In: Cuyler A, Newhouse J, eds. *The handbook of health economics*. New York, Elsevier Science: 29:1539-1627.

Chaloupka FJ, Pacula RL, Farrelly MC, et al. (1999). Do higher cigarette prices encourage youth to use marijuana? Cambridge, MA, National Bureau of Economic Research.

Chang JS, Selvin S, Metayer C, et al. (2006). Parental smoking and the risk of childhood leukemia. *Am J Epidemiol*, 163(12):1091-1100.

Chapman S (2000). Banning smoking outdoors is seldom ethically justifiable. *Tob Control*, 9(1):95-97.

Chapman S (2007). Public health advocacy and tobacco control: Making smoking history. Oxford, Blackwell Publishing.

Chapman S, Borland R, Scollo M, et al. (1999). The impact of smoke-free workplaces on declining cigarette consumption in Australia and the United States. Am J Public Health, 89(7):1018-1023.

Charlton A, While D (1994). Smoking prevalence among 16-19-year-olds related to staff and student smoking policies in sixth forms and further education. *Health Educ J*, 53(1):28-39.

Chen R, Tavendale R, Tunstall-Pedoe H (2002). Measurement of passive smoking in adults: self-reported questionnaire or serum cotinine? *J Cancer Epidemiol Prev*, 7(2):85-95.

Chen YC, Su HJ, Guo YL, et al. (2005). Interaction between environmental tobacco smoke and arsenic methylation ability on the risk of bladder cancer. Cancer Causes Control, 16(2):75-81.

Cheng YJ, Hildesheim A, Hsu M, et al. (1999). Cigarette smoking, alcohol consumption and risk of nasopharyngeal carcinoma in Taiwan. Cancer Causes Control, 10(3):201-207.

Chuang JC, Callahan PJ, Lyu CW, et al. (1999). Polycyclic aromatic hydrocarbon exposures of children in low-income families. *J Expo Anal Environ Epidemiol*, 9(2):85-98.

Clark PI, Schooley MW, Pierce B, et al. (2006). Impact of home smoking rules on smoking patterns among adolescents and young adults. *Prev Chronic Dis*, 3(2):1-13.

Clarke V, White V, Hill D (1994). School structural and policy variables associated with student smoking. *Tob Control*, 3:339-346.

Claxton LD, Morin RS, Hughes TJ, et al. (1989). A genotoxic assessment of environmental tobacco smoke using bacterial bioassays. *Mutat Res*, 222(2):81-99.

Coghlin J, Hammond S, Gann P (1989). Development of epidemiologic tools for measuring environmental tobacco smoke exposure. *Am J Epidemiol*, 130(4):696-704.

Cogliano VJ, Baan R, Straif K, et al. (2004). The science and practice of carcinogen identification and evaluation. *Environ Health Perspect*, 112(13):1269-1274.

Coker AL, Rosenberg AJ, McCann MF, et al. (1992). Active and passive cigarette smoke exposure and cervical intraepithelial neoplasia. Cancer Epidemiol Biomarkers Prev, 1(5):349-356

Coker AL, Bond S, Williams A, et al. (2002). Active and passive smoking, high-risk human papillomaviruses and cervical neoplasia. *Cancer Detect Prev.* 26(2):121-128.

Collins DJ, Lapsley HM (1996). The social costs of drug abuse in Australia in 1988 and 1992. Canberra, Australia, Australian Government Publishing Service (http://www.health.gov.au/internet/main/publishing.nsf/Content/health-publith-publicat-document-mono30-cnt.htm/).

Collins DJ, Lapsley HM (2002). Counting the cost: estimates of the social costs of drug abuse in Australia in 1998-9. National Drug Strategy Monograph Series (No 49). Canberra, Australia, Australian Government Publishing Service.

Collins DJ, Lapsley HM (2008). The costs of tobacco, alcohol and illicit drug abuse to Australian society in 2004/2005. National Drug Strategy Monograph Series (No 64). Canberra, Australia, Australian Government Publishing Service.

Conference Board of Canada (1997). Smoking and the bottom line: the costs of smoking in the workplace. Ottawa, Health Canada (http://www.hc-sc.gc.ca/hl-vs/pubs/tobac-tabac/bottomline-bilan/index-eng.php).

Conley Thomson C, Siegel M, Winickoff J, et al. (2005). Household smoking bans and adolescents' perceived prevalence of smoking and social acceptability of smoking. *Prev Med*, 41(2):349-356.

Connolly GN, Carpenter C, Hillel R, et al. (2005). Evaluation of the Massachusetts smoke-free workplace law. San Francisco, CA, University of California (http://repositories.cdlib.org/context/tc/article/1180/type/pdf/viewcontent/).

Connolly GN, Carpenter CM, Travers M, et al. (2006). How smoke-free laws improve air quality: a global study of Irish pubs. Harvard School of Public Health Division of Public Health Practice, Roswell Park Cancer Institute Department of Health Behavior, Health Service Executive-West Environmental Health Department, Research Institute for a Tobacco Free Society, Office of Tobacco Control (http://www.tobaccofreeair.org/downloads/Irishstudy 3.13.3 final5.pdf).

Coppotelli HC, Orleans CT (1985). Partner support and other determinants of smoking cessation maintenance among women. *J Consult Clin Psychol*, 53(4):455-460.

Corti B, Holman C, Donovan R, et al. (1997). Warning: attending a sport, racing or arts venue may be beneficial to your health. Aust N Z J Public Health, 21(4 Spec No):371-376.

Coultas DB, Peake GT, Samet JM (1989). Questionnaire assessment of lifetime and recent exposure to environmental tobacco smoke. *Am J Epidemiol*, 130(2):338-347.

Cowling DW, Bond P (2005). Smoke-free laws and bar revenues in California--the last call. *Health Econ*, 14(12):1273-1281.

Crawford FG, Mayer J, Santella RM, *et al.* (1994). Biomarkers of environmental tobacco smoke in preschool children and their mothers. *J Natl Cancer Inst*, 86(18):1398-1402.

Cremieux P, Ouellette P (2001). Actual and perceived impacts of tobacco regulation on restaurants and firms. *Tob Control*, 10:33-37.

Cummings KM, Markello SJ, Mahoney MC, et al. (1989). Measurement of lifetime exposure to passive smoke. Am J Epidemiol, 130(1):122-132

Cummings KM, Pechacek T, Shopland D (1994). The illegal sale of cigarettes to U.S. minors: estimates by state. *Am J Public Health*, 84(2):300-302.

Daisey JM, Mahanama KR, Hodgson AT (1998). Toxic volatile organic compounds in simulated environmental tobacco smoke: emission factors for exposure assessment. *J Expo Anal Environ Epidemiol*, 8(3):313-334.

Darling H, Reeder A (2003). Smoke-free schools? Results of a secondary school smoking policies survey 2002. *N Z Med J*, 116(1180):U560.

Darling H, Reeder A, Waa A (2006). Implementation of the smoke-free environments act (2003 Amendments) in New Zealand primary schools. *Aust N Z J Public Health*, 30(1):87.

De Bruin YB, Carrer P, Jantunen M, et al. (2004). Personal carbon monoxide exposure levels: contribution of local sources to exposures an inconvironment concentrations in Milan. *J Expo Anal Environ Epidemiol*, 14(4):312-322.

Dearlove JV, Bialous SA, Glantz SA (2002). Tobacco industry manipulation of the hospitality industry to maintain smoking in public places. *Tob Control*, 11(2):94-104.

Dee TS (1999). The complementarity of teen smoking and drinking. *J Health Econ*, 18(6):769-793.

Deloitte & Touche LLP (2003). The impact of non-smoking ordinances on restaurant financial performance. Washington, DC, Deloitte & Touche LLP.

DeMarini DM (2004). Genotoxicity of tobacco smoke and tobacco smoke condensate: a review. *Mutat Res.* 567(2-3):447-474.

den Exter Blokland EA, Hale WW, III, Meeus W, et al. (2006). Parental anti-smoking socialization. Associations between parental anti-smoking socialization practices and early adolescent smoking initiation. Eur Addict Res, 12(1):25-32.

DerSimonian R, Laird N (1986). Meta-analysis in clinical trials. *Control Clin Trials*, 7(3):177-188

Diez-Roux AV, Nieto F, Comstock G, et al. (1995). The relationship of active and passive smoking to carotid atherosclerosis 12-14 years later. *Prev Med*, 24(1):48-55.

Dinno A, Glantz S (2007). Clean indoor air laws immediately reduce heart attacks. *Prev Med*, 45(1):9-11.

Doll R, Hill AB (1950). Smoking and carcinoma of the lung; preliminary report. *Br Med J*, 2(4682):739-748.

Doll R, Hill AB (1954). The mortality of doctors in relation to their smoking habits; a preliminary report. *Br Med J*, 1(4877):1451-1455.

Doll R, Peto R (1978). Cigarette smoking and bronchial carcinoma: dose and time relationships among regular smokers and lifelong non-smokers. *J Epidemiol Community Health*, 32(4):303-313.

Donnan GA, McNeil J, Adena M, et al. (1989). Smoking as a risk factor for cerebral ischaemia. *Lancet*, 2(8664):643-647.

Drobes DJ, Tiffany ST (1997). Induction of smoking urge through imaginal and in vivo procedures: physiological and self-report manifestations. *J Abnorm Psychol*, 106(1):15-25.

Dubois G (2005). [Prevention of air pollution by indoor tobacco smoke in France]. *Bull Acad Natl Med*, 189(5):803-812.

Dunnick JK, Elwell MR, Huff J, et al. (1995). Chemically induced mammary gland cancer in the National Toxicology Program's carcinogenesis bioassay. Carcinogenesis, 16(2):173-179.

Eadie D, Heim D, Macaskill S, et al. (2008). A qualitative analysis of compliance with smokefree legislation in community bars in Scotland: implications for public health. *Addiction*, 103(6):1019-1026.

Eagan TM, Hetland J, Aaro LE (2006). Decline in respiratory symptoms in service workers five months after a public smoking ban. *Tob Control*, 15(3):242-246.

Eaton DK, Kann L, Kinchen S, et al. (2006). Youth risk behavior surveillance--United States, 2005. MMWR Surveill Summ, 55(5):1-108

Edwards R, Bullen C, O'Dea D, et al. (2007). After the smoke has cleared: evaluation of the impact of a new smokefree law (A report commissioned by the New Zealand Ministry of Health). Tob Control, 1-10.

Edwards R, Thomson G, Wilson N, et al. (2008). After the smoke has cleared: evaluation of the impact of a new national smoke-free law in New Zealand. *Tob Control*, 17(1):e2.

Eisner MD, Smith AK, Blanc PD (1998). Bartenders' respiratory health after establishment of smoke-free bars and taverns. *JAMA*, 280(22):1909-1914.

el-Bayoumy K, Chae YH, Upadhyaya P, et al. (1995). Comparative tumorigenicity of benzo[a]pyrene, 1-nitropyrene and 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine administered by gavage to female CD rats. Carcinogenesis, 16(2):431-434.

Ellingsen DG, Fladseth G, Daae H, et al. (2006). Airborne exposure and biological monitoring of bar and restaurant workers before and after the introduction of a smoking ban. J Environ Monit, 8(3):362-368.

Emaus A, Lochen M, Hoifodt R (2001). [Application of tobacco smoking regulations in restaurants in Tromso 1998]. *Tidsskr Nor Laegeforen*, 121(4):410-412.

Emmanuel SC, Phe A, Chen A (1988). The impact of the anti-smoking campaign in Singapore. *Singapore Med J*, 29(3):233-239.

Emmons KM, Hammond S, Fava J, et al. (2001). A randomized trial to reduce passive smoke exposure in low-income households with young children. *Pediatrics*, 108(1):18-24.

Emont SL, Choi WS, Novotny TE, *et al.* (1993). Clean indoor air legislation, taxation, and smoking behavior in the United States: an ecological analysis. *Tob Control*, 2:13-17.

Engelen M, Farrelly M, Hyland A (2006). The health and economic consequences of New York's Clean Indoor Air Act. Albany, NY, New York State Department of Health http://www.health.state.ny.us/prevention/tobaccocontrol/).

Enstrom JE (1979). Rising lung cancer mortality among nonsmokers. *J Natl Cancer Inst*, 62(4):755-760.

Enstrom JE, Kabat GC (2003). Environmental tobacco smoke and tobacco related mortality in a prospective study of Californians, 1960-98. *Br Med J*, 326(7398):1057-1061.

Environics Research Group (2001). *Public support for international efforts to control tobacco: A survey in five countries.* Toronto, Canada, Environics Research Group.

Eriksen MP, Gottlieb NH (1998). A review of the health impact of smoking control at the workplace. *Am J Health Promot*, 13(2):83-104.

Eriksen MP, Chaloupka FJ (2007). The economic Impact of Clean Indoor Air Laws. *CA Cancer J Clin*, 57:367-378.

Escobedo LG, Peddicord JP (1996). Smoking prevalence in US birth cohorts: the influence of gender and education. *Am J Public Health*, 86(2):231-236.

Escoffery C, Kegler M, Butler S (2008). Formative research on creating smoke-free homes in rural communities. *Health Educ* Res,1-11.

European Commission (2007). Attitudes of Europeans toward tobacco [Special Eurobarometer 272c], European Commission (http://ec.europa.eu/health/ph_publication/eb_health_en.pdf).

Evans WN, Farrelly MC, Montgomery E (1999). Do workplace smoking bans reduce smoking? *Am Econ Rev*, 89:728-747.

Ezra D (1994). Sticks and stones can break my bones, but tobacco smoke can kill me: can we protect children from parents that smoke? Saint Louis Univ Public Health Law Rev, 13:547-590

Fagerstrom KO, Schneider NG (1989). Measuring nicotine dependence: a review of the Fagerstrom Tolerance Questionnaire. *J Behav Med*, 12(2):159-182.

Fang J, Gan DK, Zheng SH, et al. (2006). [A case-control study of the risk factors for lung cancer among Chinese women who have never smoked]. Wei Sheng Yan Jiu, 35(4):464-467.

Farkas AJ, Pierce JP, Zhu SH, et al. (1996). Addiction versus stages of change models in predicting smoking cessation. *Addiction*, 91(9):1271-1280.

Farkas AJ, Gilpin EA, Distefan JM, *et al.* (1999). The effects of household and workplace smoking restrictions on quitting behaviours. *Tob Control*, 8(3):261-265.

Farkas AJ, Gilpin EA, White MM, *et al.* (2000). Association between household and workplace smoking restrictions and adolescent smoking. *JAMA*, 284(6):717-722.

Farrelly MC, Evans WN, Sfekas AE (1999). The impact of workplace smoking bans: results from a national survey. *Tob Control*, 8(3):272-277.

Farrelly MC, Bray J, Zarkin G, et al. (2001). The joint demand for cigarettes and marijuana: evidence from the national household surveys on drug abuse. *J Health Econ*, 20(1):51-68.

Farrelly MC, Niederdeppe J, Yarsevich J (2003). Youth tobacco prevention mass media campaigns: past, present, and future directions. *Tob Control*, 12(Suppl 1):i35-i47.

Farrelly MC, Nonnemaker J, Chou R, et al. (2005). Changes in hospitality workers' exposure to secondhand smoke following the implementation of New York's smoke-free law. *Tob Control*, 14(4):236-241.

Federico B, Costa G, Kunst A (2007). Educational inequalities in initiation, cessation, and prevalence of smoking among 3 Italian birth cohorts. *Am J Public Health*, 97(5):838-845.

Fernandez E, Fu M, Martinez C, et al. (2008). Secondhand smoke in hospitals of Catalonia (Spain) before and after a comprehensive ban on smoking at the national level. *Prev Med*, 47(6):624-628.

Fernandez E, Fu M, Pascual J, et al. (2009). Impact of the Spanish smoking law on exposure to second-hand smoke and respiratory health in hospitality workers: a prospective cohort study. PLoS ONE, 4(1):e4244.

Fernando D, Fowles J, Woodward A, et al. (2007). Legislation reduces exposure to second-hand tobacco smoke in New Zealand bars by about 90%. Tob Control, 16(4):235-238

Feyerabend C, Russell MA (1980). Assay of nicotine in biological materials: sources of contamination and their elimination. *J Pharm Pharmacol*, 32(3):178-181.

Fichtenberg CM, Glantz SA (2002). Effect of smoke-free workplaces on smoking behaviour: systematic review. *Br Med J*, 325(7357):188-191

Fidan F, Guven H, Eminoglu O, et al. (2005). Turkish coffeehouse kahvehane is an important tobacco smoke exposure area in Turkey. *J Toxicol Environ Health A*. 68(16):1371-1377.

Fielding JE (1991). Smoking control at the workplace. *Annu Rev Public Health*, 12:209-234

Filippini G, Maisonneuve P, McCredie M, et al. (2002). Relation of childhood brain tumors to exposure of parents and children to tobacco smoke: the SEARCH international case-control study. Surveillance of Environmental Aspects Related to Cancer in Humans. *Int J Cancer*, 100(2):206-213.

Fisher LB, Winickoff JP, Camargo CA Jr, et al. (2007). Household smoking restrictions and adolescent smoking. *Am J Health Promot*, 22(1):15-21.

Flanders WD, Lally CA, Zhu BP, et al. (2003). Lung cancer mortality in relation to age, duration of smoking, and daily cigarette consumption: results from Cancer Prevention Study II. Cancer Res. 63(19):6556-6562.

Flight Attendants Medical Research Institute (1991). (http://www.famri.org/intro/).

Fong GT, Hyland A, Borland R, et al. (2006). Reductions in tobacco smoke pollution and increases in support for smoke-free public places following the implementation of comprehensive smoke-free workplace legislation in the Republic of Ireland: findings from the ITC Ireland/UK Survey. *Tob Control*, 15(Suppl 3):iii51-iii58.

Fontham ET, Correa P, Reynolds P, *et al.* (1994). Environmental tobacco smoke and lung cancer in nonsmoking women. A multicenter study. *JAMA*, 271(22):1752-1759.

Ford L (2008). Thank you for smoking -- in a hut. *Chicago Tribune*.

Franco-Marina F. (2008). Personal communication to E. Khaykin and J. Samet.

Franco-Marina F, Villalba-Caloca J, Corcho-Berdugo A (2006). Role of active and passive smoking on lung cancer etiology in Mexico City. *Salud Publica Mex*, 48(Suppl 1):S75-S82.

Frieden TR, Mostashari F, Kerker B, et al. (2005). Adult tobacco use levels after intensive tobacco control measures: New York City, 2002-2003. Am J Public Health, 95(6):1016-1023.

Friend K, Levy D (2002). Reductions in smoking prevalence and cigarette consumption associated with mass-media campaigns. *Health Educ Res*, 17(1):85-98.

Friis RH, Safer AM (2005). Analysis of responses of Long Beach, California residents to the smoke-free bars law. *Public Health*, 119(12):1116-1121.

Fukuda K, Shibata A (1990). Exposure-response relationships between woodworking, smoking or passive smoking, and squamous cell neoplasms of the maxillary sinus. *Cancer Causes Control*, 1(2):165-168.

Galan I, Mata N, Estrada C, et al. (2007). Impact of the "tobacco control law" on exposure to environmental tobacco smoke in Spain. BMC Public Health, 7(147):224.

Galeone D, Laurendi G, Vasselli S, *et al.* (2006). Preliminary effects of Italy's ban on smoking in enclosed public places. *Tob Control*, 15(2):143.

Gallet GA, Eastman HS (2007). The impact of smoking bans on alcohol demand. *Soc Sci J*, 44(4):664-676.

Gallicchio L, Kouzis A, Genkinger J, et al. (2006). Active cigarette smoking, household passive smoke exposure, and the risk of developing pancreatic cancer. Prev Med, 42(3):200-205.

Gallus S, Pacifici R, Colombo P, et al. (2006). Prevalence of smoking and attitude towards smoking regulation in Italy, 2004. Eur J Cancer Prev, 15(1):77-81.

Gallus S, Zuccaro P, Colombo P, et al. (2007). Smoking in Italy 2005-2006: effects of a comprehensive National Tobacco Regulation. *Prev Med*, 45(2-3):198-201.

Gammon MD, Eng SM, Teitelbaum SL, *et al.* (2004). Environmental tobacco smoke and breast cancer incidence. *Environ Res*, 96(2):176-185.

Garfinkel L (1981). Time trends in lung cancer mortality among nonsmokers and a note on passive smoking. *J Natl Cancer Inst*, 66(6):1061-1066.

Garland C, Barrett-Connor E, Suarez L, et al. (1985). Effects of passive smoking on ischemic heart disease mortality of nonsmokers. A prospective study. Am J Epidemiol, 121(5):645-650.

Garvey AJ, Bliss RE, Hitchcock JL, et al. (1992). Predictors of smoking relapse among self-quitters: a report from the Normative Aging Study. Addict Behav, 17(4):367-377.

Gasparrini A, Gorini G, Marcolina D, et al. (2006). [Second-hand smoke exposure in Florence and Belluno before and after the Italian smoke-free legislation]. *Epidemiol Prev*, 30(6):348-351.

Gee IL, Watson A, Carrington J, et al. (2006). Second-hand smoke levels in UK pubs and bars: do the English Public Health White Paper proposals go far enough? J Public Health (Oxf), 28(1):17-23.

Gehrman CA, Hovell MF (2003). Protecting children from environmental tobacco smoke (ETS) exposure: a critical review. *Nicotine Tob Res*, 5(3):289-301.

Georgiadis P, Stoikidou M, Topinka J, et al. (2001a). Personal exposures to PM(2.5) and polycyclic aromatic hydrocarbons and their relationship to environmental tobacco smoke at two locations in Greece. J Expo Anal Environ Epidemiol. 11(3):169-183.

Georgiadis P, Topinka J, Stoikidou M, et al. (2001b). Biomarkers of genotoxicity of air pollution (the AULIS project): bulky DNA adducts in subjects with moderate to low exposures to airborne polycyclic aromatic hydrocarbons and their relationship to environmental tobacco smoke and other parameters. *Carcinogenesis*, 22(9):1447-1457.

Giles-Corti B, Clarkson J, Donovan R, *et al.* (2001). Creating smoke-free environments in recreational settings. *Health Educ Behav*, 28(3):341-351.

Gillespie J, Milne K, Wilson N (2005). Secondhand smoke in New Zealand homes and cars: exposure, attitudes, and behaviours in 2004. *N Z Med J*, 118(1227):U1782.

Gilpin E, Pierce J (1994). Measuring smoking cessation: problems with recall in the 1990 California Tobacco Survey. *Cancer Epidemiol Biomarkers Prev*, 3(7):613-617.

Gilpin EA, Pierce JP (2002a). Demographic differences in patterns in the incidence of smoking cessation: United States 1950-1990. *Ann Epidemiol*, 12(3):141-150.

Gilpin EA, Pierce J (2002b). The California tobacco control program and potential harm reduction through reduced cigarette consumption in continuing smokers. *Nicotine Tob Res.* 4(Suppl 2):S157-S166.

Gilpin EA, Pierce J, Cavin S, et al. (1994). Estimates of population smoking prevalence: self vs proxy reports of smoking status. Am J Public Health, 84(10):1576-1579.

Gilpin EA, Pierce JP, Farkas AJ (1997). Duration of smoking abstinence and success in quitting. *J Natl Cancer Inst*, 89(8):572-576.

Gilpin EA, White MM, Farkas AJ, et al. (1999). Home smoking restrictions: which smokers have them and how they are associated with smoking behavior. *Nicotine Tob Res*, 1(2):153-162.

Gilpin EA, Stillman F, Hartman A, et al. (2000). Index for US state tobacco control initial outcomes. *Am J Epidemiol*, 152(8):727-738.

Gilpin EA, Emery S, Farkas A, et al. (2001). The California tobacco control program: a decade of progress. Results from the California Tobacco Survey, 1990-1999. La Jolla, CA, University of California, San Diego (http://ssdc.ucsd.edu/).

Gilpin EA, Farkas A, Emery S, et al. (2002). Clean indoor air: advances in California, 1990-1999. *Am J Public Health*, 92(5):785-791.

Gilpin EA, White M, White V, et al. (2003). Tobacco control successes in California: A focus on young people, results from the California Tobacco surveys, 1990-2002. La Jolla, CA; University of California, San Diego (http://libraries.ucsd.edu/ssds/pub/CTS/cpc00007/2002FINAL RPT.pdf).

Gilpin EA, Lee L, Pierce J, et al. (2004). Support for protection from secondhand smoke: California 2002. *Tob Control*, 13(1):96.

Gilpin EA, Messer K, Pierce JP (2006). Population effectiveness of pharmaceutical aids for smoking cessation: what is associated with increased success? *Nicotine Tob Res*, 8(5):661-669.

Glantz SA (2007). Commentary: Assessing the effects of the Scottish Smokefree Law--the placebo effect and the importance of obtaining unbiased data. *Int J Epidemiol*, 36(1):155-156.

Glantz SA, Smith LRA (1994). The effect of ordinances requiring smoke-free restaurants on restaurant sales. *Am J Public Health*, 84(7):1081-1085.

Glantz SA, Parmley WW (1995). Passive smoking and heart disease. Mechanisms and risk. *JAMA*, 273(13):1047-1053.

Glantz SA, Smith LRA (1997). The effect of ordinances requiring smoke-free restaurants and bars on revenues: A follow up. *Am J Public Health*, 87(10):1687-1693.

Glantz SA, Charlesworth A (1999). Tourism and hotel revenues before and after passage of smoke-free restaurant ordinances. *JAMA*, 281(20):1911-1918.

Glantz SA, Balbach ED (2000). Tobacco war: inside the California battles. Berkeley, CA, University of California Press (http://escholarship.cdlib.org/ucpress/tobacco-war.xml).

Glantz SA, Wilson-Loots R (2003). No association of smoke-free ordinances with profits from bingo and charitable games in Massachusetts. *Tob Control*, 12:411-413.

Glantz SA, Alamar B (2005). Correction for Mandel L, Alamar BC, Glantz SA: Smokefree law did not affect revenue from gaming in Delaware. *Tob Control*, 14:360.

Glantz SA, Barnes DE, Bero L, *et al.* (1995). Looking through a keyhole at the tobacco industry. The Brown and Williamson documents. *JAMA*, 274(3):219-224.

Glaser SL, Keegan TH, Clarke CA, et al. (2004). Smoking and Hodgkin lymphoma risk in women United States. *Cancer Causes Control*, 15(4):387-397.

Glasgow RE, Cummings KM, Hyland A (1997). Relationship of worksite smoking policy to changes in employee tobacco use: findings from COMMIT. Community Intervention Trial for Smoking Cessation. *Tob Control*, 6(Suppl 2):S44-S48.

Glasgow RE, Boles S, Lichtenstein E, et al. (2004). Adoption, reach, and implementation of a novel smoking control program: analysis of a public utility-research organization partnership. *Nicotine Tob Res*, 6(2):269-274.

Goldstein AO, Sobel RA (1998). Environmental tobacco smoke regulations have not hurt restaurant sales in North Carolina. *N C Med J*, 59(5):284-287.

Gonzales M, Malcoe L, Kegler M, et al. (2006). Prevalence and predictors of home and automobile smoking bans and child environmental tobacco smoke exposure: a cross-sectional study of U.S.- and Mexico-born Hispanic women with young children. *BMC Public Health*, 6:265.

Goodman MT, Tung KH (2003). Active and passive tobacco smoking and the risk of borderline and invasive ovarian cancer (United States). Cancer Causes Control, 14(6):569-577.

Goodman P, Agnew M, McCaffrey M, et al. (2007). Effects of the Irish smoking ban on respiratory health of bar workers and air quality in Dublin pubs. Am J Respir Crit Care Med, 175:840-845.

Gorini G, Fondelli M, Lopez M, et al. (2004). [Environmental tobacco smoke exposure in public places in Florence, Italy]. *Epidemiol Prev*, 28(2):94-99.

Gower KB, Burns DM, Shanks TG, et al. (2000). Workplace smoking restrictions, rules about smoking in the home, and attitudes toward smoking restrictions in public places. In: Shopland DR, Hobart R, Burns DM, et al., eds., State and local legislative action to reduce tobacco use. Bethesda, MD, U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute. Smoking and Tobacco Control Monograph No. 11: 187-340.

Greenberg RA, Strecher V, Bauman K, et al. (1994). Evaluation of a home-based intervention program to reduce infant passive smoking and lower respiratory illness. *J Behav Med*, 17(3):273-290.

Griesbach D, Inchley J, Currie C (2002). More than words? The status and impact of smoking policies in Scottish schools. *Health Promot Int*, 17(1):31-41.

Groner JA, Ahijevych K, Grossman LK, *et al.* (2000). The impact of a brief intervention on maternal smoking behavior. *Pediatrics*, 105(1 Pt 3):267-271.

Groner JA, Hoshaw-Woodard S, Koren G, et al. (2005). Screening for children's exposure to environmental tobacco smoke in a pediatric primary care setting. Arch Pediatr Adolesc Med, 159(5):450-455.

GTSS Collaborative Group (2006). A cross country comparison of exposure to secondhand smoke among youth. *Tob Control*, 15(Suppl 2): ii4-ii19.

Gupta R, Dwyer J (2001). Focus groups with smokers to develop a smoke-free home campaign. *Am J Health Behav*, 25(6):564-571.

Hackshaw A (2003). Passive smoking: paper does not diminish conclusion of previous reports. *Br Med J*, 327(7413):501-502.

Hackshaw AK, Law MR, Wald NJ (1997). The accumulated evidence on lung cancer and environmental tobacco smoke. *Br Med J*, 315(7114):980-988.

Hahn E, Rayens M, York N, et al. (2006). Effects of a smoke-free law on hair nicotine and respiratory symptoms of restaurant and bar workers. J Occup Environ Med, 48(9):906-913

Hallamore C (2006). Smoking and the bottom line: updating the costs of smoking in the workplace. Ontario, Canada, Conference Board of Canada (http://www.conferenceboard.ca/documents.asp?rnext=1754).

Halterman JS, Fagnano M, Conn K, et al. (2006). Do parents of urban children with persistent asthma ban smoking in their homes and cars? *Ambul Pediatr*, 6(2):115-119.

Hamajima N, Hirose K, Tajima K, et al. (2002). Alcohol, tobacco and breast cancer-collaborative reanalysis of individual data from 53 epidemiological studies, including 58,515 women with breast cancer and 95,067 women without the disease. *Br J Cancer*, 87(11):1234-1245.

Hammond D, McDonald P, Fong G, et al. (2004). The impact of cigarette warning labels and smoke-free bylaws on smoking cessation: evidence from former smokers. *Can J Public Health*, 95(3):201-204.

Hammond D, Fong G, Zanna M, et al. (2006). Tobacco denormalization and industry beliefs among smokers from four countries. *Am J Prev Med*, 31(3):225-232.

Hammond EC, Horn D (1954). The relationship between human smoking habits and death rates. *JAMA*, 155:1316-1328.

Hammond SK (1999). Exposure of U.S. workers to environmental tobacco smoke. *Environ Health Perspect*, 107(Suppl 2):329-340.

Hammond SK, Emmons KM (2005). Inmate exposure to secondhand smoke in correctional facilities and the impact of smoking restrictions. *J Expo Anal Environ Epidemiol*, 15(3):205-211.

Harakeh Z, Scholte R, de Vries H, et al. (2005). Parental rules and communication: their association with adolescent smoking. *Addiction*, 100(6):862-870.

Hassan MM, Abbruzzese J, Bondy M, et al. (2007). Passive smoking and the use of noncigarette tobacco products in association with risk for pancreatic cancer: a case-control study. Cancer, 109(12):2547-2556.

Hatsukami DK, Le C, Zhang Y, et al. (2006). Toxicant exposure in cigarette reducers versus light smokers. Cancer Epidemiol Biomarkers Prev, 15(12):2355-2358.

Hatziandreu EJ, Pierce J, Fiore M, et al. (1989). The reliability of self-reported cigarette consumption in the United States. *Am J Public Health*, 79(8):1020-1023.

Haw SJ, Gruer L (2007). Changes in exposure of adult non-smokers to secondhand smoke after implementation of smoke-free legislation in Scotland: national cross sectional survey. *Br Med J (Clin Res Ed)*, 335(7619):549-552.

Haw SJ, Gruer L, Amos A, *et al.* (2006). Legislation on smoking in enclosed public places in Scotland: how will we evaluate the impact? *J Public Health* (Oxf), 28(1):24-30.

Health Canada (2006). Canadian Tobacco Use Monitoring Survey (CTUMS) 2006. Ottawa, Health Canada (http://www.hc-sc.gc.ca/hl-vs/tobac-tabac/research-recherche/stat/ctums-esutc_2006_e.html).

Healton CG, Vallone D, Martyak J, et al. (2007). Using social marketing to increase the prevalence of voluntary smoking bans in the home and car. (Austin TX. SYM7C). Austin, TX, Society for Research on Nicotine and Tobacco.

Hecht SS (2002). Human urinary carcinogen metabolites: biomarkers for investigating tobacco and cancer. *Carcinogenesis*, 23(6):907-922.

Hecht SS (2003). Tobacco carcinogens, their biomarkers and tobacco-induced cancer. *Nat Rev Cancer*, 3(10):733-744.

Hecht SS, Hoffmann D (1988). Tobaccospecific nitrosamines, an important group of carcinogens in tobacco and tobacco smoke. *Carcinogenesis*, 9(6):875-884.

Hedley AJ, Mcghee S, Repace J, et al. (2006). Risks for heart disease and lung cancer from passive smoking by workers in the catering industry. *Toxicol Sci*, 90(2):539-548.

Helakorpi S, Martelin T, Torppa J, et al. (2004). Did Finland's tobacco control act of 1976 have an impact on ever smoking? An examination based on male and female cohort trends. *J Epidemiol Community Health*, 58(8):649-654.

Helakorpi S, Martelin T, Torppa J, et al. (2008). Impact of the 1976 tobacco control act in Finland on the proportion of ever daily smokers by socioeconomic status. *Prev Med*, (4):340-345.

Helgason AR, Lund KE (2001). Environmental tobacco smoke exposure of young children-attitudes and health-risk awareness in the Nordic countries. *Nicotine Tob Res*, 3(4):341-345.

Hellberg D, Valentin J, Nilsson S (1986). Smoking and cervical intraepithelial neoplasia. An association independent of sexual and other risk factors? *Acta Obstet Gynecol Scand*, 65(6):625-631.

Heloma A, Jaakkola M (2003). Four-year followup of smoke exposure, attitudes and smoking behaviour following enactment of Finland's national smoke-free work-place law. *Addiction*, 98(8):1111-1117.

Heloma A, Jaakkola M, Kahkonen E, et al. (2001). The short-term impact of national smoke-free workplace legislation on passive smoking and tobacco use. Am J Public Health, 91(9):1416-1418.

Henderson FW, Reid HF, Morris R, et al. (1989). Home air nicotine levels and urinary cotinine excretion in preschool children. Am Rev Respir Dis. 140(1):197-201.

Henriksen L, Jackson C (1998). Anti-smoking socialization: relationship to parent and child smoking status. *Health Commun*, 10(1):87-101.

Hernandez-Mezquita M, Barrueco M, Jimenez-Ruiz C, et al. (2000). [Extent of law compliance and anti-smoking teaching at Spanish schools]. Rev Esp Salud Pública, 74(5-6):537-547.

Hetland J, Aaro L (2005). [Smoking status, attitudes towards and reported compliance with the ban on smoking at hospitality venues - a prospective panel study]. Oslo, Norway, Statens Institutt for Rusmiddelforskning. Report No. 2 (http://www.sirus.no/files/pub/201SIRUSskrifter0205.pdf).

Hill AB (1965). The environment and disease: association or causation? *Proc R Soc Med*, 58:295-300.

Hill D, Borland R (1991). Adults' accounts of onset of regular smoking: influences of school, work, and other settings. *Public Health Rep*, 106(2):181-185.

Hill SE, Blakely T, Kawachi I, et al. (2007). Mortality among lifelong nonsmokers exposed to secondhand smoke at home: cohort data and sensitivity analyses. Am J Epidemiol, 165(5):530-540.

Hinds WC (1982). Aerosol technology: properties, behavior, and measurement of airborne particles. New York, John Wiley & Sons

Hirayama T (1981). Non-smoking wives of heavy smokers have a higher risk of lung cancer: a study from Japan. *Br Med J (Clin Res Ed)*, 282(6259):183-185.

Hirayama T (1984). Cancer mortality in nonsmoking women with smoking husbands based on a large-scale cohort study in Japan. *Prev Med*, 13(6):680-690.

Hirayama T (1985). Passive smoking--a new target of epidemiology. *Tokai J Exp Clin Med*, 10(4):287-293.

Hirose K, Tajima K, Hamajima N, et al. (1995). A large-scale, hospital-based case-control study of risk factors of breast cancer according to menopausal status. *Jpn J Cancer Res*, 86(2):146-154.

Hirose K, Tajima K, Hamajima N, *et al.* (1996). Subsite (cervix/endometrium)-specific risk and protective factors in uterus cancer. *Jpn J Cancer Res*, 87(9):1001-1009.

Holm AL, Davis RM (2004). Clearing the airways: advocacy and regulation for smoke-free airlines. *Tob Control*, 13(Suppl 1):i30-i36.

Hopkins DP, Briss PA, Ricard CJ, et al. (2001). Reviews of evidence regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke. Am J Prev Med, 20(2 Suppl):16-66.

Horwitz MB, Hindi-Alexander M, Wagner TJ (1985). Psychosocial mediators of abstinence, relapse, and continued smoking: a one-year follow-up of a minimal intervention. *Addict Behav*, 10(1):29-39.

Hovell MF, Zakarian J, Matt G, et al. (2000). Effect of counseling mothers on their children's exposure to environmental tobacco smoke: randomised controlled trial. *Br Med J*, 321(72 57):337-342.

Hovell MF, Meltzer S, Wahlgren D, et al. (2002). Asthma management and environmental tobacco smoke exposure reduction in Latino children: a controlled trial. *Pediatrics*, 110(5): 946-956.

Howard G, Burke G, Szklo M, et al. (1994). Active and passive smoking are associated with increased carotid wall thickness. The atherosclerosis risk in communities study. *Arch Intern Med*, 154(11):1277-1282.

Howard G, Wagenknecht L, Burke G, et al. (1998). Cigarette smoking and progression of atherosclerosis: the Atherosclerosis Risk in Communities (ARIC) study. *JAMA*, 279(2):119-124.

Howell F (2004). Ireland's workplaces, going smoke free. *Br Med J*, 328(7444):847-848.

Hu J, Mao Y, Ugnat AM (2000). Parental cigarette smoking, hard liquor consumption and the risk of childhood brain tumors--a case-control study in northeast China. *Acta Oncol*, 39(8):979-984.

Hughes SC, Corcos IA, Hofstetter CR, et al. (2008). Children's exposure to secondhand smoke at home in Seoul, Korea. Asian Pac J Cancer Prev. 9(3):491-495.

Hurley SF, McNeil JJ, Donnan GA, et al. (1996). Tobacco smoking and alcohol consumption as risk factors for glioma: a case-control study in Melbourne, Australia. J Epidemiol Community Health, 50(4):442-446.

Hyland A, Cummings K (1999). Restaurateur reports of the economic impact of the New York City Smoke-free Air Act. *J Public Health Manag Pract*, 5(1):37-42.

Hyland A, Cummings K, Nauenberg E (1999). Analysis of taxable sales receipts: was New York City's Smoke-free Air Act bad for business? *J Public Health Manag Pract*, 5(1):14-21.

Hyland A, Travers M, Dresler C, *et al.* (2008a). A 32-country comparison of tobacco smoke derived particle levels in indoor public places. *Tob Control*, 17(3):159-165.

Hyland A, Higbee C, Hassan L, et al. (2008b). Does smoke-free Ireland have more smoking inside the home and less in pubs than the United Kingdom? Findings from the International Tobacco Control Policy Evaluation Project. Eur J Public Health, 18(1):63-65.

Hyland A, Higbee C, Travers MJ, et al. (2009). Smoke-free homes and smoking cessation and relapse in a longitudinal population of adults. Nicotine Tob Res..

IARC (1986). IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol. 38, Tobacco Smoke and Involuntary Smoking. Lyon, IARC Press.

IARC (2004). IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol. 83, Tobacco Smoke and Involuntary Smoking. Lyon, International Agency for Research on Cancer.

IARC (2008). IARC Handbooks of Cancer Prevention, Tobacco Control, Vol. 12: Methods for Evaluating Tobacco Control Policies. Lyon, International Agency for Research on Cancer.

Institute of Medicine (2001). Clearing the smoke: assessing the science base for tobacco harm reduction. Washington DC, National Academy Press.

Invernizzi G, Ruprecht A, Mazza R, et al. (2002). [Real-time measurement of indoor particulate matter originating from environmental tobacco smoke: a pilot study]. *Epidemiol Prev*, 26(1):30-34.

Jaakkola MS, Jaakkola JJ (1997). Assessment of exposure to environmental tobacco smoke. *Eur Respir J.* 10(10):2384-2397.

Jackson C, Henriksen L (1997). Do as I say: parent smoking, antismoking socialization, and smoking onset among children. *Addict Behav*, 22(1):107-114.

Jackson C, Dickinson D (2003). Can parents who smoke socialise their children against smoking? Results from the smoke-free kids intervention trial. *Tob Control*, 12(1):52-59.

Jarvis MJ, Goddard E, Higgins V, et al. (2000). Children's exposure to passive smoking in England since the 1980s: cotinine evidence from population surveys. *Br Med J*, 321(7257):343-345.

Javitz HS, Zbikowski S, Swan G, et al. (2006). Financial burden of tobacco use: an employer's perspective. *Clin Occup Environ Med*, 5(1):9-29, vii.

Jee SH, Ohrr H, Kim IS (1999). Effects of husbands' smoking on the incidence of lung cancer in Korean women. *Int J Epidemiol*, 28(5):824-828.

Jenkins RA, Maskarinec M, Counts R, et al. (2001). Environmental tobacco smoke in an unrestricted smoking workplace: area and personal exposure monitoring. J Expo Anal Environ Epidemiol, 11(5):369-380.

Jha P, Chaloupka F (1999). Curbing the epidemic. Washington, DC, The World Bank (http://www1.worldbank.org/tobacco/book/html/cover2a.html).

Ji M, Hofstetter C, Hovell M, et al. (2005). Smoking cessation patterns and predictors among adult Californians of Korean descent. *Nicotine Tob Res*, 7(1):59-69.

Jiang X, Yuan JM, Skipper PL, et al. (2007). Environmental tobacco smoke and bladder cancer risk in never smokers of Los Angeles County. Cancer Res, 67(15):7540-7545.

Johansson A, Hermansson G, Ludvigsson J (2004). How should parents protect their children from environmental tobacco-smoke exposure in the home? *Pediatrics*, 113(4):e291-e295.

Johnson KC (2005). Accumulating evidence on passive and active smoking and breast cancer risk. *Int J Cancer*, 117(4):619-628.

Johnsson T, Tuomi T, Riuttala H, et al. (2006). Environmental tobacco smoke in Finnish restaurants and bars before and after smoking restrictions were introduced. Ann Occup Hyg, 50(4):331-341.

Jones K, Wakefield M, Turnball D (1999). Attitudes and experiences of restaurateurs regarding smoking bans in Adelaide, South Australia. *Tob Control*, 8(1):62-66.

Joseph AM, Knapp JM, Nichol KL, *et al.* (1995). Determinants of compliance with a national smoke-free hospital standard. *JAMA*, 274(6):491-494.

Junker MH, Danuser B, Monn C, et al. (2001). Acute sensory responses of nonsmokers at very low environmental tobacco smoke concentrations in controlled laboratory settings. *Environ Health Perspect*, 109(10):1045-1052.

Juster HR, Loomis BR, Hinman TM, et al. (2007). Declines in hospital admissions for acute myocardial infarction in New York State after implementation of a comprehensive smoking ban. Am J Public Health, 97(11):2035-2039.

Kaiserman MJ (1997). The cost of smoking in Canada, 1991. Chronic Dis Can, 18(1):13-19.

Kallio K, Jokinen E, Raitakari O, et al. (2007). Tobacco smoke exposure is associated with attenuated endothelial function in 11-year-old healthy children. *Circulation*, 115(25):3205-3212.

Kandel DB, Wu P (1995). The contributions of mothers and fathers to the inter-generational transmission of cigarette smoking in adolescence. *J Res Adolesc*, 5:225-252.

Kang JM, Jiang Y, Lin XG, et al. (2007). [Study on the level of tobacco-generated smoke in several restaurants and bars in Beijing, China]. Zhonghua liu xing bing xue za zhi, 28(8):738-741.

Keeler TE, Hu T, Barnett P, et al. (1993). Taxation, regulation, and addiction: a demand function for cigarettes based on time-series evidence. J Health Econ. 12(1):1-18.

Kegler MC, Malcoe LH (2002). Smoking restrictions in the home and car among rural Native American and white families with young children. *Prev Med*, 35(4):334-342.

Kegler MC, Escoffery C, Groff A, et al. (2007). A qualitative study of how families decide to adopt household smoking restrictions. Fam Community Health, 30(4):328-341.

Kegler MC, Escoffery C, Butler S (2008). A qualitative study on establishing and enforcing smoking rules in family cars. Nicotine *Tob Res*, 10(3):493-497.

Kellogg JH (1922). *Tobaccoism, or, how tobacco kills*. Battle Creek, Michigan, The Modern Medicine Publishing Company.

Kennett L. (1987). GI: *The American soldier in World War II*. New York, NY, Scribner.

Kessler G (2006). Civil Action #99-2496. USA vs. Philip Morris Inc.

Khuder SA, Milz S, Jordan T, et al. (2007). The impact of a smoking ban on hospital admissions for coronary heart disease. *Prev Med*, 45(1):3-8.

King G, Mallett R, Kozlowski L, et al. (2005). Personal space smoking restrictions among African Americans. *Am J Prev Med*, 28(1):33-40.

Kinne S, Kristal A, White E, et al. (1993). Worksite smoking policies: their population impact in Washington State. Am J Public Health, 83(7):1031-1033.

Klein EG, Forster JL, McFadden B, et al. (2007). Minnesota tobacco-free park policies: Attitudes of the general public and park officials. *Nicotine Tob Res*, 9(Suppl 1):S49-S55.

Klepeis NE, Ott WR, Repace JL (1999). The effect of cigar smoking on indoor levels of carbon monoxide and particles. *J Expo Anal Environ Epidemiol*, 9(6):622-635.

Klepeis NE, Ott WR, Switzer P (2007). Realtime measurement of outdoor tobacco smoke particles. *J Air Waste Manag Assoc*, 57(5):522-534

Klerman L (2004). Protecting children: reducing their environmental tobacco smoke exposure. *Nicotine Tob Res*, 6(Suppl 2):S239-S253.

Kluger R (1996). Ashes to ashes: America's hundred-year cigarette war, the public health, and the unabashed triumph of Philip Morris. New York, NY, Alfred A. Knopf.

Kodl MM, Mermelstein R (2004). Beyond modeling: parenting practices, parental smoking history, and adolescent cigarette smoking. *Addict Behav*, 29(1):17-32.

Komro KA, McCarty MC, Forster JL, *et al.* (2003). Parental, family, and home characteristics associated with cigarette smoking among adolescents. *Am J Health Promot*, 17(5):291-299.

KPMG (1998). The impact of California's smoking ban on bars, taverns and night clubs: a survey of owners and managers. Washington, DC, The American Beverage Institute (http://www.tobaccoscam.ucsf.edu/pdf/068-KPMG.pdf).

KPMG (2001). Proposed smoking ban: impacts on Hong Kong hospitality businesses. Hong Kong Catering Industry Association (http://www.tobaccoscam.ucsf.edu/pdf/089-HongKongHospitality.pdf).

Kumar R, O'Malley P, Johnston L (2005). School tobacco control policies related to students' smoking and attitudes toward smoking: national survey results, 1999-2000. Health Educ Behav, 32(6):780-794.

Kunyk D, Els C, Predy G, et al. (2007). Development and introduction of a comprehensive tobacco control policy in a Canadian regional health authority. Prev Chronic Dis, 4:A30.

Kurahashi N, Inoue M, Liu Y, et al. (2008). Passive smoking and lung cancer in Japanese non-smoking women: a prospective study. *Int J Cancer*, 122(3):653-657.

Kuusimaki L, Peltonen K, Vainiotalo S (2007). Assessment of environmental tobacco smoke exposure of Finnish restaurant workers, using 3-ethenylpyridine as marker. *Indoor Air*, 17(5):394-403.

LaKind JS, Jenkins RA, Naiman DQ, et al. (1999). Use of environmental tobacco smoke constituents as markers for exposure. *Risk Anal*, 19(3):359-373.

Lal A, Siahpush M, Scollo M (2003). The economic impact of smoke-free policies on sales in restaurants and cafés in Victoria. *Aust N Z J Public Health*, 27(5):557-558.

Lal A, Siahpush M, Scollo M (2004). The economic impact of smoke-free legislation on sales turnover in restaurants and pubs in Tasmania. *Tob Control*, 13(4):454-455.

Lal A, Siahpush M (2008). The effect of smokefree policies on electronic gaming machine expenditure in Victoria, Australia. *J Epidemiol Community Health*, 62(1):11-15.

Lam TH, Janghorbani M, Hedley AJ, et al. (2002). Public opinion on smoke-free policies in restaurants and predicted effect on patronage in Hong Kong. *Tob Control*, 11(3):195-200.

Laugesen M, Swinburn B (2000). New Zealand's tobacco control programme 1985-1998. *Tob Control*, 9(2):155-162.

Lawn S, Pols R (2005). Smoking bans in psychiatric inpatient settings? A review of the research. *Aust N Z J Psychiatry*, 39(10):866-885

Lazcano-Ponce E, Benowitz N, Sanchez-Zamorano L, et al. (2007). Secondhand smoke exposure in Mexican discotheques. *Nicotine Tob Res*, 9(10):1021-1026.

Leaderer BP, Hammond SK (1991). Evaluation of vapour-phase nicotine and respirable suspended particle mass as markers for environmental tobacco smoke. *Environ Sci Technol*, 25(4):770-777.

Leatherdale ST, Smith P, Ahmed R (2008). Youth exposure to smoking in the home and in cars: how often does it happen and what do youth think about it? *Tob Control*, 17(2):86-92.

Lee CW, Kahende J (2007). Factors associated with successful smoking cessation in the United States, 2000. *Am J Public Health*, 97(8):1503-1509.

Lee K, Hahn E, Riker C, et al. (2007). Immediate impact of smoke-free laws on indoor air quality. South Med J, 100(9):885-889.

Lee LY, Gerhardstein DC, Wang AL, *et al.* (1993). Nicotine is responsible for airway irritation evoked by cigarette smoke inhalation in men. *J Appl Physiol*, 75(5):1955-1961.

Lee PN (1988). *Misclassification of smoking habits and passive smoking*. Berlin, Springer Verlag.

Lee PN (1998). Difficulties in assessing the relationship between passive smoking and lung cancer. *Stat Methods Med Res*, 7(2):137-163.

Lee PN, Chamberlain J, Alderson MR (1986). Relationship of passive smoking to risk of lung cancer and other smoking-associated diseases. *Br J Cancer*, 54(1):97-105.

Lee SC, Chan LY, Chiu MY (1999). Indoor and outdoor air quality investigation at 14 public places in Hong Kong. *Environ Int*, 22:443-450.

Lemstra M, Neudorf C, Opondo J (2008). Implications of a public smoking ban. *Can J Public Health*, 99(1):62-65.

Leung GM, Ho LM, Lam TH (2004). Secondhand smoke exposure, smoking hygiene, and hospitalization in the first 18 months of life. *Arch Pediatr Adolesc Med*, 158(7):687-693.

Levin ML, Goldstein H, Gerhardt PR (1950). Cancer and tobacco smoking. *JAMA*, 143:336-338.

Levy DT, Friend KB (2003). The effects of clean indoor air laws: what do we know and what do we need to know? *Health Educ Res*, 18(5):592-609

Lewit EM, Hyland A, Kerrebrock N, et al. (1997). Price, public policy, and smoking in young people. *Tob Control*, 6(Suppl 2):S17-S24.

Li D, Wang M, Dhingra K, et al. (1996). Aromatic DNA adducts in adjacent tissues of breast cancer patients: clues to breast cancer etiology. Cancer Res, 56(2):287-293.

Li WM, Lee SC, Chan LY (2001). Indoor air quality at nine shopping malls in Hong Kong. *Sci Total Environ*, 273(1-3):27-40.

Lichtenstein E, Andrews J, Lee M, et al. (2000). Using radon risk to motivate smoking reduction: evaluation of written materials and brief telephone counseling. *Tob Control*, 9(3):320-326.

Lilla C, Verla-Tebit E, Risch A, et al. (2006). Effect of NAT1 and NAT2 genetic polymorphisms on colorectal cancer risk associated with exposure to tobacco smoke and meat consumption. Cancer Epidemiol Biomarkers Prev. 15(1):99-107.

Lissowska J, Brinton LA, Zatonski W, et al. (2006). Tobacco smoking, NAT2 acetylation genotype and breast cancer risk. *Int J Cancer*, 119(8):1961-1969.

Lissowska J, Brinton LA, Garcia-Closas M (2007). Re: more data regarding the effects of passive smoking on breast cancer risk among younger women. *Int J Cancer*, 120(11):2517-2518.

Little CC (1961). Some phases of the problem of smoking and lung cancer. *N Engl J Med*, 264:1241-1245.

Lo AC, Soliman AS, El-Ghawalby N, *et al.* (2007). Lifestyle, occupational, and reproductive factors in relation to pancreatic cancer risk. *Pancreas*, 35(2):120-129.

Lofroth G (1989). Environmental tobacco smoke: overview of chemical composition and genotoxic components. *Mutat Res*, 222(2):73-80.

Lombard HL, Doering CR (1980). Cancer studies in Massachusetts. 2. Habits, characteristics and environment of individuals with and without cancer. *CA Cancer J Clin*, 30(2):115-122.

Longo DR, Brownson R, Johnson J, et al. (1996). Hospital smoking bans and employee smoking behavior: Results of a national survey. *JAMA*, 275(16):1252-1257.

Longo DR, Johnson J, Kruse R, et al. (2001). A prospective investigation of the impact of smoking bans on tobacco cessation and relapse. *Tob Control*, 10(3):267-272.

Lopez MJ, Nebot M, Albertini M, et al. (2008). Secondhand smoke exposure in hospitality venues in Europe. *Environ Health Perspect*, 116:1469-1472.

Lubin JH (1999). Estimating lung cancer risk with exposure to environmental tobacco smoke. *Environ Health Perspect*, 107(Suppl 6):879-883.

Luk R, Ferrence R, Gmel G (2006). The economic impact of a smoke-free bylaw on restaurant and bar sales in Ottawa, Canada. *Addiction*, 101:738-745.

Lund KE (2006). The introduction of smokefree hospitality venues in Norway. Impact on revenues, frequency of patronage, satisfaction and compliance. Oslo, Norway, Norwegian Institute for Alcohol and Drug Research (SIRUS) (http://www.sirus.no/files/pub/375/ SIRUSskrifter0206eng.pdf). Lund KE, Rise J (2004). Mediekampanje om roykfrie serveringssteder varen 2004 [with English language summary on p8]. Oslo, Norwegian Institute for Alcohol and Drug Research (SIRUS) (http://www.sirus.no/).

Lund KE, Helgason AR (2005). Environmental tobacco smoke in Norwegian homes, 1995 and 2001: changes in children's exposure and parent's attitudes and health risk awareness. *Eur J Public Health*, 15(2):123-127.

Lund KE, Lund M (2006). The impact of smoke-free hospitality venues in Norway. Eurohealth, 12(4):22-24 (http://www.euro.who.int/Document/Obs/Eurohealth12 4.pdf).

Lund M, Lindbak R (2007). Norwegian tobacco statistics 1973-2006. Oslo, Norwegian Institute for Alcohol and Drug Research (SIRUS) (http://www.sirus.no/internett/forsiden?factory=index).

Lund M, Lund K, Rise J, et al. (2005). Smoke-free bars and restaurants in Norway. Oslo, Norway, The Norwegian Institute for Alcohol and Drug Research (http://www.globalink.org/documents/2005smokefreebarsandrestaurantsinNorway.pdf).

Lundborg P (2007). Does smoking increase sick leave? Evidence using register data on Swedish workers. *Tob Control*, 16(2):114-118.

Lung SC, Wu MJ, Lin CC (2004). Customers' exposure to PM2.5 and polycyclic aromatic hydrocarbons in smoking/nonsmoking sections of 24-h coffee shops in Taiwan. *J Expo Anal Environ Epidemiol*, 14(7):529-535.

Lushchenkova O, Fernandez E, Lopez M, et al. (2008). [Secondhand smoke exposure in Spanish adult non-smokers following the introduction of an anti-smoking law]. Rev Esp Cardiol. 61(7):687-694.

Mackenzie CR, Campbell GD (1963). Cigarettes and disease. *S Afr Med J*, 37:974-975.

Maclure M, Katz RB, Bryant MS, et al. (1989). Elevated blood levels of carcinogens in passive smokers. *Am J Public Health*, 79(10):1381-1384.

MacMahon B, Trichopoulos D, Cole P, et al. (1982). Cigarette smoking and urinary estrogens. N Engl J Med, 307(17):1062-1065.

Madden D (2003). The cost of employee smoking in Ireland. Dublin, University College Dublin (http://otc.ie/Uploads/Download%20Da vid%20Maddens%20Presentation.pdf).

Mandel LL, Alamar BC, Glantz SA (2005). Smoke-free law did not affect revenue from gaming in Delaware. *Tob Control*, 14(1):10-12.

Manor R (1997). Where there's smoke: there'll soon be a lounge, at Lambert Airport. *St. Louise Post*-Dispatch. A1-A6.

Mao Y, Hu J, Semenciw R, et al. (2002). Active and passive smoking and the risk of stomach cancer, by subsite, in Canada. Eur J Cancer Prev, 11(1):27-38.

Markowitz S (2000). The price of alcohol, wife abuse, and husband abuse. South Econ J, 67(2):279-303.

Marlatt GA, Curry S, Gordon JR (1988). A longitudinal analysis of unaided smoking cessation. *J Consult Clin Psychol*, 56(5):715-720

Martin J, George R, Andrews K, et al. (2006). Observed smoking in cars: a method and differences by socioeconomic area. *Tob Control.* 15(5):409-411.

Martinez-Donate AP, Hovell MF, Hofstetter CR, et al. (2005). Smoking, exposure to secondhand smoke, and smoking restrictions in Tijuana, Mexico. *Rev Panam Salud Publica*, 18(6):412-417.

Martinez-Donate AP, Hovell MF, Hofstetter CR, et al. (2007). Correlates of home smoking bans among Mexican-Americans. *Am J Health Promot*, 21(4):229-236.

Maskarinec MP, Jenkins RA, Counts RW, et al. (2000). Determination of exposure to environmental tobacco smoke in restaurant and tavern workers in one US city. J Expo Anal Environ Epidemiol. 10(1):36-49.

Matanoski G, Kanchanaraksa S, Lantry D, et al. (1995). Characteristics of nonsmoking women in NHANES I and NHANES I epidemiologic follow-up study with exposure to spouses who smoke. Am J Epidemiol, 142(2):149-157.

Matt GE, Wahlgren DR, Hovell MF, et al. (1999). Measuring environmental tobacco smoke exposure in infants and young children through urine cotinine and memory-based parental reports: empirical findings and discussion. *Tob Control*, 8(3):282-289.

Matt GE, Quintana PJ, Hovell MF, et al. (2004). Households contaminated by environmental tobacco smoke: sources of infant exposures. *Tob Control*, 13(1):29-37.

Mauderly JL, Gigliotti AP, Barr EB, et al. (2004). Chronic inhalation exposure to mainstream cigarette smoke increases lung and nasal tumor incidence in rats. *Toxicol Sci*, 81(2):280-292.

Maziak W, Ali R, Fouad M, et al. (2008). Exposure to secondhand smoke at home and in public places in Syria: a developing country's perspective. *Inhal Toxicol*, 20(1):17-24.

Mcghee SM, Hedley AJ, Ho LM (2002). Passive smoking and its impact on employers and employees in Hong Kong. *Occup Environ Med*, 59(12):842-846.

McKee M, Gilmore A, Novotny T (2003). Smoke free hospitals. *Br Med J*, 326(7396):941-942.

McMillen RC, Winickoff JP, Klein JD, et al. (2003). US adult attitudes and practices regarding smoking restrictions and child exposure to environmental tobacco smoke: changes in the social climate from 2000-2001. *Pediatrics*, 112:e55-e60.

McMorrow MJ, Foxx RM (1983). Nicotine's role in smoking: an analysis of nicotine regulation. *Psychol Bull*, 93(2):302-327.

McMullen KM, Brownson R, Luke D, et al. (2005). Strength of clean indoor air laws and smoking related outcomes in the USA. *Tob Control*, 14(1):43-48.

McNabola A, Broderick B, Johnston P, et al. (2006). Effects of the smoking ban on benzene and 1,3-butadiene levels in pubs in Dublin. J Environ Sci Health A Tox Hazard Subst Environ Eng. 41(5):799-810.

Medical Research Council (MRC) (1957). Tobacco smoking and cancer of the lung. *Br Med J.* 1:1523-1524.

Menegaux F, Steffen C, Bellec S, et al. (2005). Maternal coffee and alcohol consumption during pregnancy, parental smoking and risk of childhood acute leukaemia. Cancer Detect Prev. 29(6):487-493.

Menegaux F, Ripert M, Hemon D, et al. (2007). Maternal alcohol and coffee drinking, parental smoking and childhood leukaemia: a French population-based case-control study. *Pediatr Perinat Epidemiol*, 21(4):293-299.

Menzies D, Nair A, Williamson P, et al. (2006). Respiratory symptoms, pulmonary function, and markers of inflammation among bar workers before and after a legislative ban on smoking in public places. *JAMA*, 296(14):1742-1748.

Mermelstein R, Lichtenstein E, McIntyre K (1983). Partner support and relapse in smoking-cessation programs. *J Consult Clin Psychol*, 51(3):465-466.

Merom D, Rissel C (2001). Factors associated with smoke-free homes in NSW: results from the 1998 NSW Health Survey. *Aust N Z J Public Health*, 25(4):339-345.

Messer K, Trinidad D, Al-Delaimy W, et al. (2008a). Smoking cessation rates in the United States: a comparison of young adult and older smokers. *Am J Public Health*, 98(2):317-322.

Messer K, Mills A, White M, et al. (2008b) The effect of smoke-free homes on smoking behavior in the United States. Am J Prev Med, 35(3):210-216.

Metzger KB, Mostashari F, Kerker B (2005). Use of pharmacy data to evaluate smoking regulations' impact on sales of nicotine replacement therapies in New York City. *Am J Public Health*, 95(6):1050-1055.

Michnovicz JJ, Hershcopf RJ, Naganuma H, et al. (1986). Increased 2-hydroxylation of estradiol as a possible mechanism for the anti-estrogenic effect of cigarette smoking. N Engl J Med, 315(21):1305-1309.

Miller C (2002). Evaluation of the South Australian smoke-free homes and cars project. In: The Cancer Council South Australia. Tobacco Control Research and Evaluation Report, 1998-2001. Adelaide, Tobacco Control Research Evaluation Unit, The Cancer Council South Australia (http://www.cancersa.org.au/cms_resources/documents/TCRE/TCRE_Eval Report Vol1 Ch9.pdf).

Mills CA, Porter MM (1950). Tobacco smoking habits and cancer of the mouth and respiratory system. *Cancer Res*, 10(9):539-542.

Ministry of Health (1990). Smoke-free law in New Zealand (http://www.moh.govt.nz/smoke freelaw).

Ministry of Health (2003). Il rapporto sull'impatto della legge 16 gennaio 2003, n. art.51 "Tutela della salute dei non fumatori" (http://www.ministerosalute.it/resources/static/primopiano/255/conferenzaFumo.pdf).

Ministry of Health (2007). New Zealand tobacco use survey 2006. Wellington, Ministry of Health.

Ministry of Health & Care Services (1973). Act No.14 of 9 March 1973 relating to prevention of the harmful effects of tobacco. Oslo, Norway (http://www.regieringen.no/en/dep/hod/Subjects/The-Department-of-Public-Health/Act-No-14-of-9-March-1973-relating-to-Prevention-of-the-Harmful-Effects-of-Tobacco.html?id=224962).

Moher M, Ley K, Lancaster T (2005). Workplace interventions for smoking cessation (Review). *The Cochrane Library*, 2:1-79.

Moore L, Roberts C, Tudor-Smith C (2001). School smoking policies and smoking prevalence among adolescents: multilevel analysis of cross-sectional data from Wales. *Tob Control*, 10(2):117-123.

Moore M, Hrushow A (2004). Increasing compliance with California Smoke-free workplace law in San Francisco: Final evaluation report. San Francisco, CA

Moore RS, Lee JP, Antin TM, et al. (2006). Tobacco free workplace policies and low socioeconomic status female bartenders in San Francisco. J Epidemiol Community Health, 60(Suppl 2):51-56.

Morabia A, Bernstein M, Heritier S, et al. (1996). Relation of breast cancer with passive and active exposure to tobacco smoke. Am J Epidemiol, 143(9):918-928.

Morozumi R, li M (2006). The impact of smokefree workplace policies on smoking behavior in Japan. *Appl Econ*, 13:549-555.

Moshammer H, Neuberger M, Nebot M (2004). Nicotine and surface of particulates as indicators of exposure to environmental tobacco smoke in public places in Austria. *Int J Hyg Environ Health*, 207(4):337-343.

Moskowitz JM, Lin Z, Hudes E (2000). The impact of workplace smoking ordinances in California on smoking cessation. *Am J Public Health*, 90(5):757-761.

Mudarri D (1994). The costs and benefits of smoking restrictions: an assessment of the Smoke-Free Environment Act of 1993. (H.R. 3434) Washington, DC, US Environmental Protection Agency.

Mulcahy M, Evans D, Hammond S, et al. (2005). Secondhand smoke exposure and risk following the Irish smoking ban: an assessment of salivary cotinine concentrations in hotel workers and air nicotine levels in bars. *Tob Control*, 14(6):384-388.

Muller FH (1939). Tabakmissbrauch und lungercarcinoma. Z Krebsforsch, 49:57-85.

Mumford EA, Levy DT, Romano EO (2004). Home smoking restrictions. Problems in classification. *Am J Prev Med*, 27(2):126-131.

Nagao M, Ushijima T, Wakabayashi K, et al. (1994). Dietary carcinogens and mammary carcinogenesis. Induction of rat mammary carcinomas by administration of heterocyclic amines in cooked foods. Cancer, 74(3 Suppl): 1063-1069.

Nagle AL, Schofield MJ, Redman S (1996). Smoking on hospital grounds and the impact of outdoor smoke-free zones. *Tob Control*, 5(3):199-204.

Nardini S, Pacifici R, Mortali C, *et al.* (2003). A survey on policies of smoking control in Italian hospitals. *Monaldi Arch Chest Dis*, 59(4):310-313.

National Cancer Institute (1999). Health effects of exposure to environmental tobacco smoke: the report of the California Environmental Protection Agency. Smoking and Tobacco Control Monograph 10. NIH Pub. No. 99-4645. Bethesda, MD, U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute.

National Cancer Institute (2000). State and local legislative action to reduce tobacco use. Smoking and Tobacco Control Monograph 11. NIH Pub. No. 00-4804. Bethesda, MD, U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute.

National Health and Medical Research Counciln (NHMRC) (1987). Effects of passive smoking on health. Report of the NHMRC working party on the effect of passive smoking on health. Canberra, Australian Government Publishing Service.

National Health and Medical Research Council (NHMRC) (1997). The health effects of passive smoking: A scientific information paper. Canberra, Australian Government Publishing Service.

National Health Service (2006). The smoking, health and social care (Scotland) Act 2005 (Commencement No.4), Order 2006 (http://www.legislation.gov.uk/legislation/scotland/ssi2006/20060121.htm).

National Libraries of Medicine (2007). The Reports of the Surgeon General: chronological listing by primary document type. National Libraries of Medicine (http://profiles.nlm.nih.gov/NN/ListByDate.html).

National Research Council (1986). Environmental tobacco smoke: measuring exposures and assessing health effects. Washington, DC, National Academy Press. (http://www.nap.edu/openbook.php?isbn=0309037301).

Navas-Acien A, Peruga A, Breysse P, et al. (2004). Secondhand tobacco smoke in public places in Latin America, 2002-2003. *JAMA*, 291(22):2741-2745.

Nebot M, Lopez M, Gorini G, et al. (2005). Environmental tobacco smoke exposure in public places of European cities. *Tob Control*, 14(1):60-63.

New Zealand Ministry of Health (2005). The smoke is clearing: Anniversary report. Wellington, New Zealand, Ministry of Health (http://www.moh.govt.nz/publications).

Nishino Y, Tsubono Y, Tsuji I, et al. (2001). Passive smoking at home and cancer risk: a population-based prospective study in Japanese nonsmoking women. *Cancer Causes Control*, 12(9):797-802.

Noonan G (1976). Passive smoking in enclosed public places. *Med J Aust*, 2(2):68-70.

Norman GJ, Ribisl KM, Howard-Pitney B, *et al.* (1999). Smoking bans in the home and car: Do those who really need them have them? *Prev Med*, 29(6 Pt 1):581-589.

Norman GJ, Ribisl KM, Howard-Pitney B, *et al.* (2000). The relationship between home smoking bans and exposure to state tobacco control efforts and smoking behaviors. *Am J Health Promot*, 15(2):81-88.

Norway Ministry of Health and Care Services (1973). Act No. 14 of 9 March 1973 relating to prevention of the harmful effects of tobacco (http://www.regjeringen.no/en/dep/hod/Subjects/The-Department-of-Public-Health/Act-No-14-of-9-March-1973-relating-to-Prevention-of-the-Harmful-Effects-of-Tobacco.html?id=224962).

Office of Tobacco Control (2005). Smoke-free workplaces in Ireland: a one-year review. Clane, Co Kildare, Ireland, Office of Tobacco Control (http://www.otc.ie/Uploads/1_Year_Report_FA.pdf).

Office of Tobacco Control (2007). Annual Report (http://www.otc.ie/uploads/Annual_Report 2007.pdf).

Okah FA, Choi WS, Okuyemi KS, et al. (2002). Effect of children on home smoking restriction by inner-city smokers. *Pediatrics*, 109(2):244-249.

Okah FA, Okuyemi K, McCarter K, et al. (2003). Predicting adoption of home smoking restriction by inner-city black smokers. *Arch Pediatr Adolesc Med*, 157(12):1202-1205.

Osthus S, Pape H, Lund K (2007). [Smoking restrictions and smoking prevalence in high schools]. *Tidsskr Nor Laegeforen*, 127(9):1192-1194.

Otsuka R, Watanabe H, Hirata K, et al. (2001). Acute effects of passive smoking on the coronary circulation in healthy young adults. *JAMA*, 286(4):436-441.

Ott WR (1999). Mathematical models for predicting indoor air quality from smoking activity. *Environ Health Perspect*, 107(Suppl 2):375-381.

Pakko MR (2006). Smoke-free law did affect revenue from gaming in Delaware. *Tob Control*, 15(1):68-69.

Palinkas LA, Pierce J, Rosbrook BP, et al. (1993). Cigarette smoking behavior and beliefs of Hispanics in California. Am J Prev Med, 9(6):331-337.

Pang D, McNally R, Birch JM (2003). Parental smoking and childhood cancer: results from the United Kingdom Childhood Cancer Study. *Br J Cancer*, 88(3):373-381.

Parrott S, Godfrey C, Raw M (2000). Costs of employee smoking in the workplace in Scotland. *Tob Control*, 9(2):187-192.

Patten CA, Gilpin E, Cavin S, et al. (1995). Workplace smoking policy and changes in smoking behavior in California: a suggested association. *Tob Control*, 4:36-41.

Payne TJ, Smith P, Sturges L, et al. (1996). Reactivity to smoking cues: mediating roles of nicotine dependence and duration of deprivation. Addict Behav, 21(2):139-154.

Pell JP, Haw S, Cobbe S, et al. (2008). Smoke-free legislation and hospitalizations for acute coronary syndrome. *N Engl J Med*, 359(5):482-491.

Penner M, Penner S (1990). Excess insured health care costs from tobacco-using employees in a large group plan. *J Occup Med*, 32(6):521-523.

Petrakis NL, Gruenke LD, Beelen TC, et al. (1978). Nicotine in breast fluid of nonlactating women. *Science*, 199(4326):303-305.

Petrakis NL, Miike R, King EB, et al. (1988). Association of breast fluid coloration with age, ethnicity, and cigarette smoking. *Breast Cancer Res Treat*, 11(3):255-262.

Phillips K, Bentley M, Howard D, et al. (1996). Assessment of air quality in Stockholm by personal monitoring of nonsmokers for respirable suspended particles and environmental tobacco smoke. Scand J Work Environ Health, 22(Suppl 1):1-24.

Phillips LE, Longstreth WT Jr, Koepsell T, et al. (2005). Active and passive cigarette smoking and risk of intracranial meningioma. *Neuroepidemiology*, 24(3):117-122.

Pickett MS, Schober SE, Brody DJ, et al. (2006). Smoke-free laws and secondhand smoke exposure in US non-smoking adults, 1999-2002. *Tob Control*, 15(4):302-307.

Pickett W, Northrup D, Ashley M (1999). Factors influencing implementation of the legislated smoking ban on school property in Ontario. *Prev Med*, 29(3):157-164.

Picone GA, Sloan F, Trogdon JG (2004). The effect of the tobacco settlement and smoking bans on alcohol consumption. *Health Econ*, 13(10):1063-1080.

Pierce JP, Gilpin EA (1995). A historical analysis of tobacco marketing and the uptake of smoking by youth in the United States: 1890-1977. *Health Psychol*, 14(6):500-508.

Pierce JP, Leon M (2008). Effectiveness of smoke-free policies. *Lancet Oncol*, 9(7):614-615.

Pierce JP, Dwyer T, DiGiusto E, *et al.* (1987). Cotinine validation of self-reported smoking in commercially run community surveys. *J Chronic Dis*, 40(7):689-695.

Pierce JP, Fiore M, Novotny T, et al. (1989). Trends in cigarette smoking in the United States. Educational differences are increasing. *JAMA*, 261(1):56-60.

Pierce JP, Naquin M, Gilpin E, et al. (1991). Smoking initiation in the United States: a role for worksite and college smoking bans. *J Natl Cancer Inst*, 83(14):1009-1013.

Pierce J, Evans D, Farkas A, et al. (1994). Tobacco use in California. An evaluation of the Tobacco Control Program 1989-1993. San Diego, CA, La Jolla, University of California.

Pierce JP, Choi WS, Gilpin EA, et al. (1996). Validation of susceptibility as a predictor of which adolescents take up smoking in the United States. Health Psychol, 15(5):355-361.

Pierce J, Gilpin E, Emery S, et al. (1998a). Tobacco control in California: Who's winning the war? An evaluation of the tobacco control program 1989-1996. San Diego, CA, La Jolla, University of California.

Pierce JP, Gilpin E, Emery S, *et al.* (1998b). Has the California tobacco control program reduced smoking? *JAMA*, 280(10):893-899.

Pierce JP, Gilpin E, Farkas AJ (1998c). Can strategies used by statewide tobacco control programs help smokers make progress in quitting? Cancer Epidemiol Biomarkers Prev, 7(6):459-464.

Pierce JP, Farkas AJ, Gilpin EA (1998d). Beyond stages of change: the quitting continuum measures progress towards successful smoking cessation. *Addiction*, 93(2):277-286.

Pierce JP, White MM, Messer K (2009). Changing age-specific patterns of cigarette consumption in the United States, 1992-2002: Association with smoke-free homes and statelevel tobacco control activity. *Nicotine Tob Res*, 11(2):171-177.

Pikora T, Phang J, Karro J, et al. (1999). Are smoke-free policies implemented and adhered to at sporting venues? Aust N Z J Public Health, 23(4):407-409.

Pinilla J, Gonzalez B, Barber P, et al. (2002). Smoking in young adolescents: an approach with multilevel discrete choice models. *J Epidemiol Community Health*, 56(3):227-232.

Pion M, Givel M (2004). Airport smoking rooms don't work. *Tob Control*, 13(Suppl 1):i37-i40.

Pirkle JL, Flegal KM, Bernert JT, et al. (1996). Exposure of the US population to environmental tobacco smoke: the Third National Health and Nutrition Examination Survey, 1988 to 1991. JAMA, 275(16):1233-1240.

Pirkle JL, Bernert JT, Caudill SP, et al. (2006). Trends in the exposure of nonsmokers in the U.S. population to secondhand smoke: 1988-2002. Environ Health Perspect, 114(6):853-858.

Pizacani BA, Martin DP, Stark MJ, et al. (2003). Household smoking bans: which households have them and do they work? *Prev Med*, 36(1):99-107.

Pizacani BA, Martin DP, Stark MJ, et al. (2004). A prospective study of household smoking bans and subsequent cessation related behaviour: the role of stage of change. Tob Control, 13(1):23-28.

Porter RS, Gowda V, Kotchou K, et al. (2001). Tobacco use among adults: Arizona, 1996 and 1999. MMWR Morb Mortal Wkly Rep, 50(20):402-406.

Poulsen LH, Osler M, Roberts C, et al. (2002). Exposure to teachers smoking and adolescent smoking behaviour: analysis of cross sectional data from Denmark. *Tob Control*, 11(3):246-251

Prabhat J, Chaloupka F (2000). Tobacco control in developing countries. Oxford, UK, Oxford University Press (http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6VBF-45JPF9S-1&_user=382452&_rdoc=1&_fmt=&_orig=search&_sort=d&view=c&_acct=C000055308&_version=1&_urlVersion=0&_userid=382452&md5=0674ca2b4a6aa0dc70_ald70240e9641a).

Proescholdbell RJ, Chassin L, MacKinnon DP (2000). Home smoking restrictions and adolescent smoking. *Nicotine Tob Res*, 2(2):159-167.

Proescholdbell SK, Foley KL, Johnson J, et al. (2008). Indoor air quality in prisons before and after implementation of a smoking ban law. *Tob Control*, 17(2):123-127.

Pron GE, Burch JD, Howe GR, et al. (1988). The reliability of passive smoking histories reported in a case-control study of lung cancer. Am J Epidemiol, 127(2):267-273.

Pruss-Ustun A, Corvalan C, eds. (2006). Preventing disease through health environments. Towards an estimate of the environmental burden of disease. Geneva, World Health Organization.

Przewozniak K, Zatonski W (2002). Sharp decline in smoking prevalence in Polish female doctors: they reach world standards [Abstract]. Abstract book and programme of the 3rd European Conference on Tobacco or Health. Warsaw, Poland (http://eng.pw.edu.pl/).

Przewozniak K, Gumkowski J, Zagroba M, et al. (2007). ETS exposure in public places in Poland: A look at PM2.5 levels and implications for policy. Abstract book of the 4th European Conference on Tobacco or Health. Basel, Switzerland.

Przewozniak K, Gumkowski J, Zatonski W (2008). Passive smoking in Poland - exposure, health effects and prevention [In Polish with English language summary]. In: Proceedings of the National Conference on Environment and Health. Warsaw, Poland, Warsaw Technical University (http://eng.pw.edu.pl/).

Pyles MK, Mullineaux DJ, Okoli CT, et al. (2007). Economic effect of a smoke-free law in a tobacco-growing community. *Tob Control*, 16(1):66-68.

Rainio SU, Rimpela AH (2008). Home smoking bans in Finland and the association with child smoking. *Eur J Public Health*, 18(3):306-311.

Reeder A, Glasgow H (2000). Are New Zealand schools smoke-free? Results from a national survey of primary and intermediate school principals. *N Z Med J*, 113(1104):52-54.

Rees VW, Connolly GN (2006). Measuring air quality to protect children from secondhand smoke in cars. *Am J Prev Med*, 31(5):363-368.

Renaud T (2007). Smoking ban in public areas. Health Policy Monitor, (April):1-7.

Repace JL (2004). Respirable particles and carcinogens in the air of Delaware hospitality venues before and after a smoking ban. *J Occup Environ Med*, 46(9):887-905.

Repace JL, Lowrey AH (1980). Indoor air pollution, tobacco smoke, and public health. *Science*, 208(4443):464-472.

Repace JL, Lowrey AH (1990). Risk assessment methodologies for passive smoking-induced lung cancer. *Risk Anal*, 10(1):27-37.

Repace JL, Lowrey AH (1993). An enforceable indoor air quality standard for environmental tobacco smoke in the workplace. *Risk Anal*, 13(4):463-475.

Repace JL, Al-Delaimy W, Bernert J (2006a). Correlating atmospheric and biological markers in studies of secondhand tobacco smoke exposure and dose in children and adults. *J Occup Environ Med*, 48(2):181-194.

Repace JL, Hyde JN, Brugge D (2006b). Air pollution in Boston bars before and after a smoking ban. *BMC Public Health*, 6:266.

Reynolds P, Hurley S, Goldberg DE, et al. (2004). Active smoking, household passive smoking, and breast cancer: evidence from the California Teachers Study. J Natl Cancer Inst, 96(1):29-37.

Rigotti NA (1989). Trends in the adoption of smoking restrictions in public places and worksites. *N Y State J Med*, 89(1):19-26.

Rise J, Lund K (2005). Predicting children's level of exposure to environmental tobacco smoke based on two national surveys in Norway in 1995 and 2001. *Addict Behav*, 30(6):1267-1271.

Rissel C, McLellan L, Bauman A (2000). Factors associated with delayed tobacco uptake among Vietnamese/Asian and Arabic youth in Sydney, NSW. Aust N Z J Public Health, 24(1):22-28.

Roberts LM, Wakefield MA, Reynolds CS (1996). Children's exposure to environmental tobacco smoke in private vehicles. *Med J Aust*, 165(6):350.

Robinson J, Kirkcaldy A (2007). Disadvantaged mothers, young children and smoking in the home: mothers' use of space within their homes. *Health Place*, 13(4):894-903.

Roddam AW, Pirie K, Pike MC, et al. (2007). Active and passive smoking and the risk of breast cancer in women aged 36-45 years: a population based case-control study in the UK. Br J Cancer, 97(3):434-439.

Rodriguez D, Tscherne J, Udrain-McGovern J (2007). Contextual consistency and adolescent smoking: testing the indirect effect of home indoor smoking restrictions on adolescent smoking through peer smoking. *Nicotine Tob Res.* 9(11):1155-1161.

Rogers T, Feighery EC, Haladjian HH (2008). Current practices in enforcement of California laws regarding youth access to tobacco products and exposure to secondhand smoke. Sacramento, CA (http://www.publichealth.pitt.edu/docs/Land_conference_committee_testimony_3-13-08.pdf).

Roseby R, Waters E, Polnay A, et al. (2004). Family and carer smoking control programs for reducing children's exposure to ETS (Cochrane Review). Cochrane Library, Chichester: John Wiley & Sons Ltd (http://www.thecochranelibrary.com).

Rosenlund M, Berglind N, Gustavsson A, *et al.* (2001). Environmental tobacco smoke and myocardial infarction among never-smokers in the Stockholm Heart Epidemiology Program (SHEEP). *Epidemiology*, 12(5):558-564.

Ross H (2006). Economics of smoke-free policies. In: Smokefree Partnership, ed. Lifting the smokescreen: 10 reasons for a smoke free Europe. Brussels, European Respiratory Society: 1-148.

Royal College of Physicians of London (1962). Smoking and health. Summary of a report of the Royal College of Physicians of London on smoking in relation to cancer of the lung and other diseases. London, Pitman Medical Publishing Co., Ltd.

Rubenstein D, Jesty J, Bluestein D (2004). Differences between mainstream and sidestream cigarette smoke extracts and nicotine in the activation of platelets under static and flow conditions. *Circulation*, 109(1):78-83.

Ruprecht A, Boffi R, Mazza R, et al. (2006). [A comparison between indoor air quality before and after the implementation of the smoking ban in public places in Italy]. *Epidemiol Prev*, 30(6):334-337.

Ryan P, Lee MW, North B, et al. (1992). Risk factors for tumors of the brain and meninges: results from the Adelaide Adult Brain Tumor Study. Int J Cancer, 51(1):20-27.

Saloojee Y, Dagli E (2000). Tobacco industry tactics for resisting public policy on health. *Bull World Health Organ*, 78(7):902-910.

Samanic C, Kogevinas M, Dosemeci M, et al. (2006). Smoking and bladder cancer in Spain: effects of tobacco type, timing, environmental tobacco smoke, and gender. Cancer Epidemiol Biomarkers Prev, 15(7):1348-1354.

Samet JM (2006). Smoking bans prevent heart attacks. *Circulation*, 114:1450-1451.

Sandler DP, Everson RB, Wilcox AJ (1985a). Passive smoking in adulthood and cancer risk. *Am J Epidemiol*, 121(1):37-48.

Sandler DP, Everson RB, Wilcox AJ, et al. (1985b). Cancer risk in adulthood from early life exposure to parents' smoking. *Am J Public Health*, 75(5):487-492.

Sandler DP, Comstock GW, Helsing KJ, et al. (1989). Deaths from all causes in non-smokers who lived with smokers. Am J Public Health, 79(2):163-167.

Sargent RP, Shepard RM, Glantz SA (2004). Reduced incidence of admissions for myocardial infarction associated with public smoking ban: before and after study. *Br Med J*, 328(7446):977-980.

Scherer G (1999). Smoking behavior and compensation: a review of the literature. *Psychopharmacology*, 145:1-20.

Schiaffino A, Fernandez E, Kunst A, *et al.* (2007). Time trends and educational differences in the incidence of quitting smoking in Spain (1965-2000). *Prev Med*, 45(2-3):226-232.

Schick S, Gvinianidze K, Tsereteli D, et al. (2008). Pilot study of compliance with healthcare facility smoking laws in Georgia. Georgian Med News, (154):47-52.

Schofield MJ, Edwards K (1995). Community attitudes to bans on smoking in licensed premises. *Aust J Public Health*, 19(4):399-402.

Schrek R, Baker LA, Ballard GP, et al. (1950). Tobacco smoking as an etiologic factor in disease: cancer. *Cancer Res*, 10:49-58.

Schulze A, Mons M (2005). Trends in cigarette smoking initiation and cessation among birth cohorts of 1926-1970 in Germany. *Eur J Cancer Prev*, 14(5):477-483.

Scientific Committee on Tobacco and Health (1998). Report of the Scientific Committee on Tobacco and Health. United Kingdom, The Stationary Office (http://www.archive.official-documents.co.uk/).

Scollo M, Lal A (2002). Summary of studies assessing the economic impact of smokefree policies in the hospitality industry – includes studies produced to 31 August 2002. Melbourne, Australia, VicHealth Centre for Tobacco Control (http://www.vctc.org.au/publ/reports/hospitality_paper summary.pdf).

Scollo M, Lal A (2005). Summary of studies assessing the economic impact of smoke-free policies in the Hospitality industry - includes studies produced to July 2005. Melbourne, VicHealth Centre for Tobacco Control http://www.vctc.org.au/tc-res/Hospitalitysummary.pdf).

Scollo M, Lal A (2008). Summary of studies assessing the economic impact of smoke-free policies in the hospitality industry. Melbourne, Australia, VicHealth Centre for Tobacco Control (http://www.vctc.org.au/tc-res/Hospitalitysummary.pdf).

Scollo M, Lal A, Hyland A (2003). Review of the quality of studies on the economic effects of smoke-free policies on the hospitality industry. *Tob Control*, 12:13-20.

Scottish Executive (2004). A breath of fresh air for Scotland. Improving Scotland's health: the challenge. Tobacco Control Action Plan. Edinburgh, Scotland, Scottish Executive, St Andrew's House http://www.scotland.gov.uk/Publications/2004/01/18736/31541).

Scottish Government (2005). Clearing the air. Enforcement protocol (https://www.clearing theairscotland.com/fags/enforcement.html).

Scottish Government (2008). Smoke-free legislation - National compliance data: summary (http://www.clearingtheairscotland.com/latest/index.html).

Scottish Parliament. (2004). Prohibition of smoking in regulated areas (Scotland) bill (http://www.scottish.parliament.uk/business/bills/20-prohibitionSmoking/b20s2-introd.pdf).

Scottish Parliament. (2006). Prohibition of smoking in certain premises (Scotland) regulations (http://www.opsi.gov.uk/legislation/scotland/ssi2006/20060090.htm).

Sebrie EM, Glantz SA (2007). "Accommodating" smoke-free policies: tobacco industry's Courtesy of Choice programme in Latin America. *Tob Control*, 16(5):e6.

Sebrie EM, Schoj V, Glantz SA (2008). Smoke free environments in Latin America: on the road to real change? *Prev Control*, 3:21-35.

Sellstrom E, Bremberg S (2006). Is there a "school effect" on pupil outcomes? A review of multilevel studies. *J Epidemiol Community Health*, 60(2):149-155.

Semple S, Maccalman L, Naji A, et al. (2007a). Bar workers' exposure to second-hand smoke: the effect of Scottish smoke-free legislation on occupational exposure. Ann Occup Hyg, 51(7):571-580.

Semple S, Creely K, Naji A, et al. (2007b). Secondhand smoke levels in Scottish pubs: the effect of smoke-free legislation. *Tob Control*, 16(2):127-132.

Semple S, Naji A, Haw S, *et al.* Care home workers' exposure to SHS: A short summary of findings. *Occup Environ Med*, in press.

Seo DC, Torabi MR (2007). Reduced admissions for acute myocardial infarction associated with a public smoking ban: matched controlled study. *J Drug Educ*, 37(3):217-226.

Shavers VL, Fagan P, Alexander L, et al. (2006). Workplace and home smoking restrictions and racial/ethnic variation in the prevalence and intensity of current cigarette smoking among women by poverty status, TUS-CPS 1998-1999 and 2001-2002. J Epidemiol Community Health, 60(Suppl 2):ii34-ii43.

Shelley D, Nguyen N, Yerneni R, et al. (2008). Tobacco use behaviors and household smoking bans among Chinese Americans. Am J Health Promot, 22(3):168-175.

Shields M (2005). The journey to quitting smoking. *Health Rep*, 16(3):19-36.

Shields M (2007). Smoking bans: influence on smoking prevalence. *Health Rep*, 18(3):9-24.

Shiffman S, Gnys M, Richards T, et al. (1996). Temptations to smoke after quitting: a comparison of lapsers and maintainers. *Health Psychol*, 15(6):455-461.

Shopland DR, Anderson CM, Burns DM (2006). Association between home smoking restrictions and changes in smoking behaviour among employed women. *J Epidemiol Community Health*, 60(Suppl 2):ii44-ii50.

Shrubsole MJ, Gao YT, Dai Q, et al. (2004). Passive smoking and breast cancer risk among non-smoking Chinese women. *Int J Cancer*, 110(4):605-609.

Siahpush M, Scollo M (2001). Trends in public support for smoking bans in public places in Australia. *Aust N Z J Public Health*, 25(5):473.

Siahpush M, Borland R, Scollo M (2003). Factors associated with smoking cessation in a national sample of Australians. *Nicotine Tob Res.* 5(4):597-602.

Siegel M (2002). The effectiveness of statelevel tobacco control interventions: a review of program implementation and behavioral outcomes. *Annu Rev Public Health*, 23:45-71.

Siegel M, Albers A, Cheng D, et al. (2005). Effect of local restaurant smoking regulations on progression to established smoking among vouths. *Tob Control*, 14(5):300-306.

Siegel M, Albers A, Cheng D, et al. (2008). Local restaurant smoking regulations and the adolescent smoking initiation process: results of a multi-level contextual analysis among Massachusetts youths. Arch Pediatr Adolesc Med, 162(5):477-483.

Skeer M, Land M, Cheng D, et al. (2004). Smoking in Boston bars before and after a 100% smoke-free regulation: an assessment of early compliance. *J Public Health Manag Pract*, 10(6):501-507.

Skogstad M, Kjaerheim K, Fladseth G, et al. (2006). Cross shift changes in lung function among bar and restaurant workers before and after implementation of a smoking ban. Occup Environ Med, 63(7):482-487.

Slattery ML, Robison LM, Schuman KL, *et al.* (1989). Cigarette smoking and exposure to passive smoke are risk factors for cervical cancer. *JAMA*, 261(11):1593-1598.

Slattery ML, Edwards S, Curtin K, et al. (2003). Associations between smoking, passive smoking, GSTM-1, NAT2, and rectal cancer. Cancer Epidemiol Biomarkers Prev, 12(9):882-889

Slattery ML, Curtin K, Giuliano AR, et al. (2008). Active and passive smoking, IL6, ESR1, and breast cancer risk. *Breast Cancer Res Treat*, 109(1):101-111.

Sly PD, Pronczuk J (2007). Guest editorial: susceptibility of children to pollutants. *Paediatr Respir Rev.* 8(4):273-274.

Sly PD, Deverell M, Kusel MM, et al. (2007). Exposure to environmental tobacco smoke in cars increases the risk of persistent wheeze in adolescents. *Med J Aust*, 186(6):322.

Smith SJ, Deacon JM, Chilvers CE (1994). Alcohol, smoking, passive smoking and caffeine in relation to breast cancer risk in young women. UK National Case-Control Study Group. *Br J Cancer*, 70(1):112-119.

Smoke Free Partnership (2006). Lifting the smokescreen: 10 reasons for a smokefree Europe (http://www.european-lung-foundation.org/uploads/Document/WEB_CHEMIN_282_1142435970.pdf).

Sobel R (1978). They satisfy: the cigarette in American life. Garden City, NJ, Anchor Press/Doubleday.

Sobti RC, Kaur S, Kaur P, et al. (2006). Interaction of passive smoking with GST (GSTM1, GSTT1, and GSTP1) genotypes in the risk of cervical cancer in India. Cancer Genet Cytogenet, 166(2):117-123.

Soliman S, Pollack H, Warner K (2004). Decrease in the prevalence of environmental tobacco smoke exposure in the home during the 1990s in families with children. *Am J Public Health*, 94(2):314-320.

Sorahan T, Lancashire RJ (2004). Parental cigarette smoking and childhood risks of hepatoblastoma: OSCC data. *Br J Cancer*, 90(5):1016-1018.

Spencer N, Blackburn C, Bonas S, et al. (2005). Parent reported home smoking bans and toddler (18-30 month) smoke exposure: a cross-sectional survey. Arch Dis Child, 90(7):670-674.

Stayner L, Bena J, Sasco AJ, et al. (2007). Lung cancer risk and workplace exposure to environmental tobacco smoke. Am J Public Health, 97(3):545-551.

Stephen GA, McRill C, Mack M, et al. (2003). Assessment of respiratory symptoms and asthma prevalence in a U.S.-Mexico border region. *Arch Environ Health*, 58(3):156-162.

Stephens T, Pederson L, Koval J, et al. (1997). The relationship of cigarette prices and nosmoking bylaws to the prevalence of smoking in Canada. Am J Public Health, 87(9):1519-1521

Stephens T, Pederson L, Koval J, et al. (2001). Comprehensive tobacco control policies and the smoking behaviour of Canadian adults. *Tob Control*, 10(4):317-322.

Stephens YD, English G (2002). A statewide school tobacco policy review: process, results, and implications. *J Sch Health*, 72(8):334-338.

Stillman F, Navas-Acien A, Ma J, et al. (2007). Second-hand tobacco smoke in public places in urban and rural China. *Tob Control*, 16(4):229-234

Study Group on Smoking and Health (1957). Smoking and health: joint report of the study group on smoking and health. *Science*, 1129-1133.

Sumida H, Watanabe H, Kugiyama K, *et al.* (1998). Does passive smoking impair endothelium-dependent coronary artery dilation in women? *J Am Coll Cardiol*, 31(4):811-815.

Sweda L (2004). Lawsuits and secondhand smoke. *Tob Control*, 13(Suppl 1):i61-i66.

Szabo E, White V, Hayman J (2006). Can home smoking restrictions influence adolescents' smoking behaviors if their parents and friends smoke? *Addict Behav*, 31(12):2298-2303.

Tan AS, Arulanandam S, Chng CY, et al. (2000). Overview of legislation and tobacco control in Singapore. *Int J Tuberc Lung Dis*, 4(11):1002-1008.

Tang H, Cowling D, Lloyd J, et al. (2003). Changes of attitudes and patronage behaviors in response to a smoke-free bar law. Am J Public Health, 93(4):611-617.

Tang H, Cowling D, Stevens C, et al. (2004). Changes of knowledge, attitudes, beliefs, and preference of bar owner and staff in response to a smoke-free bar law. *Tob Control*, 13(1):87-89.

Tauras JA (2005). Can public policy deter smoking escalation among young adults? *J Policy Anal Manage*, 24(4):771-784.

Tauras JA, Chaloupka FJ (1999a). Price, clean indoor air laws, and cigarette smoking: evidence from longitudinal data for young adults, National Bureau of Economic Research. NBER Working Paper Series. Working Paper #6937 (http://www.nber.org/).

Tauras JA, Chaloupka FJ (1999b). Determinants of smoking cessation: an analysis of young adult men and women, National Bureau of Economic Research. NBER Working Paper Series. Working Paper #726. http://www.nber.org/).

Tay SK, Tay KJ (2004). Passive cigarette smoking is a risk factor in cervical neoplasia. *Gynecol Oncol*, 93(1):116-120.

Taylor AE, Johnson DC, Kazemi H (1992). Environmental tobacco smoke and cardiovascular disease. A position paper from the Council on Cardiopulmonary and Critical Care, *American Heart Association. Circulation*, 86(2):699-702.

Taylor R, Najafi F, Dobson A (2007). Metaanalysis of studies of passive smoking and lung cancer: effects of study type and continent. *Int J Epidemiol*, 36(5):1048-1059.

The Scotland Office (1999). Devolution. <u>(http://www.scotlandoffice.gov.uk/devolution/).</u>

Thompson B, Coronado G, Chen L, et al. (2006). Preferred smoking policies at 30 Pacific Northwest colleges. *Public Health Rep*, 121:586-593.

Thomson GW, Wilson N (2004). Public attitudes about tobacco smoke in workplaces: the importance of workers' rights in survey questions. *Tob Control*, 13(2):206-207.

Thomson G, Wilson N (2006). One year of smokefree bars and restaurants in New Zealand: Impacts and responses. *BMC Public Health*, 6(64):1-9.

Thomson G, Wilson N, Howden-Chapman P (2006). Population level policy options for increasing the prevalence of smokefree homes. *J Epidemiol Community Health*, 60(4):298-304.

Tobacco Control Research and Evaluation (TCRE) (2007). Community support for smokefree cars legislation, findings from new South Australian research. Adelaide, Tobacco Control Research and Evaluation Program (http://www.cancersa.org.au/aspx/home.aspx).

Tominz R, Murolo G, Montina G, et al. (2006). [Exposure to passive smoking in local health units of northern Italy before and after the enforcement of the smoking ban]. *Epidemiol Prev*, 30(6):338-342.

Tong EK, Nguyen TT, Vittinghoff E, et al. (2008). Smoking behaviors among immigrant Asian Americans: rules for smoke-free homes. *Am J Prev Med*, 35(1):64-67.

Travers M, Cummings K, Hyland A, et al. (2004). Indoor air quality in hospitality venues before and after the implementation of a Clean Indoor Air Law - Western New York, 2003. MMWR Morb Mortal Wkly Rep, 53(44):1038-1041.

Travers MJ, Hyland A, Higbee C, et al. (2007). Tobacco smoke pollution exposure in hospitality venues around the U.S. and the effect of smokefree air policies. In: Proceedings of 13th annual meeting 21-24 February 2007, Austin TX. Madison, WI, Society for research on nicotine and tobacco. POS2-156: 114.

Trichopoulos D, Kalandidi A, Sparros L, et al. (1981). Lung cancer and passive smoking. Int J Cancer, 27(1):1-4.

Trichopoulos D, Kalandidi A, Sparros L (1983). Lung cancer and passive smoking: conclusion of Greek study. *Lancet*, 2(8351):677-678.

Trimble CL, Genkinger JM, Burke AE, et al. (2005). Active and passive cigarette smoking and the risk of cervical neoplasia. *Obstet Gynecol*, 105(1):174-181.

Trinidad DR, Gilpin EA, Pierce JP (2005). Compliance and support for smoke-free school policies. *Health Educ Res*, 20(4):466-475.

Trotter L, Wakefield M, Borland R (2002). Socially cued smoking in bars, nightclubs, and gaming venues: a case for introducing smokefree policies. *Tob Control*, 11(4):300-304.

Tsai FC, Smith KR, Vichit-Vadakan N, et al. (2000). Indoor/outdoor PM10 and PM2.5 in Bangkok, Thailand. J Expo Anal Environ Epidemiol, 10(1):15-26.

Tsai HT, Tsai Y, Yang SF, et al. (2007). Lifetime cigarette smoke and second-hand smoke and cervical intraepithelial neoplasm--a community-based case-control study. *Gynecol Oncol*, 105(1):181-188.

Tyrrell I (1999). Deadly enemies: tobacco and its opponents in Australia. Sydney, Australia, UNSW Press [http://books.google.fr/books?id=np8GnBlHkcAC&pg=PR5&lpg=PR3&dq=%22deadly+enemies+tobacco+and+its+opponents+in+australia%22&psp=1&output=html&sig=SfCuBlmkQwtDC5c_ME6uYS45QXE).

U.S. Department of Health Education and Welfare (USDHEW) (1964). Smoking and health: report of the Advisory Committee to the Surgeon General of the Public Health Service. Washington, DC, U.S. Public Health Service. Publication No. 1103.

U.S. Department of Health Education and Welfare (USDHEW) (1972). The health consequences of smoking. A report of the Surgeon General: 1972. Washington, DC, U.S. Public Health Service. Publication No. (HSM) 72-7516 (http://profiles.nlm.nih.gov/NN/B/B/P/M/_/nnbbpm.pdf).

U.S. Department of Health and Human Services (USDHHS) (1982). The health consequences of smoking: cancer. A report of the Surgeon General. Rockville, MD, U.S. Department of Health and Human Services, Center for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health (http://profiles.nlm.nih.gov/NN/B/C/D/W//nnbcdw.pdf).

U.S. Department of Health and Human Services (USDHHS) (1986). The health consequences of involuntary smoking: a report of the Surgeon General. Rockville, MD, U.S. Department of Health and Human Services, Center for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health (http://www.surgeongeneral.gov/library/secondhandsmoke/report/).

U.S. Department of Health and Human Services (USDHHS) (1988). The health consequences of nicotine addiction. A report of the Surgeon General. Rockville, MD, U.S. Department of Health and Human Services, Center for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. CDC - 88-8406 (http://www.surgeongeneral.gov/library/secondhandsmoke/report/).

U.S. Department of Health and Human Services (USDHHS) (1989). Reducing the health consequences of smoking, 25 years of progress. Rockville, MD, U.S. Department of Health and Human Services, Center for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.

- U.S. Department of Health and Human Services (USDHHS) (1990). The health benefits of smoking cessation: a report of the Surgeon General. Rockville, MD, U.S. Department of Health and Human Services, Center for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health (http://profiles.nlm.nih.gov/NN/B/B/C/T/_/nnbbct.pdf).
- U.S. Department of Health and Human Services (USDHHS) (2000). Healthy people 2010. Volume 1, 2nd ed. Rockville, MD, U.S. Department of Health and Human Services, Center for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. (http://www.healthypeople.gov/Document/tableofcontents.htm#volume1).
- U.S. Department of Health and Human Services (USDHHS) (2001). Women and smoking: a report of the Surgeon General. Rockville, MD, U.S. Department of Health and Human Services, Center for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health (http://www.cdc.gov/tobacco/data_statistics/sgr/sgr_2001/#full).
- U.S. Department of Health and Human Services (USDHHS) (2004). The health consequences of smoking: a report of the Surgeon General. Atlanta, GA, U.S. Department of Health and Human Services, Center for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.
- U.S. Department of Health and Human Services (USDHHS) (2006). Control of secondhand smoke exposure. In: The health consequences of involuntary exposure to tobacco smoke: a report of the surgeon general. Atlanta, GA, U.S. Department of Health and Human Services, Center for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.
- U.S. Environmental Protection Agency (EPA) (1992). Respiratory health effects of passive smoking: lung cancer and other disorders. Washington, DC, Office of Health and Environmental Assessment, Office of Research and Development, U.S Environmental Protection Agency. Publication No. EPA/600/6-90/006F.

Valente P, Forastiere F, Bacosi A, et al. (2007). Exposure to fine and ultrafine particles from secondhand smoke in public places before and after the smoking ban, Italy 2005. *Tob Control*, 16(5):312-317.

van Walbeek C, Blecher H, van Graan M (2007). Effects of the tobacco products control amendment Act of 1999 on restaurant revenues in South Africa--a survey approach. *S Afr Med J*, 97(3):208-211.

Viehbeck SM, McDonald PW (2004). An examination of the relationship between municipal smoke-free bylaw strength and the odds of being a former smoker. *Can J Public Health*, 96(1):42-44.

Villeneuve PJ, Johnson KC, Mao Y, et al. (2004). Environmental tobacco smoke and the risk of pancreatic cancer: findings from a Canadian population-based case-control study. Can J Public Health, 95(1):32-37.

Vineis P (2008). Personal communication to E. Khaykin and J. Samet.

Vineis P, Hoek G, Krzyzanowski M, et al. (2007). Lung cancers attributable to environmental tobacco smoke and air pollution in non-smokers in different European countries: a prospective study. Environ Health, 6:1-7.

Wakefield MA, Chaloupka F (2000). Effectiveness of comprehensive to bacco control programmes in reducing teenage smoking in the USA. *Tob Control*, 9(2):177-186.

Wakefield MA, Wilson D, Owen N, et al. (1992). Workplace smoking restrictions, occupational status, and reduced cigarette consumption. *J Occup Med*, 34(7):693-697.

Wakefield MA, Roberts L, Owen N (1996). Trends in prevalence and acceptance of workplace smoking bans among indoor workers in South Australia. *Tob Control*, 5(3):205-208.

Wakefield MA, Chaloupka F, Kaufman N, et al. (2000a). Effect of restrictions on smoking at home, at school, and in public places on teenage smoking: cross sectional study. Br Med J, 321(7257):333-337.

Wakefield MA, Banham D, Martin J, et al. (2000b). Restrictions on smoking at home and urinary cotinine levels among children with asthma. Am J Prev Med, 19(3):188-192.

Wakefield MA, Siahpush M, Scollo M, et al. (2002). The effect of a smoke-free law on monthly restaurant retail turnover in South Australia. Aust N Z J Public Health, 26(4):375-382.

Wald NJ, Nanchahal K, Thompson SG, *et al.* (1986). Does breathing other people's tobacco smoke cause lung cancer? *Br Med J (Clin Res Ed)*, 293(6556):1217-1222.

Walsh RA, Tzelepis F (2003). Support for smoking restrictions in bars and gaming areas: review of Australian studies. *Aust N Z J Public Health*, 27(3):310-322.

Walsh RA, Paul CL, Tzelepis F (2000). Overwhelming support for smoking bans. *Aust N Z J Public Health*, 24(6):640-641.

Wamboldt FS, Balkissoon RC, Rankin AE, et al. (2008). Correlates of household smoking bans in low-income families of children with and without asthma. Fam Process, 47(1):81-94.

Wang Z, Tang L, Sun G, et al. (2006). Etiological study of esophageal squamous cell carcinoma in an endemic region: a population-based case control study in Huaian, China. *BMC Cancer*, 6:287

Waring MS, Siegel JA (2007). An evaluation of the indoor air quality in bars before and after a smoking ban in Austin, Texas. *J Expo Sci Environ Epidemiol*. 17(3):260-268.

Warren CW, Riley L, Asma S, *et al.* (2000). Tobacco use by youth: a surveillance report from the Global Youth Tobacco Survey project. *Bull World Health Organ*, 78(7):868-876.

Wartenberg D, Calle EE, Thun MJ, et al. (2000). Passive smoking exposure and female breast cancer mortality. *J Natl Cancer Inst*, 92(20):1666-1673.

Wasserman J, Manning W, Newhouse J, et al. (1991). The effects of excise taxes and regulations on cigarette smoking. J Health Econ, 10(1):43-64.

Weber MD, Bagwell DA, Fielding JE, et al. (2003). Long term compliance with California's Smoke-Free Workplace Law among bars and restaurants in Los Angeles County. *Tob Control*, 12(3):269-273.

Weiss B (1989). Behavior as an endpoint for inhaled toxicants. In: McClellan RO, Henderson RF, eds. Concepts in inhalation toxicology. New York, Hemisphere Publishing: 475-493.

Weiss ST (1986). Passive smoking and lung cancer. What is the risk? *Am Rev Respir Dis*, 133(1):1-3.

Wewers ME, Uno M (2002). Clinical interventions and smoking ban methods to reduce infants' and children's exposure to environmental tobacco smoke. *J Obstet Gynecol Neonatal Nurs*, 31(5):592-598.

Whincup PH, Gilg JA, Emberson JR, et al. (2004). Passive smoking and risk of coronary heart disease and stroke: prospective study with cotinine measurement. *Br Med J*, 329(74 59):200-205.

White JR, Froeb HF (1980). Small-airways dysfunction in nonsmokers chronically exposed to tobacco smoke. *N Engl J Med*, 302(13):720-723

WHO (1970). WHA23.32 Health consequences of smoking. Smoking and Health, WHO Chronicle 24,365. Geneva, Switzerland, World Health Organization.

WHO (1975). Smoking and its effects on health. Report of a WHO expert committee. Technical report series 568. Geneva, Switzerland, World Health Organization (http://whqlibdoc.who.int/trs/WHO_TRS_568.pdf).

WHO (1986). WHA39.14 Tobacco or health. Resolution WHA39.14. Geneva, Switzerland, World Health Organization.

WHO (1990). WHA 43.16 Tobacco or health. Resolution WHA43.16. Geneva, Switzerland, World Health Organization.

WHO (1999). International consultation on environmental tobacco smoke (ETS) and child health. Geneva, Switzerland, World Health Organization (http://www.who.int/tobacco/research/en/ets_report.pdf).

WHO (2005). WHO Framework Convention on Tobacco Control. Geneva, Switzerland, World Health Organization_(http://www.who.int/tobacco/framework/WHO_FCTC_english.pdf).

WHO (2007a). Guidelines for Article 8 of the Framework Convention on Tobacco Control. Geneva, Switzerland, World Health Organization.

WHO (2007b). Protection from exposure to secondhand tobacco smoke: policy recommendations. Geneva, Switzerland, World Health Organization_(http://www.who.int/tobacco/resources/publications/wntd/2007/who_protection_exposure_final_25June2007.pdf).

WHO (2008). WHO report on the global tobacco epidemic: the MPOWER package. Geneva, Switzerland, World Health Organization (http://www.who.int/tobacco/mpower/mpower_report_full_2008.pdf).

Wilkinson JD, Arheart KL, Lee DJ (2006). Accuracy of parental reporting of secondhand smoke exposure: The National Health and Nutrition Examination Survey III. *Nicotine Tob Res.* 8(4):591-597.

Wilson N, Thomson G, Edwards R (2007). Lessons from Hong Kong and other countries for outdoor smokefree areas in New Zealand? *N Z Med J.* 120:U2624.

Winkelstein ML, Tarzian A, Wood RA (1997). Parental smoking behavior and passive smoke exposure in children with asthma. *Ann Allergy Asthma Immunol*, 78(4):419-423.

Wipfli H, Avila-Tang E, Navas-Acien A, et al. (2008). Secondhand smoke exposure among women and children: evidence from 31 countries. Am J Public Health, 98(4):672-679.

Wold B, Currie C, Roberts C, *et al.* (2004a). National legislation on school smoking restrictions in eight European countries. *Health Promot Int*, 19(4):482-488.

Wold B, Torsheim T, Currie C, et al. (2004b). National and school policies on restrictions of teacher smoking: a multilevel analysis of student exposure to teacher smoking in seven European countries. Health Educ Res, 19(3):217-226.

Wong S (2002). York region no-smoking by-law #A-0285-2000-105. Ontario, Canada, Ontario Campaign for Action on Smoking http://www.york.ca/services/public+health+and+safety/tobacco+free+living/york+region+no-smoking+by-law.htm).

Wong GC, Berman B, Hoang T, et al. (2002). Children's exposure to environmental tobacco smoke in the home: comparison of urine cotinine and parental reports. Arch Environ Health, 57(6):584-590.

Wong GC, Bernaards CA, Berman BA, et al. (2004). Do children with asthma and their parents agree on household ETS exposure? Implications for asthma management. Patient Educ Couns, 53(1):19-25.

Woodruff TJ, Rosbrook B, Pierce J, et al. (1993). Lower levels of cigarette consumption found in smoke-free workplaces in California. *Arch Intern Med*, 153(12):1485-1493.

Woodward A, Laugesen M (2001). How many deaths are caused by second hand cigarette smoke? *Tob Control*, 10(4):383-388.

Wu AH (1999). Exposure misclassification bias in studies of environmental tobacco smoke and lung cancer. *Environ Health Perspect*, 107(Suppl 6):873-877.

Wu MT, Lee L, Ho C, et al. (2003). Lifetime exposure to environmental tobacco smoke and cervical intraepithelial neoplasms among nonsmoking Taiwanese women. Arch Environ Health, 58(6):353-359.

Wynder EL, Graham EA (1950). Tobacco smoking as a possible etiologic factor in bronchiogenic carcinoma; a study of 684 proved cases. *J Am Med Assoc*, 143(4):329-336.

Wynder EL, Graham EA, Croninger AB (1953). Experimental production of carcinoma with cigarette tar. *Cancer Res*, 13(12):855-864.

Wynder WL (1961). An appraisal of the smoking-lung-cancer issue. *N Engl J Med*, 264:1235-1240.

Yang G, Fan L, Tan J, *et al.* (1999). Smoking in China: findings of the 1996 National Prevalence Survey. *JAMA*, 282(13):1247-1253.

Yang T, Wu Y, Abdullah A, et al. (2007). Attitudes and behavioral response toward key tobacco control measures from the FCTC among Chinese urban residents. BMC Public Health. 7(1):248.

Yañez E (2002). Clean indoor air and communities of color: challenges and opportunities. Washington, DC, The Praxis Project (http://www.thepraxisproject.org/tools/CIA and CoC.pdf).

You RX, Thrift AG, McNeil J, et al. (1999). Ischemic stroke risk and passive exposure to spouses' cigarette smoking. Melbourne Stroke Risk Factor Study (MERFS) Group. Am J Public Health, 89(4):572-575.

Young D, Borland R, Siahpush M, et al. (2007). Australian smokers support stronger regulatory controls on tobacco: findings from the ITC Four-Country Survey. Aust N Z J Public Health, 31(2):164-169.

Yousey YK (2006). Household characteristics, smoking bans, and passive smoke exposure in young children. *J Pediatr Health Care*, 20(2):98-105.

Yu MC, Garabrant DH, Huang TB, et al. (1990). Occupational and other non-dietary risk factors for nasopharyngeal carcinoma in Guangzhou, China. Int J Cancer, 45(6):1033-1039.

Yuan JM, Wang XL, Xiang YB, et al. (2000). Non-dietary risk factors for nasopharyngeal carcinoma in Shanghai, China. *Int J Cancer*, 85(3):364-369.

Yurekli AA, Zhang P (2000). The impact of clean indoor-air laws and cigarette smuggling on demand for cigarettes: an empirical model. *Health Econ*, 9(2):159-170.

Zeegers MP, Goldbohm RA, van den Brandt PA (2002). A prospective study on active and environmental tobacco smoking and bladder cancer risk (The Netherlands). *Cancer Causes Control*, 13(1):83-90.

Zhang X, Shu X, Yang G, *et al.* (2005). Association of passive smoking by husbands with prevalence of stroke among Chinese women nonsmokers. *Am J Epidemiol*, 161(3): 213-218

Zheng W, McLaughlin J, Chow W, et al. (1993). Risk factors for cancers of the nasal cavity and paranasal sinuses among white men in the United States. Am J Epidemiol, 138(11):965-972