



**RED MEAT AND  
PROCESSED MEAT**

VOLUME 114

This publication represents the views and expert opinions of an IARC Working Group on the Evaluation of Carcinogenic Risks to Humans, which met in Lyon, 6–13 October 2015

LYON, FRANCE - 2018

**IARC MONOGRAPHS  
ON THE EVALUATION  
OF CARCINOGENIC RISKS  
TO HUMANS**

# 1. EXPOSURE DATA

---

## 1.1 Identification of the agents

The focus of this *Monograph* is the consumption of red meat and processed meat. These terms are defined below.

### 1.1.1 Red meat

Red meat refers to fresh unprocessed mammalian muscle meat (e.g. beef, veal, pork, lamb, mutton, horse, or goat meat), which may be minced or frozen, and is usually consumed cooked.

### 1.1.2 Offal

Mammalian offal refers to the internal organs and entrails of a butchered animal (e.g. scrotum, small intestine, heart, brain, kidney, liver, thymus, pancreas, testicle, tongue, tripe, or stomach) consumed as such. Mammalian offal is considered to be a specific food category in food consumption surveys ([FAO, 2015](#)); however, mammalian offal is reported together with red meat in some epidemiological studies.

### 1.1.3 Processed meat

Processed meat refers to any meat that has been transformed through one or several of the following processes: salting, curing, fermentation, smoking, or other processes to enhance flavour or improve preservation. Most processed meats are made from pork or beef, but may also include other red meats, poultry, offal, or meat

by-products such as blood. It is also important to distinguish between industrial processing and household preparations. As there is a huge variety of processed meat products, it is difficult to sort them into categories ([Santarelli et al., 2008](#)). However, based on recommendations by the Food and Agriculture Organization of the United Nations (FAO) ([Heinz & Hautzinger, 2007](#)), different groups of industrial processed meats can be proposed.

#### (a) Cured meat pieces

Examples of cured meat include raw beef, raw ham, cooked beef, cooked ham, corned beef, and bacon.

Curing is a process by which the meat is treated with a small amount of salt (sodium chloride, NaCl, with or without potassium chloride, KCl), with or without nitrate or nitrite salts. Curing enhances shelf-life by preserving and preventing the spoilage of meat. Cured meat cuts are made of entire pieces of muscle meat and can be subdivided into two groups: cured raw meats and cured cooked meats ([Pearson & Gillett, 1996](#); [Heinz & Hautzinger, 2007](#); [Honikel, 2010](#)). Cured raw meats are consumed uncooked. They do not undergo any heat treatment during production, which involves curing, fermentation, and ripening in controlled conditions to make the meats palatable. For cured cooked meats, the raw muscle meat is always cured and then undergoes treatment to achieve the desired palatability ([Heinz & Hautzinger, 2007](#)).

(b) *Fresh industrial processed meat products*

Examples of fresh industrial processed meat products include sausage and kebab.

These products are mixtures composed of comminuted muscle meat and animal fat in varying proportions. Products are salted only, not cured. Non-meat ingredients are added in smaller quantities for improvement of flavour and binding, or in larger quantities for volume extension (reducing costs). All meat and non-meat ingredients are mixed when raw. If the fresh meat mixture is packed into casings, the product is defined as sausage. Heat treatment is applied immediately before consumption to make the products palatable ([Heinz & Hautzinger, 2007](#)). [The Working Group noted that a hamburger is considered as belonging to this category of processed meat when fat, salt, or other additives are added to the hamburger meat, but is considered as red meat when it contains minced beef only.]

(c) *Precooked ready-to-eat products*

Examples of precooked ready-to-eat products include frankfurter, mortadella, liver sausage, blood sausage, canned corned beef, and liver pâté.

These products are prepared from muscle meat, fat, and other edible meat by-products (blood and liver) or non-meat ingredients. These products are processed raw through comminuting and mixing. Sometimes the raw meat material can be precooked before it is ground or chopped, and other ingredients are added. The resulting mixture is portioned and then submitted to heat treatment to induce protein coagulation. This leads to the typical firm, elastic texture of precooked ready-to-eat products, as well as a desired palatability and a certain degree of bacterial stability ([Heinz & Hautzinger, 2007](#)).

(d) *Fermented sausages*

Examples of fermented sausages include salami, chorizo, pepperoni, and traditional Asian products such as nem.

Fermented sausages are uncooked meat products, and consist of coarse mixtures of lean meats and fatty tissues combined with salt, nitrite (curing agent), sugar, spices, and other non-meat ingredients packed into casings. Their characteristic properties (flavour, firm texture, and red curing colour) originate from the fermentation process. Short or long ripening phases, combined with moisture reduction, are necessary to develop the typical flavour and texture of the final product. The fermented sausages are not subjected to any heat treatment during processing and, in most cases, are distributed and consumed raw ([Heinz & Hautzinger, 2007](#)).

(e) *Dried meat*

Examples of dried meat include dried meat strips or flat pieces.

Drying, or drying in combination with smoking, is practised all over the world and is probably the most ancient method of meat preservation.

Dried meat products result from dehydration or drying of lean meat in natural or artificial conditions ([Zukál & Incze, 2010](#)). Pieces of lean meat without adherent fat are cut to a specific uniform shape that permits the gradual and equal drying of whole batches of meat. Salt, nitrite, and sugar may be added to the meat before the drying process. Many of the nutritional properties of meat, particularly the protein content, remain unchanged through drying. Common dried meat products are beef jerky from the USA, biltong from South Africa, and tasajo from South America ([Heinz & Hautzinger, 2007](#)).

Meat may be smoked raw or after salting, marinating, cooking, or other treatments. There are many types of smoking, leading to products with very different sensory properties and

shelf-lives. Warm or cold smoking can be used. Warm smoking is carried out at temperatures of 23–45 °C. Cold smoking is carried out at temperatures of 12–25 °C and is used in the manufacturing of raw fermented sausages made from cured meats.

Drying and smoking are used to improve the shelf-life and organoleptic properties of meat products. In developing countries such as Africa, where extending shelf-life is the priority, drying is the most used process. In parallel with simple drying, west Africa has refined the hot smoking process to further improve shelf-life through the preservative and antibacterial effects of smoke substances. To lower the cost of meat products, African countries have also developed traditional products consisting of mixtures of meat and vegetables. Central and southern American countries have adapted European meat processing techniques for local meat products, especially for barbecuing (e.g. chorizo criollo or morcilla) ([Heinz & Hautzinger, 2007](#)).

## 1.2 Meat composition

### 1.2.1 Red meat

#### (a) Main components

The animal carcass consists of muscle, connective tissue, fat and bone, and about 75% water, depending on the species, breed, size, and age. For a given species, the muscle is relatively constant in composition ([Table 1.1](#)). Red meat contains high biological value proteins and essential micronutrients, including vitamins and minerals ([Table 1.2](#); [Williams, 2007](#)). The composition of the meat varies based on the animal species, sex, age, and diet, as well as the climate and activity during its growth ([Lorenzo et al., 2010](#)). Total nitrogen, fat, and iron levels increase as the animal approaches maturity. In addition, the ratio of polyunsaturated fatty acids (PUFAs) to saturated fatty acids (SFAs) decreases with the maturity of the animal. The nutritional

value of meat is also significantly affected by the livestock production system ([Lorenzo et al., 2010, 2014](#)).

#### (i) Protein

Red meat contains 20–25 g of protein per 100 g. The proteins are highly digestible (94%) and provide all essential amino acids (lysine, threonine, methionine, phenylalanine, tryptophan, leucine, isoleucine, and valine) ([Williams, 2007](#)).

#### (ii) Fat

Red meat is also a source of fatty acids. Fat in red meat is subcutaneous, intramuscular, or intermuscular, and the composition will vary according to the animal's age, sex, breed, and diet, as well as the cut of meat ([Wood & Enser, 1997](#)). For example, the amount of fat in raw cattle longissimus muscle can range from 0.59% to 16%, depending on the breed ([Barnes et al., 2012](#)). Fat in meat includes SFAs, mono-unsaturated fatty acids (MUFAs), and PUFAs. The typical fatty acid composition of fat in beef is reported to be 46.5, 48.9, and 4.59 g per 100 g of total fatty acids for SFAs, MUFAs, and PUFAs, respectively. While these proportions are similar in all red meats, exact amounts depend on the type of meat ([Givens, 2005](#)). The main SFAs present in red meat are palmitic acid and stearic acid, and the main MUFA is oleic acid. Red meat also contains n-3 PUFAs, such as  $\alpha$ -linolenic acid, and n-6 PUFAs, such as linoleic acid. The animal's diet strongly influences PUFA levels in meat. For example, meat from foals raised by extensive production systems on wood pastures has higher levels of n-3 PUFAs than meat from foals fed concentrate ([Lorenzo et al., 2010, 2014](#)). The last category of fat found in the red meat of ruminants is conjugated linoleic acids, the levels of which also depend on feeding practices ([Wood et al., 1999](#); [Givens, 2005](#)).

**Table 1.1 Chemical composition of typical mammalian muscle (red meat) for consumption**

Main component	Constituents	Wet weight (%)
<i>Water</i>		75.00
<i>Protein</i>		19.00
Myofibrillar:		11.50
	Myosin	5.50
	Actin	2.50
	Connectin	0.90
	Nebulin (N2 line protein)	0.30
	Tropomyosins	0.60
	Troponins, C, I and T	0.60
	$\alpha, \beta, \gamma$ Actinins	0.50
	Myomesin (M-line protein) and C proteins	0.20
	Desmin, filamin, F- and I-proteins, etc.	0.40
Sarcoplasmic:		5.50
	Glyceraldehyde phosphate dehydrogenase	1.20
	Aldolase	0.60
	Creatine kinase	0.50
	Other glycolytic enzymes	2.20
	Myoglobin	0.20
	Haemoglobin and other unspecified extracellular proteins	0.60
Connective tissue and organelles:		2.00
	Collagen	1.00
	Elastin	0.05
	Mitochondrial etc. (including cytochrome c and insoluble enzymes)	0.95
<i>Lipid</i>		2.50
Neutral lipid; phospholipids; fatty acids; fat-soluble substances		2.50
<i>Carbohydrate</i>		1.20
	Lactic acid	0.90
	Glucose-6-phosphate	0.15
	Glycogen	0.10
	Glucose, traces of other glycolytic intermediates	0.05
<i>Miscellaneous, soluble non-protein substances</i>		2.30
Nitrogenous:		1.65
	Creatinine	0.55
	Inosine monophosphate	0.30
	Di- and tri-phosphopyridine nucleotides	0.10
	Amino acids	0.35
	Carnosine, anserine	0.35
Inorganic:		0.65
	Total soluble phosphorus	0.20
	Potassium	0.35
	Sodium	0.05
	Magnesium	0.02
	Calcium, zinc, trace metals	0.03
<i>Vitamins</i>		
Various fat- and water soluble vitamins		Minute

This table was published in Lawrie's Meat Science 6th edition, [Lawrie \(1998\)](#), Page No. 59, Copyright Elsevier (1998)

**Table 1.2 Average nutrient composition (per 100 g) of the lean component of red meat**

Nutrient	Beef	Veal	Lamb	Mutton
Moisture (g)	73.1	74.8	72.9	73.2
Protein (g)	23.2	24.8	21.9	21.5
Fat (g)	2.8	1.5	4.7	4.0
Energy (kj)	498	477	546	514
Cholesterol (mg)	50	51	66	66
Thiamin (mg)	0.04	0.06	0.12	0.16
Riboflavin (mg)	0.18	0.20	0.23	0.25
Niacin (mg)	5.0	16.0	5.2	8.0
Vitamin B6 (mg)	0.52	0.8	0.10	0.8
Vitamin B12 (µg)	2.5	1.6	0.96	2.8
Pantothenic acid (mg)	0.35	1.50	0.74	1.33
Vitamin A (µg)	< 5	< 5	8.6	7.8
Beta-carotene (µg)	10	< 5	< 5	< 5
Alpha-tocopherol (mg)	0.63	0.50	0.44	0.20
Sodium (mg)	51	51	69	71
Potassium (mg)	363	362	344	365
Calcium (mg)	4.5	6.5	7.2	6.6
Iron (mg)	1.8	1.1	2.0	3.3
Zinc (mg)	4.6	4.2	4.5	3.9
Magnesium (mg)	25	26	28	28
Phosphorus (mg)	215	260	194	290
Copper (mg)	0.12	0.08	0.12	0.22
Selenium (µg)	17	< 10	14	< 10

Adapted from [Williams \(2007\)](#). *Nutrition & Dietetics*, John Wiley & Sons

### (iii) Vitamins

The only natural source of vitamin B12 is in food derived from animal products. Red meat is a rich source of B vitamins such as B6, B12, niacin, and thiamine ([Gille & Schmid, 2015](#)). For example, 100 g of lean beef meat will provide 2.5 µg of vitamin B12, corresponding to 79% of the recommended dietary intake for this nutrient. The older the animal, the richer its meat will be in B vitamins ([Williams, 2007](#)). Pork contains a high level of thiamine compared with other meats ([Bender, 1992](#)). While the concentration of vitamin E in red meat is low, it is higher in fattier cuts of meat. Vitamin A and folate are found at higher levels in liver than in lean muscle meat ([Bender, 1992](#)).

### (iv) Minerals

Red meat is one of the richest sources of minerals such as iron or zinc, and has a higher mineral bioavailability than plant products ([Williams, 2007](#)). For example, 100 g of lean beef meat will provide approximately 1.8 mg of iron and 4.6 mg of zinc, corresponding to approximately 14% and 42%, respectively, of the recommended dietary intake for these nutrients ([Williams, 2007](#)). Red meat is also a good source of selenium. For example, 100 g of lean beef meat will provide about 17 µg of selenium, corresponding to approximately 26% of the recommended dietary intake for this nutrient ([Williams, 2007](#)).

*(v) Creatine*

Creatine levels in skeletal muscle average 350 mg per 100 g of red meat ([Purchas & Busboom, 2005](#); [Williams, 2007](#)). Cooking of the muscle meat transforms creatine into creatinine through non-enzymatic conversion. Creatine and creatinine in meat are critical precursors in the formation of heterocyclic aromatic amines (HAAs) ([Skog et al., 1998, 2000](#)).

*(b) Effect of slaughtering and storage post mortem*

Preslaughter handling of the animal can have an impact on the composition and quality of the meat. For example, stressed or fatigued animals have depleted glycogen. Before slaughtering, animals are usually stunned and then exsanguinated. Blood is drained from the carcass, leading to a loss of oxygen and a depletion of adenosine triphosphate, as well as the combination of the proteins actin and myosin to form actinomyosin to cause muscle contraction. After slaughtering, glycogen is converted to lactic acid, and pH levels fall to approximately 5.5 over a period of 24–36 hours and can have an impact on microbial content ([Lawrie, 1998](#); [Lawrie & Ledward, 2006](#)).

A major safety concern in meat production and storage is bacterial contamination. Although muscle is usually sterile, bacterial contamination from gastrointestinal contents and butchering instruments is common. Bacterial contamination is minimized by low temperatures, low temperatures and packaging in a controlled atmosphere. Minced and comminuted meats with larger surface areas may be more likely to become contaminated than large cuts of meat ([Lawrie, 1998](#); [Lawrie & Ledward, 2006](#)). Physical and chemical methods, such as spray washing with hot water, can be used for microbial decontamination. Water may be chlorinated and combined with weak organic acids, phosphates, hydrogen peroxide, or ozone to improve antimicrobial activities. Improving hygiene levels by means of

antibiotics (chlorotetracycline and oxytetracycline) is increasingly discouraged and regulated. Irradiation is permitted in the USA, but it can induce free radical formation ([Isam et al., 2007](#)). Freezing of red meat results in little to no loss of nutrients, apart from vitamin E, which is oxidized. Proteins remain unchanged during frozen storage, but fats are susceptible to changes from oxidation. Pork meat, which is richer in unsaturated fatty acids than other red meats, is the most susceptible to these changes. Once meat is defrosted, the juices containing soluble proteins, vitamins, and minerals are lost ([Rahman, 2007](#)).

*(c) Chemical contaminants and residues*

Residues of drugs (e.g. antibiotics and hormones), pesticides, and agricultural chemicals can be found in meat and meat products (e.g. as a result of exposure of the animals to chemicals used on buildings, grazing areas, and crops) ([Fig. 1.1](#); [Engel et al., 2015](#)). Additionally, several hundred substances may be used to treat animals, to preserve animal health, and to improve animal production, including antimicrobial agents, anticoccidial agents, anthelmintics, steroids, anti-inflammatory agents, tranquillizers, vasodilators, analgesics, and anaesthetics ([FDA, 2005](#)).

*(i) Veterinary drugs*

Veterinary drugs given to animals are strictly regulated in most developed countries ([FDA, 2005](#)), and maximum residue limits (MLRs) have been established for some of these drugs by the Codex Alimentarius.

*Antibiotics:* In the European Union (EU), the only antibiotics allowed as feed additives are coccidiostats and histomonostats ([European Commission, 2003](#)), as other antibiotics, especially if they are also used for humans, could induce antimicrobial resistance in consumers ([Chattopadhyay, 2014](#)). Cooking procedures degrade the residues of several antibacterial drugs, depending on the amount of heat

treatment involved, principally cooking time and temperature ([Heshmati, 2015](#)).

*Hormones:* In several countries, such as the USA, the use of hormones, including testosterone propionate, estradiol, estradiol benzoate, and progesterone, and compounds that display a high affinity for human hormone receptors are approved for food animal production. This raises concerns because these hormones, or their biologically active metabolites, may accumulate in edible tissues, potentially exposing consumers ([Nachman & Smith, 2015](#)). Cooking reduces, but does not eliminate, the potential for dietary exposure to hormones, such as estradiol, in ground beef ([Braekevelt et al., 2011](#)). [Table 1.3](#) lists the amounts of steroid hormones ingested via the diet from hormone-treated or non-hormone-treated animals, and the amounts of these hormones produced daily in the human body. [The Working Group noted that the ingestion of estradiol, progesterone, and testosterone from meat appears to be minor relative to what is biosynthesized in humans ([Table 1.3](#)).]

Environmental and phytosanitary contaminants can also occur in meat products.

#### (ii) *Pesticide residues*

Pesticide residues used for phytosanitary treatments may be present in meat products. Animals consume plants treated with pesticides or contaminated by persistent pesticides in the environment. However, vegetable consumption remains by far the main dietary source of human exposure to pesticides ([Kan & Meijer, 2007](#)).

#### (iii) *Dioxins and dioxin-like products*

Dioxins and dioxin-like products are divided into three groups: polychlorinated dibenzodioxins (PCDDs), polychlorinated dibenzofurans (PCDFs), and polychlorinated biphenyls (PCBs) (see [Fig. 1.1](#)).

These contaminants, which are mainly produced by industrial processes, are ubiquitous in foods of animal origin, and accumulate in the fatty tissues of animals and humans ([Larsen, 2006](#); [IARC, 2012a, 2016](#)). Food, including meat, remains the primary source of human exposure to these contaminants in the general population ([IARC, 2012a, 2016](#)).

#### (iv) *Brominated flame retardants*

Brominated flame retardants (BFRs) are widely used in plastic materials, textiles, electric and electronic equipment, and of construction materials for livestock buildings. There are five classes of BFRs: polybrominated diphenyl ethers (PBDEs), hexabromocyclododecanes (HBCDs), tetrabromobisphenol A, and other phenols, polybrominated biphenyls (PBBs) ([IARC, 2016](#)), and unclassified BFRs ([ANSES, 2011, 2012](#); [IARC, 2016](#)). The persistence of BFRs in the environment is a public health concern ([AFSSA, 2005](#); [ANSES, 2011](#); [EPA, 2010](#)). The main source of human exposure to BFRs is the consumption of fish and meat products ([Lyche et al., 2015](#)). Studies have shown that the cooking process and, to a greater extent, the type of meat item influence levels of PBDEs.

#### (v) *Polycyclic aromatic hydrocarbons*

Polycyclic aromatic hydrocarbons (PAHs) can be generated in the environment or during the processing of foods; this is discussed further in Section 1.2.3(a)(iii). PAHs are closely monitored by health agencies in developed countries ([IARC 2010a](#); [Schroeder, 2010](#)). Furthermore, PAH levels can increase depending on cooking conditions.

#### (vi) *Heavy metals*

Contamination by heavy metals such as cadmium, lead, arsenic, or mercury largely occurs from industrial wastes ([IARC, 2012b](#)). Meat consumption is a significant source of human exposure to lead and cadmium ([Kan & Meijer, 2007](#)).



**Table 1.3 Comparison of the amounts of steroid hormones produced daily in the human body and ingested via the diet from hormone-treated animals**

Hormone	Total daily production (µg/day) ( <a href="#">JECFA, 2000</a> ; <a href="#">EFSA, 2007</a> )	Residue in muscle (µg/kg) ( <a href="#">Paris et al., 2006</a> )		Ingested amount via intake of muscle from treated animals <sup>a</sup> (µg/day)
		Non-treated animals	Treated animals	
Estradiol	< 14 (prepubertal boys) 10–24 (prepubertal girls) 27–68 (adult men) 30–470 (adult women)	0.003–0.035	0.011–0.28	0.0033–0.084
Progesterone	150–250 (prepubertal children) 416–750 (adult men, premenopausal women)	0.0–0.9	0.23–0.77	0.069–0.231
Testosterone	30–100 (prepubertal children) 210–480 (adult female) 2100–6900 (adult male)	0.006–0.029	0.031–0.36	0.0093–0.108

<sup>a</sup> Calculated according to an intake of 300 g/day of muscle

Reprinted from [Jeong et al. \(2010\)](#) © 2010 The Korean Society of Toxicology. License: CC BY-NC 3.0

### (vii) *Mycotoxins*

Mycotoxins, metabolites produced by mould, are toxic and may be carcinogenic to animals and humans. Livestock contamination by mycotoxins occurs via their diet, and human exposure results from consumption of contaminated livestock. However, cereals and oil seeds are the main sources of human exposure to mycotoxins ([FDA, 2008](#); [Marroquín-Cardona et al., 2014](#)). Mycotoxin residues accumulate in the blood, liver, and kidney, and, to a lesser degree, in muscle-derived meat products ([Kan & Meijer, 2007](#)). Mycotoxins, such as aflatoxin, are not destroyed by normal industrial processing or cooking since they are heat-stable ([Awadt et al., 2012](#)).

## 1.2.2 Processed meat

### (a) *Ingredients*

There are three major reasons for processing meat: reduction in microbial contamination, production of attractive products, and reduction of waste by reconstitution of muscle meat scraps or offal. Therefore, along with the main components, which are meat and animal



fat, a wide range of non-meat substances are used in processed meat products ([Bender, 1992](#); [Heinz & Hautzinger, 2007](#); [Toldrá, 2010](#); [Weiss et al., 2010](#)).

#### (i) *Non-meat ingredients of animal origin*


Although not commonly applied, non-meat ingredients of animal origin may be used to improve water binding and prevent fat separation during heat treatment. Some of these ingredients can also be considered as meat extenders. The most commonly used non-meat ingredients of animal origin are milk caseinate; whole milk or non-fat, dried milk; gelatine; blood plasma; eggs; and transglutaminase ([Sun & Holley, 2011](#)).

#### (ii) *Ingredients of plant origin*


Spices are predominantly functional ingredients, and are used in small quantities to provide or add flavour and taste to meat products. The most commonly used ingredients of plant origin are isolated soy protein ( $\leq 90\%$  protein) and wheat gluten ( $\leq 80\%$  protein). The most common ingredients used as fillers (if rich in carbohydrate) or meat extenders (if rich in protein) are soy flour or concentrate, cereal flour or cereals, starches, breadcrumbs, vegetables, and fruits ([Asgar et al.,](#)

**Fig. 1.1 Examples of environmental micropollutants potentially found in red and processed meats**2,3,7,8-Tetrachlorodibenzo-*p*-dioxin (TCDD)



Polychlorinated dibenzofuran (PCDFs)



Polychlorinated biphenyls (PCBs)



Polybrominated diphenyl ethers (PBDEs)

Benzo[*a*]pyrene (BaP)

Lindane (pesticide)

Adapted from *Meat Science*, Volume 109, E. Engel, J. Ratel, J. Bouhlef, C. Planche, M. Meurillon, Novel approaches to improving the chemical safety of the meat chain towards toxicants, Pages No. 75–85, Copyright (2015), with permission from Elsevier ([Engel et al., 2015](#))

[2010](#)). Several plant derivatives can also be used as fat replacers, antioxidants, or antimicrobials ([Hygreeva et al., 2014](#)).

### (iii) Chemical substances used as additives

There are a limited number of chemical substances allowed for meat processing, as the substances need to be safe for consumers and improve the quality of the final product. The most commonly used substances are salt (NaCl or NaCl plus KCl) for taste, impact on meat proteins, and shelf-life; nitrate and nitrite for curing, colour, flavour, and shelf-life; ascorbic acid for accelerated curing; phosphates for protein structuring and water binding; chemical preservatives for shelf-life; antioxidants for flavour and shelf-life; monosodium glutamate for enhancement of

flavour; and food colourings. Chemical additives have exclusively functional properties. They are used in small amounts, usually below 1%, with nitrate as low as 0.05% and with only salt in the range of 2% ( $\leq 4\%$  in some fermented dried products) ([Heinz & Hautzinger, 2007](#)).

### (b) Processing methods

Standard technical processing methods for meat products, such as cutting, comminuting, mixing, tumbling, or stuffing, are an important part of the manufacturing process ([Heinz & Hautzinger, 2007](#)). However, as these processes do not influence the formation of potentially carcinogenic process-induced toxicants, they will not be further detailed in this section.

Microbial inactivation can be achieved by “sous vide”, a method whereby foods are vacuum-packaged and then slow-cooked (temperature, 55–60 °C), as well as by cooking, canning (temperature, up to 121 °C), irradiation (chilled temperature, 0–4 °C), and high-pressure processing (300–600 MPa). Microbial inactivation can also be achieved by the addition of artificial preservatives such as nitrate or nitrite, weak organic acids, and/or salt or sugar. Canning is probably the most efficient meat preservation method, as it ensures the destruction of pathogens and food spoilage microorganisms, and allows foods to be easily handled and transported ([Guerrero Legarreta, 2010](#)).

The most common approaches to retard lipid oxidation, a major limiting factor in the shelf-life of dehydrated muscle tissue, is the addition of antioxidants and the use of appropriate packaging techniques ([Rahman, 2007](#)).

Chemical processing methods essentially include curing, smoking, and fermentation.

#### (i) Curing

Meat curing, in the narrow sense, is the addition of salt (NaCl or NaCl plus KCl), with or without nitrate or nitrite, during the manufacturing of meat products. Nitrate and nitrite are not used as sole curing agents. Each is always applied with salt. In meat products, the concentrations of nitrate and nitrite are usually in the range of 100–200 mg/kg, while salt is 2000 mg/kg or more. Salt lowers the water activity and enhances food safety. Salt also changes the protein structures in meat. Nitrate and nitrite support the safety action of salt, and improve the appearance and flavour. Nitrate must undergo reduction to nitrite to be effective. During the curing process, myoglobin is converted to nitrosomyoglobin, resulting in the characteristic cured meat colour ([EFSA, 2003](#); [Honikel, 2008, 2010](#)). Over the past few decades, ascorbic acid or its salt, ascorbate (e.g. isoascorbate or erythorbate), has been used in cured meat batters. Ascorbate reacts with oxygen

to form dehydroascorbate, and thus prevents the oxidation of nitrite to nitrate. Ascorbate is also added to reduce the formation of nitrosamines. Ascorbate, together with nitrite and salt, has an effective antimicrobial effect, particularly against *Clostridium botulinum* ([Honikel, 2010](#); [Sindelar & Milkowski, 2012](#)). Citric acid or sodium citrate may replace up to half of either form of the ascorbate/erythorbate reductants, but may not be used without the reductants ([Sindelar et al., 2010](#)). Nitrite addition is strictly regulated by international standards, and the amount allowed in cured meat is decreasing (see Section 1.5).

#### (ii) Smoking

Smoking refers to the exposure of meat to the smoke of burning wood ([Sikorski & Kalakowski, 2010](#)). Many cured products are also smoked, or contain soluble components of wood smoke, mainly to add flavour and increase shelf-life. Smoking gives meat a brown colour. It changes its flavour and improves its preservation, as smoke contains a wide variety of polyphenolic compounds as well as aldehydes and carboxylic acids, which have antimicrobial properties. Smoking can be done at different temperatures, depending on the end product ([Sikorski & Kalakowski, 2010](#)). However, wood pyrolysis may be hazardous, as the process is difficult to control and can lead to the generation of PAHs. Modifications to traditional wood pyrolysis processes are being studied to reduce the production and deposition of PAHs in processed meat ([Roseiro et al., 2011](#); [Ledesma et al., 2014](#)). An alternative is to use liquid smoke flavouring solutions produced from different wood products, under specific pyrolysis conditions and as per extraction protocols aimed at strongly reducing the concentration of PAHs ([Sikorski & Kalakowski, 2010](#)).

### (iii) Fermentation

Fermentation refers to a low-energy, biological acidulation (by cultured or wild microorganisms) and preservation method that results in a distinctive flavour and palatability, colour, and tenderness, as well as in enhanced microbiological safety. Since this process involves microorganisms, it is influenced by many environmental factors, including raw meat quality, sanitation, time, temperature, and humidity, all of which need to be strictly controlled during production. The reduction of pH and the lowering of water activity are microbial hurdles that aid in producing a safe product. Both natural and controlled fermentation processes involve lactic acid bacteria. Fermented sausages often have a long storage life, due to added salt, nitrate, and/or nitrite, and a low pH, due to lactic acid production by bacteria in the early stages of storage and later stages of drying ([Ockermann & Basu, 2010](#)).

### 1.2.3 Changes in meat composition due to cooking methods

Cooking can have a positive and negative impact on food quality. Cooking is important to inactivate pathogenic microorganisms, and improve palatability and digestibility ([Santé-Lhoutellier et al., 2008](#); [Bax et al., 2012, 2013](#)).

Generally, cooking reduces, but does not eliminate, meat contaminants such as hormones, antibiotics, chemicals (e.g. PCBs, PCDFs, and PCDDs), or metals (e.g. arsenic, cadmium, mercury, and lead) ([Hori et al., 2005](#); [Perelló et al., 2008](#); [Perelló et al., 2010](#); [Braekevelt et al., 2011](#); [Zeitoun & Ahmed, 2011](#); [Heshmati, 2015](#)). Furthermore, cooking can lead to the production of potential carcinogens.

The different cooking methods used to prepare red and processed meat may have varying influences on the production of potential carcinogens ([Table 1.4](#)). Cooking methods differ based on cooking temperature, direct or indirect contact with the heating source (flame), and use

of fat. The method has an impact on the formation of carcinogenic compounds such as HAAs or PAHs ([Skog et al., 1998](#); [Giri et al., 2015](#)). At low temperatures (around 100 °C), steaming, boiling, or stewing generate much lower levels of these carcinogenic compounds. For baking and roasting, temperatures are higher (up to 200 °C), but as there is limited direct contact with a hot surface, the formation of these carcinogenic compounds is also low ([Rohrmann et al., 2002](#)). Barbecuing, grilling, and pan-frying expose meat products to high temperatures, and to a hot surface or to direct flame, and thus can produce an appreciable level of these carcinogenic compounds ([American Institute for Cancer Research/World Cancer Research Fund, 1997](#); [Sinha et al., 1998a, b](#)).

#### (a) Red meat

This part of the section focuses on the toxicants found in red meat that are mostly produced by certain heating and cooking conditions.

##### (i) N-Nitroso compounds

N-Nitroso compounds (NOCs) are mainly formed endogenously in human organisms. No data report their formation in red meat during heat treatment; they are mainly considered processed meat toxicants (see Section 1.2.3(b)(i)).

##### (ii) Heterocyclic aromatic amines

HAAs are a family of heat-induced food toxicants that were discovered about 30 years ago by Professor Sugimura. Currently, about 25 HAAs have been identified in cooked meat, fish, and poultry products ([Sugimura et al., 2004](#)), as well as in cigarette smoke and diesel exhaust ([Manabe et al., 1991](#)). HAAs can be divided into two distinct families: aminoimidazoazaarenes and carbolines or pyrolytic HAAs ([Fig. 1.2](#) and [Table 1.5](#)). Aminoimidazoazaarenes are formed by Maillard reaction (a chemical reaction between amino acids, creatine/creatinine, and sugars), whereas carbolines and pyrolytic HAAs are formed at

**Table 1.4 Definition of cooking methods<sup>a</sup>**

Cooking method <sup>b</sup>	Definition
Baked	Cooked by dry heat in an oven, covered or uncovered, no additional fat used for cooking
Barbecued	Cooked on grill bars over burning charcoal, wood or gas
Battered and baked	Covered by batter (flour, milk, and egg mixture) and baked
Battered and fried	Covered by batter (flour, milk, and egg mixture) and fried
Boiled	Cooked in boiling liquid
Breaded and baked	Covered by an outer layer of breadcrumbs and baked
Breaded and fried	Covered by an outer layer of breadcrumbs and fried
Breaded and griddled	Covered by an outer layer of breadcrumbs and griddled
Coated and fried	Covered by an outer layer and fried: includes battered and fried, breaded and fried, in flour and fried
Deep fried	Cooked in hot fat or oil by immersing the food entirely
Fried	Generic descriptor for cooked in heated fat, usually over a direct source of heat
Griddled	Cooked on a heated flat metal surface over a source of direct heat; a little fat or oil may be used to grease the metal surface
Grilled	Cooked rapidly without moisture, on grill bars under or over intense direct heat, no fat used
In flour and fried	Covered by an outer layer of flour and fried
Microwaved	Cooked or reheated in a microwave oven; no fat used
Poached	Cooked by dropping in boiling liquid
Reheated	Made hot; no liquid nor fat is added
Roasted	Cooked by dry heat in an oven or over a fire
Shallow fried	Cooked in a shallow layer of heated fat
Steamed	Cooked by steam, in pressure cooker or cooked suspended above boiling water
Stewed	Cooked by boiling or simmering in liquid contained in an enclosed vessel; the food is cooked over a low heat for a long period of time
Stir fried/sautéed	Cooked by frying food over high heat, by stirring constantly to avoid sticking
Toasted	Cooked with direct heat until the surface of the food is browned

<sup>a</sup> Definitions based on the EPIC Study, [Rohrmann et al. \(2002\)](#)

<sup>b</sup> Cooking method is defined as the preparation of meat items just before consumption

Adapted by permission from Macmillan Publishers Ltd: [Rohrmann et al. \(2002\)](#). Cooking of meat and fish in Europe--results from the European Prospective Investigation into Cancer and Nutrition (EPIC). *European Journal of Clinical Nutrition*, Volume 56, issue 12, pages 1216–1230, copyright (2002)


elevated temperatures ([Murkovic, 2004](#)). The main source of human exposure to HAAs is via cooked proteinaceous foods; however, the levels of HAAs are highly dependent on the type of meat, cooking time, and cooking temperature, and generally increase with the level of “done-ness” ([Skog et al., 2000](#)).

The cooking method also influences HAA formation; it has been shown that high-temperature methods (pan-frying, grilling, and barbecuing) cause the highest HAA concentrations, especially for 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) ([Alaejos & Afonso, 2011](#)). The concentrations of HAAs in different cooked

meats are given in [Table 1.6](#). The concentrations of HAAs are highly variable. For a comprehensive review, see [Alaejos & Afonso \(2011\)](#). A series of linear tricyclic ring HAAs containing the 2-amino-1-methylimidazo[4,5-*g*]quinoxaline (IgQx) skeleton are formed in cooked meats at concentrations that are relatively high compared with the concentrations of their angular tricyclic ring isomers or related HAAs ([Ni et al., 2008](#)), such as 2-amino-3,8-dimethylimidazo[4,5-*f*]quinoxaline (MeIQx) and PhIP, which are known experimental animal carcinogens and potential human carcinogens ([IARC, 1993](#)). The toxicological properties of these recently discovered IgQx



**Fig. 1.2 Structures of heterocyclic aromatic amines found in cooked red and/or processed meats**

Principal aminoimidazoazaarenes found in red and processed meats




R = H (IQ)

R = Me (MeIQ)


 $R_1, R_2 = H, R_3 = Me$  (8-MeIQx) $R_1, R_3 = Me, R_2 = H$  (4,8-DiMeIQx)

PhIP


Principal pyrolytic HAAs found in red and processed meats




Harman




Norharman




Trp-P-1



Trp-P-2



AαC



MeAαC

The full chemical names of these compounds are given in [Table 1.5](#)

HAA, heterocyclic aromatic amines

Reprinted from *Cancer Science*, Volume 95, Takashi Sugimura, Keiji Wakabayashi, Hitoshi Nakagama, Minako Nagao, Heterocyclic amines: Mutagens/carcinogens produced during cooking of meat and fish, Pages No. 290–299, Copyright (2004), with permission from John Wiley & Sons ([Sugimura et al., 2004](#))

derivatives warrant further investigation and assessment.

Some methods to decrease the levels of HAAs in cooked meats have been described. For example, microwave pretreatment followed by disposal of the resulting liquid before frying of hamburger patties reduces the formation of some aminoimidazoazaarenes ([Felton et al., 1992](#)). Various studies have also emphasized the role of added antioxidants with phenolic or polyphenolic

moiety in the limitation of HAA formation – via their scavenging capacity for reactive radicals involved in the HAA mechanism of formation ([Balogh et al., 2000](#); [Vitaglione & Fogliano, 2004](#); [Gibis & Weiss, 2010, 2012](#)). Other compounds, such as organosulfur compounds, contained in garlic or onion, have also been shown to have an inhibitory effect on HAA formation ([Shin et al., 2002](#)).

**Table 1.5 Chemical names of heterocyclic aromatic amines potentially found in cooked red and processed meats**

Common abbreviation	Full name
IQ	2-amino-3-methylimidazo[4,5- <i>f</i> ]quinoline
MeIQ	2-amino-3,4-dimethylimidazo[4,5- <i>f</i> ]quinoline
IQx	2-amino-3-methylimidazo[4,5- <i>f</i> ]quinoxaline
MeIQx	2-amino-3,8-dimethylimidazo[4,5- <i>f</i> ]quinoxaline
4,8-DiMeIQx	2-amino-3,4,8-trimethylimidazo[4,5- <i>f</i> ]quinoxaline
7,8-DiMeIQx	2-amino-3,7,8-trimethylimidazo[4,5- <i>f</i> ]quinoxaline
4-CH <sub>2</sub> OH-8-MeIQx	2-amino-4-hydroxymethyl-3,8-dimethylimidazo[4,5- <i>f</i> ]quinoxaline
PhIP	2-amino-1-methyl-6-phenylimidazo[4,5- <i>b</i> ]pyridine
4'-hydroxy-PhIP	2-amino-6-(4-hydroxyphenyl)-1-methylimidazo[4,5- <i>b</i> ] pyridine
Trp-P-1	3-amino-1,4-dimethyl-5 <i>H</i> -pyrido[4,3- <i>b</i> ]indole
Trp-P-2	3-amino-1-methyl-5 <i>H</i> -pyrido[4,3- <i>b</i> ]indole
AαC	2-amino-9 <i>H</i> -pyrido[2,3- <i>b</i> ]indole
MeAαC	2-amino-3-methyl-9 <i>H</i> -pyrido[2,3- <i>b</i> ]indole
Glu-P-1	2-amino-6-methyldipyrido[1,2- <i>a</i> :3'2'- <i>d</i> ]imidazole
Glu-P-2	2-aminodipyridol[1,2- <i>a</i> :3'2'- <i>d</i> ]imidazole
Harman	1-methyl-9 <i>H</i> -pyrido[3,4- <i>b</i> ]indole
Norharman	9 <i>H</i> -pyrido[3,4- <i>b</i> ]indole
IgQx	2-amino-1-methylimidazo[4,5- <i>g</i> ]quinoxaline

Note: the chemical structure of some of these compounds is given in [Fig. 1.2](#)

Adapted with permission from [Sugimura et al. \(2004\)](#) and [Alaejos & Afonso \(2011\)](#)

### (iii) Polycyclic aromatic hydrocarbons

The main source of non-occupational human exposure, for non-smoking individuals, is food consumption ([Kazerouni et al., 2001](#)). PAHs can be formed by pyrolysis of organic materials, direct contact of fat with a flame, or incomplete combustion of charcoal, so they are present in grilled meats ([Chen & Lin, 1997](#); [Alomirah et al., 2011](#)). More than 30 PAHs have been identified; among them is benzo[*a*]pyrene (BaP), which is classified as a Group 1 human carcinogen ([IARC, 2012a](#)). The main PAHs found in processed meats are presented in [Fig. 1.3](#) and [Table 1.7](#). Representative concentrations of PAHs in different processed meat samples are given in [Table 1.8](#).

By avoiding the direct contact of meat with a flame, PAH levels can be lowered. The amount of fat can also influence PAH levels. The more fat that is contained in meat, the more PAHs are

produced. This may be related to the pyrolysis of fat, which drips onto the heat source ([Mottier et al., 2000](#)).

Heat treatment of red and processed meat can also produce other toxicants, such as acrylamide ([Tareke et al., 2002](#)) and *N*-methylacrylamide ([Yaylayan et al., 2004](#)).

### (iv) Iron

Iron is a trace element essential for human health that can be found in foods of animal and plant origin. In food, iron can be found in two forms: haem iron and non-haem iron. Haem iron, which is more bioavailable than non-haem iron, is only found in animal products ([Schonfeldt & Hall, 2011](#)). Haem iron is contained in myoglobin and haemoglobin, whereas non-haem iron is associated with small molecules such as phosphate, ascorbate, or free amino acids to form salts. The amount of iron in meat, and the ratio between haem and non-haem iron, depends on

**Table 1.6 Concentrations of heterocyclic aromatic amines in different cooked meats**

Cooked meat	Concentrations of HAAs (ng/g)									
	IQ	MeIQ	8-MeIQx	4,8-DiMeIQx	PhIP	Trp-P-1	Trp-P-2	Harman	Norharman	AaC
Minced beef (fried, grilled, and barbecued)	ND-12	ND-8	ND-7	ND-3	ND-34	ND to < 1.45	ND-2	ND-28	ND-30	ND-21
Beef (roasted and oven-broiled)	ND to < 0.2	ND to < 0.2	ND-17.5	ND-3.4	ND-32.4	ND-0.01	ND	ND-240 <sup>a</sup>	ND-205 <sup>b</sup>	ND-0.11
Beef extract (products commercially cooked)	ND-75	ND-10	ND-38	ND-6	ND-10	ND-13	ND-14	NQ-377	NQ-94	NQ to < 8.1
Lamb (grilled and fried)	< 0.1	< 0.1	ND-3	ND-2	ND-11 <sup>b</sup>	ND-1	< 0.3	ND-7	ND-9	ND-0.5
Pork (grilled and fried)	ND-7	ND-11	ND-21	ND-28	ND-32	ND-1	ND-5	ND-25 <sup>a</sup>	ND-51 <sup>a</sup>	ND-3
Sausage (fried, roasted, and barbecued)	ND-5	ND-2	ND-5	ND-3	ND-6	ND-1	ND-2	ND-3	ND-10	< 0.03
Bacon (fried)	ND-11	ND-2	ND-27	ND-9	ND-106 <sup>b</sup>	0.6	< 0.29	ND-33	ND-60	NQ to < 0.5
Pan scrapings from different meats	< 2		29-63	4-15	83-144					3-77


<sup>a</sup> The highest levels of harman and norharman were found in commercially roasted beef ([Khan et al., 2008](#))

<sup>b</sup> A study in the Republic of Korea reported very high concentrations of PhIP (258 ng/g), harman (990 ng/g), and norharman (413 ng/g) in griddled pork loin, and of PhIP (168 ng/g) in griddled bacon ([Back et al., 2009](#))

4,8-DiMeIQx, 2-amino-3,4,8-trimethylimidazo[4,5-*f*]quinoxaline; 8-MeIQx, 2-amino-3,8-dimethylimidazo[4,5-*f*]quinoxaline; AaC, 2-amino-9*H*-pyrido[2,3-*b*]indole; HAA, heterocyclic aromatic amines; harman, 1-methyl-9*H*-pyrido[3,4-*b*]indole; IQ, 2-amino-3-methylimidazo[4,5-*f*]quinoline; MeIQ, 2-amino-3,4-dimethylimidazo[4,5-*f*]quinoline; ND, not detected; norharman, 9*H*-pyrido[3,4-*b*]indole; NQ, not quantified; PhIP, 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine; Trp-P-1, 3-amino-1,4-dimethyl-5*H*-pyrido[4,3-*b*]indole; Trp-P-2, 3-amino-1-methyl-5*H*-pyrido[4,3-*b*]indole

Table compiled using data from the review of [Alaejos & Afonso \(2011\)](#), indicating the lowest and highest values found for the different HAAs in different heat-processed meats



**Fig. 1.3 Structures of polycyclic aromatic hydrocarbons found in red and/or processed meats**

the species and the type of muscle ([Lombardi-Boccia et al., 2002](#); [Table 1.9](#)). Red meat contains more total iron and haem iron than white meat. Beef, lamb, and horse meat are richer in haem iron and total iron than pork meat. The age of the animal is also important in iron intake, as older animals contain more iron. During cooking, part of haem iron is converted to non-haem iron, depending on the cooking parameters, such as time and temperature ([Lombardi-Boccia et al., 2002](#); [Purchas & Busboom, 2005](#); [Purchas et al., 2006](#)).

(v) *Advanced glycation end products*


Advanced glycation end products (AGEPs) are heat-induced food toxicants, which are protein-bound Maillard reaction products. AGEPs constitute a group of heterogeneous moieties produced endogenously from the non-enzymatic glycation of proteins, lipids, and nucleic acids ([Krause et al., 2003](#); [Goldberg et al., 2004](#)). They are present in several heated foods, such as canned foods and meat products ([Goldberg et al., 2004](#); [Uribarri et al., 2010](#)). The formation of AGEPs is part of the normal metabolism, but if their levels are very high in tissues and in the circulation, they can become pathogenic. Carboxymethyllysine is one of the best-characterized AGEPE compounds, and is frequently used as a marker of AGEPE formation in food. In meat products, carboxymethyllysine ranges from 0.01 to 6.87 mg per 100 g of food (mean, 0.86), and in meat dishes, it ranges from 0.10 to 42.39 mg per 100 g of food (mean, 2.42) ([Hull et al., 2012](#)). AGEPE levels depend on red meat composition ([Goldberg et al., 2004](#); [Chen & Smith, 2015](#)). Indeed, foods high in

protein and lipid content show the highest AGEPE levels, probably due to the large quantity of free radicals released via the various lipid peroxidation reactions that catalyse the formation of AGEPEs during the cooking of meat products. AGEPE formation depends on temperature, method, and duration of heating. The higher the cooking temperature, the more AGEPEs are formed in red and processed meat. Different studies have shown that oven-frying produces more AGEPEs than deep-frying, and broiling produces more AGEPEs than roasting. Boiling produces less AGEPEs ([Goldberg et al., 2004](#); [Chen & Smith, 2015](#)). Cooking duration seems to be less important than the temperature and method, as shown in [Table 1.10](#).

(vi) *N-Glycolylneuraminic acid*

Sialic acids are a family of sugars with a nine-carbon sugar acid. *N*-Glycolylneuraminic acid (Neu5Gc) ([Fig. 1.4](#)) is one of the most common sialic acids and is found in almost all mammals. Humans are genetically deficient in Neu5Gc production and instead metabolically accumulate it from dietary sources, particularly red meat and milk products. However, metabolically accumulated dietary Neu5Gc results in the production of circulating anti-Neu5Gc antibodies, leading to chronic local inflammation ([Hedlund et al., 2008](#)). It has been shown that the amount of Neu5Gc is high in red meats compared with other dietary sources, with beef being the most Neu5Gc-enriched compared with other red meats ([Tangvoranuntakul et al., 2003](#); [Samraj et al., 2015](#); [Table 1.11](#)).

**Fig. 1.4 Structure of *N*-glycolylneuraminic acid (Neu5Gc)**



(b) *Processed meat*

Processed meat can contain additional toxicants, apart from the heat toxicants described for red meat. The addition of nitrate and nitrite generates NOCs, and smoking can generate PAHs.

(i) *N-Nitroso compounds*

Processed meat products can be contaminated with NOCs such as *N*-nitrosamines, which result from the reaction between a nitrosating agent, originating from nitrite or smoke, and a secondary amine, derived from protein and lipid degradation (Preussmann & Stewart, 1984; De Mey et al., 2015). *N*-Nitrosamine production is dependent on reaction conditions (e.g. low pH and high temperature), and on meat composition and processing (e.g. ageing, ripening, fermentation, smoking, heat treatment, and storage) (Stadler & Lineback, 2009; Sindelar & Milkowski, 2012; De Mey et al., 2015). NOCs can also be formed endogenously after consumption of red or processed meat (Santarelli et al., 2008).

The most commonly found *N*-nitrosamines in processed meat are *N*-nitrosodimethylamine (NDMA), *N*-nitrosodiethylamine (NDEA), *N*-nitrosopiperidine (NPIP), and *N*-nitrosopyrrolidine (NPYR) (Fig. 1.5; Table 1.12). The concentrations of some of these NOCs in

representative processed meats are given in Table 1.13.

A recent study detected *N*-nitrosamines in dry fermented sausages; only NPIP and *N*-nitrosomorpholine (NMOR) were detected in a high number of samples ( $n = 101$ ; 22% and 28%, respectively). When *N*-nitrosamines were detected, their total amount remained below 5.5  $\mu\text{g}/\text{kg}$ , with only one exception at 14  $\mu\text{g}/\text{kg}$  (De Mey et al., 2014).


The addition of sodium ascorbate to meat, and to a lesser extent NaCl, was shown to decrease *N*-nitrosamine levels (e.g. NDMA and NDEA) in processed meat. On the contrary, baking processes increased *N*-nitrosamine levels (Rywotycki, 2007). [The Working Group noted that since the levels of nitrate and nitrite allowed in cured products are being lowered in many countries, a decrease in NOC formation is expected compared with previous decades.]

(ii) *Polycyclic aromatic hydrocarbons*

Traditional commercial smoking techniques, in which smoke from incomplete wood burning comes into direct contact with the product, can lead to significant contamination by PAHs if the process is not adequately monitored. Temperature, time, humidity, type of smoke used, and even the design of the smokehouse are crucial parameters in controlling PAH formation (EFSA, 2008; Roseiro et al., 2011). The concentrations of selected PAHs in different smoked meats are given in Table 1.8.

PAHs have also been found in dry fermented sausages in Portugal. The concentrations of chrysene, benzo[*a*]anthracene, BaP, and benzo[*a*]fluoranthene were 5.1–38.11, 8–32.9, 1.2–6.6, and 0.63–7.4  $\mu\text{g}/\text{kg}$  dry matter, respectively (Roseiro et al., 2011).

The use of liquid smoke flavouring might reduce PAH levels in commercially smoked meat products (EFSA, 2008).

**Fig. 1.5 Structures of N-nitroso compounds commonly found in processed meats**

The full chemical names of these compounds are given in [Table 1.12](#)

## 1.3 Exposure via food intake

### 1.3.1 Data description

Consumption for a given food depends on two parameters: size of the portion and frequency of eating. In addition, the overall dietary pattern is based on types of foods consumed, which depends on socioeconomic factors (e.g. age, ethnicity, geographical origin, religion, level of education, and income). As a result of these difficulties, food consumption can be estimated using two different techniques: per capita studies and individual surveys, which can, respectively, underestimate or overestimate long-term dietary exposures.

Food consumption results can also be generated using household budget surveys, which correspond to per capita estimates at the household level. However, as the data for household budget surveys are weak, they will not be further considered in this *Monograph*.

In epidemiological studies, food frequency questionnaires (FFQs) are typically used for ranking subjects according to food or nutrient intake, rather than for estimating absolute levels of intake ([Beaton, 1994](#); [Kushi, 1994](#); [Sempos et al., 1999](#)). These questionnaires are further discussed in Section 1.4.1.

### (a) Per capita consumption from economic surveys

The per capita consumption is calculated as follows: national production figures plus imports, minus exports, divided by the total number of individuals in the population. The average values are collected by the Food and Agriculture Organization of the United Nations Statistical Databases (FAOSTAT) ([FAO, 2015](#)) on a yearly basis, and may provide a superior estimate of long-term consumption. However, the per capita data underestimate the true consumption of food items, as less than 100% of the population are consumers, and the whole population is used to calculate the data. On the contrary, for food items consumed by 100% of the population, the data correctly account for both the amount consumed and the frequency of consumption. Based on the FAO per capita data, the World Health Organization (WHO) generated the Global Environment Monitoring System (GEMS) cluster diets ([WHO, 2015a](#)) using a mathematical technique to group countries with similar dietary patterns ([Sy et al., 2013](#)). Consumption values were calculated for each cluster as the average consumption of the food commodity in each country of the cluster. The range of values was therefore narrower than those for FAO national per capita consumption.

**Table 1.7 Polycyclic aromatic hydrocarbons cited in this *Monograph***

Common name (name used in this volume)	CAS registry No.
Benz[ <i>a</i> ]anthracene	56-55-3
Benzo[ <i>b</i> ]fluoranthene	205-99-2
Benzo[ <i>j</i> ]fluoranthene	205-82-3
Benzo[ <i>k</i> ]fluoranthene	207-08-9
Benzo[ <i>c</i> ]fluorene	205-12-9
Benzo[ <i>ghi</i> ]perylene	191-24-2
Benzo[ <i>a</i> ]pyrene	50-32-8
Chrysene	218-01-9
Cyclopenta[ <i>cd</i> ]pyrene	27208-37-3
Dibenz[ <i>a,h</i> ]anthracene	53-70-3
Indeno[1,2,3- <i>cd</i> ]pyrene	193-39-5
Dibenzo[ <i>a,e</i> ]pyrene	192-65-4
Dibenzo[ <i>a,h</i> ]pyrene	189-64-0
Dibenzo[ <i>a,i</i> ]pyrene	189-55-9
Dibenzo[ <i>a,l</i> ]pyrene	191-30-0
Indeno[1,2,3- <i>cd</i> ]pyrene	193-39-5
5-Methylchrysene	3697-24-3

Note: the chemical structure of some of these compounds is given in [Fig. 1.3](#)

### (b) Individual food consumption data

Individual food consumption data are generated from surveys based on recall or recording of daily consumption over 1–7 days. This method allows the distribution of consumption across a population and the consumption of high consumers to be estimated. The method overestimates long-term consumption by extrapolating data collected over a short period of time ([Tran et al., 2004](#); [IPCS, 2009](#)).

### 1.3.2 Results

#### (a) Total meat consumption

For total per capita meat consumption worldwide in 2011, important differences were observed between regions consuming high quantities of meat (i.e. Oceania, 318 g/day; north America, 315 g/day; south America, 215 g/day; Europe, 208 g/day; central America, 148 g/day) and regions consuming low quantities

of meat (i.e. Asia, 86 g/day; Africa, 51 g/day) ([FAO, 2015](#)).

In the European Prospective Investigation into Cancer and Nutrition (EPIC) study, surveys not representative of the national population were conducted in 10 European countries. Food consumption was estimated based on one 24-hour dietary recall ([Linseisen et al., 2002](#)). This study concluded that for total meat, the lowest mean consumption in Europe was observed in Greece (47 g/day for women and 79 g/day for men), and the highest mean consumption was observed in Spain (124 g/day for women and 234 g/day for men) ([Linseisen et al., 2002](#)).

According to FAOSTAT, from 2003 to 2011, meat consumption increased in all regions, but most significantly in Asia (16%) and in Africa (20%). These figures were for both red and poultry meats, and for both processed and unprocessed meats ([FAO, 2015](#)).

**Table 1.8 Concentration levels ( $\mu\text{g}/\text{kg}$ ) of selected polycyclic aromatic hydrocarbon in samples of white, red, and processed meat**

PAH	Cooking/ processing method	Beef		Pork		Chicken	
		Range	Mean	Range	Mean	Range	Mean
Benzo[ <i>k</i> ]fluoranthene (BkF)	Smoked	1.03–3.35	2.57	0.65–4.69	2.96	1.13–4.01	3.54
	Grilled	0.35–2.04	1.87	0.22–3.56	1.37	< 0.10–1.95	1.29
	Boiled	< 0.10–1.81	1.09	0.36–1.45	1.01	0.12–1.54	1.19
	Unprocessed	ND	ND	ND	ND	ND	ND
Benzo[ <i>a</i> ]pyrene (BaP)	Smoked	< 0.10–5.43	5.34	0.50–10.02	1.28	< 0.10–5.91	2.91
	Grilled	0.17–2.93	2.74	0.21–5.73	1.75	0.48–3.73	1.82
	Boiled	0.27–1.30	0.87	0.17–1.45	0.94	< 0.10–1.66	0.99
	Unprocessed	1.71–2.42	0.34	ND	ND	ND	ND
Indeno[123- <i>cd</i> ]pyrene (IP)	Smoked	1.82–27.59	5.10	8.81–31.11	5.29	1.40–7.17	1.39
	Grilled	1.34–8.48	0.62	1.65–8.59	4.01	1.07–3.42	0.61
	Boiled	0.41–1.22	0.54	0.54–1.81	0.97	0.34–1.19	0.45
	Unprocessed	1.32–7.86	3.16	0.27–3.06	1.73	0.21–1.08	0.45
Benzo[ <i>ghi</i> ]perylene (BghiP)	Smoked	< 0.30–2.55	1.42	< 0.30–3.18	1.09	0.88–3.41	2.68
	Grilled	0.61–1.64	1.50	0.78–2.66	1.84	< 0.30–2.56	1.34
	Boiled	0.36–1.19	0.82	< 0.30–1.62	0.93	< 0.30–1.87	1.12
	Unprocessed	ND	ND	ND	ND	ND	ND

Adapted from *Food chemistry*, Volume 156, [Olatunji et al. \(2014\)](#). Determination of polycyclic aromatic hydrocarbons [PAHs] in processed meat products using gas chromatography – Flame ionization detector, Pages No. 296–300, Copyright (2014), with permission from Elsevier

(b) *Association between consumption of red meat and consumption of other foods*

Food categories are not independent in regard to consumption. In the field of nutrition, nutrient intake is estimated by combining consumption data with food nutrient composition databases. Thereafter, homogeneous subgroups of consumers with comparable nutrient intakes (dietary patterns) are identified by using classical statistical clustering techniques ([Pryer et al., 2001](#); [Hu, 2002](#)). The association between food categories can also be observed by using principal component analysis. For example, intake of processed meat was associated with intake of French fries, sweets, cakes, desserts, snacks, and alcoholic beverages ([Fung et al., 2003](#); [Dixon et al., 2004](#); [Kesse et al., 2006](#)).

Whereas clustering is based on nutrient intake, it is very difficult to a posteriori identify foods that contribute by a majority to a given dietary pattern. Zetlaoui et al. proposed the use of

principal component analysis for food clustering ([Zetlaoui et al., 2011](#)). Based on this approach, and its application in the FAO per capita data set (i.e. 415 food products in 179 countries), 30 consumption systems leading to 17 cluster diets have been described ([Sy et al., 2013](#)). According to this publication, the consumption of pork meat seemed to be associated with the consumption of barley beer, poultry meat, wheat flour, and refined sugar. The consumption of cattle meat seemed to be associated with cow milk and wheat flour ([Sy et al., 2013](#)).

(c) *Red meat consumption*

According to FAOSTAT in 2011, the cumulated mean per capita consumption of beef, mutton, goat, and pig meat was 30, 60, 130, 140, and 200 g/day, respectively, for Africa, Asia, America, Europe, and Oceania ([FAO, 2015](#)). From the WHO/GEMS clusters, the average

**Table 1.9 Total iron and percentage of haem iron in raw and cooked meat**

Meats	Total iron (mg/100 g)		% Haem iron		% Loss
	Raw	Cooked	Raw	Cooked	
<i>Red meat</i>					
Beef					
Sirloin	2.07	3.59	83	74	11
Fillet	2.35	3.38	90	85	6
Roasted beef	2.04	3.74	87	84	3
Topside	1.93	2.88	87	66	24
Mean	2.09	3.39	87	78	11
Veal					
Fillet	0.85	1.58	84	83	1
Lamb					
Chop	2.23	3.20	75	70	7
Horse					
Fillet	2.21	3.03	79	71	11
Pork					
Loin	0.36	0.46	56	46	18
Chump chop	0.49	0.79	66	69	(+4)
Mean	0.42	0.64	62	61	7
<i>White meat</i>					
Chicken					
Breast	0.40	0.58	30	28	7
Leg (thigh)	0.70	1.34	30	22	27
Leg (lower part)	0.63	1.20	46	35	24
Wing	0.63	0.92	44	25	43
Mean	0.59	1.01	38	28	28
Turkey					
Breast	0.50	0.79	28	27	4
Leg (thigh)	0.99	1.46	50	39	22
Leg (lower part)	0.88	1.51	49	38	22
Mean	0.79	1.25	42	35	18

Adapted from [Lombardi-Boccia et al. \(2002\)](#)

total red meat consumption ranged from 15 to 147 g/day ([WHO, 2013](#)).

In a systematic assessment, the Global Burden of Diseases Nutrition and Chronic Diseases Expert Group (NutriCoDE) evaluated the global consumption of key dietary items (foods and nutrients) by region, nation, age, and sex in 1990 and 2010 ([Imamura et al., 2015](#)). Consumption data were evaluated from 325 surveys (71.7% nationally representative) covering 88.7% of the global adult population. According to the analysis, the median of mean consumption

of red meat worldwide ranged from 23 g/day (2.6–28 g/day) for the first quintile to 84 g/day (71–138 g/day) for the fifth quintile ([Imamura et al., 2015](#)).

Individual food consumption surveys provide the distribution of consumption for consumers only (i.e. high percentiles of consumption as well as percentages of consumers by country) ([FAO/WHO, 2015](#); [FCID, 2015](#)). Worldwide detailed data on red meat consumption (g/kg bw per day) are presented in [Table 1.14](#) and [Table 1.15](#) for adults and children, respectively.

**Table 1.10 Advanced glycation end product content in red meat, processed meat, and chicken<sup>a</sup>**

Meat	Cooking/processing method	Advanced glycation end product (kU/110 g)	
Beef	Raw	707	
	Roast	6 071	
	Steak, raw	800	
	Steak, broiled	7 479	
	Steak, grilled 4 min	7 416	
	Steak, microwaved, 6 min	2 687	
	Steak, pan fried w/olive oil	10 058	
	Steak, strips, 450°F, 15 min	6 851	
	Steak, strips, stir fried with 1 T canola oil, 15 min	9 522	
	Steak, strips, stir fried without oil, 7 min	6 973	
	Stewed	2 443	
	Frankfurter, boiled in water, 212° F, 7 min	7 484	
	Frankfurter, broiled 450°F, 5 min	11 270	
	Ground, 20% fat, pan/cover	5 527	
	Hamburger patty, olive oil 180°F, 6 min	2 639	
	Meatball, potted (cooked in liquid), 1 h	4 300	
	Meatball, w/sauce	2 852	
	Meatloaf, crust off, 45 min	1 862	
	Pork	Bacon, fried 5 min no added oil	91 577
		Bacon, microwaved, 2 slices, 3 min	9 023
Ham, deli, smoked		2 349	
Liverwurst		633	
Chop, pan fried, 7 min		4 752	
Ribs, roasted		4 430	
Roast (Chinese take-out)		3 544	
Sausage, beef and pork links, pan fried		5 426	
Sausage, Italian, raw		1 861	
Sausage, Italian, barbecued		4 839	
Sausage, pork links, microwaved, 1 min		5 943	
Lamb		Leg, raw	826
	Leg, boiled, 30 min	1 218	
	Leg, broiled, 450°F, 30 min	2 431	
	Leg, microwave, 5 min	1 029	
Veal	Stewed	2 858	
Chicken	Ground, white meat, raw	877	
	Meatball, potted (cooked in liquid) 1 h	1 501	
	Potted (cooked in liquid) with onion and water	3 329	
	Roasted	6 020	
	Skin, back of thigh, roasted then barbecued	18 520	

<sup>a</sup> Glycation end product content based on carboxymethyllysine content

Adapted from *Journal of the American Dietetic Association*, Volume 110, issue 6, Jaime Uribarri, Sandra Woodruff, Susan Goodman, Weijing Cai, Xue Chen, Renata Pyzik, Angie Yong, Gary E. Striker, Helen Vlassara, Advanced Glycation End Products in Foods and a Practical Guide to Their Reduction in the Diet, Pages No. 911-916.e12, Copyright (2010), with permission from Elsevier ([Uribarri et al., 2010](#))

**Table 1.11 Content of *N*-glycolylneuraminic acid in red or processed meat, raw or cooked under different conditions**

Meat	Neu5Gc content (µg/g)
Ground beef	25
Beef steak (raw)	134
Beef steak (baked)	210
Beef steak (boiled)	231
Beef steak (fried)	199
Ground lamb	14
Lamb steak (raw)	57
Lamb steak (baked)	50
Lamb steak (boiled)	47
Lamb steak (fried)	19
Ground pork	19
Pork chop (raw)	25
Pork chop (baked)	40
Pork chop (boiled)	36
Pork chop (fried)	29
Pork bratwurst	11
Pork bacon	7

Neu5Gc, *N*-glycolylneuraminic acid

From [Samraj et al. \(2015\)](#), with permission of the editor

### (i) Europe

The European Food Safety Authority (EFSA) collected in a harmonized way the results from national food consumption surveys of more than 20 member states of the EU. The median of mean meat consumption for adults was 35 g/day, ranging from about 10 g/day (Sweden) to 110 g/day (Austria). At the 95th percentile, meat consumption ranged from 20 g/day (Sweden, 21% of consumers) to 237 g/day (Austria, 88% of consumers). Similar results were found for adolescents, both in terms of amount consumed and percentage of the population that are consumers. For infants and toddlers, the median of mean meat consumption was about 50 g/day, ranging from 20 to 80 g/day, and the percentage of consumers ranged from about 5% (the Netherlands) to 100% (Germany). At the 95th percentile, the meat consumption ranged from 40 g/day (the Netherlands) to about 190 g/day (Belgium) ([EFSA, 2011](#)).

The EPIC study concluded that red meat consumption ranged from 24 to 57 g/day for women and from 40 to 121 g/day for men based on 24-hour recall ([Linseisen et al., 2002](#)).

### (ii) Americas

Few representative national surveys were available for the Americas. In the USA ([FCID, 2015](#)), the mean consumption of total red meat was 86 g/day for adult consumers and 242 g/day at the 95th percentile for the same population (72% consumers). For children aged between 1 and < 3 years, mean consumption was 31 g/day and 89 g/day at the 95th percentile (62% consumers). For children aged between 3 and 16 years, mean consumption was 60 g/day and 176 g/day at the 95th percentile (71% consumers) ([FCID, 2015](#)). Similarly, in Brazil, the mean consumption of beef was 92 g/day for the general population and 232 g/day at the 95th percentile for the same population (69% consumers). No data were available for children in Brazil, and no data were available for other countries in Latin America. However, according to the GEMS Cluster diets, the dietary patterns in this region seemed homogeneous ([FAO/WHO, 2015](#); [Table 1.14](#) and [Table 1.15](#)).

### (iii) Africa

Data were scarce and incomplete for Africa. Fortunately, individual food consumption surveys were performed for adult women and children in Burkina Faso and Uganda. In these two countries, the percentage of consumers of red meat was less than 5% of the population. However, for these adult consumers, the mean consumption was between 23 and 90 g/day, and consumption at the 95th percentile was between 28 and 147 g/day. For children, the percentage of consumers of red meat was below 4% of this population. Similarly, compared with adults, consumption for child consumers was close to that observed in developed countries, with a mean between 13 and 62 g/day, and a high consumption



**Table 1.12 N-Nitroso compounds commonly found in processed meat**

Common abbreviation	Full name	CAS registry No.
NMDA	N-Nitrosodimethylamine	62-75-9
NDEA	N-Nitrosodiethylamine	55-18-5
NPIP	N-Nitrosopiperidine	100-75-4
NPYR	N-Nitrosopyrrolidine	930-55-2
NDBA	N-Nitrosodi-n-butylamine	924-16-3
–	N-Nitrosomethylethylamine	10595-95-6
–	N-Nitrosoproline	7519-36-0
–	N-Nitrosohydroxyproline	30310-80-6
NMOR	N-Nitrosomorpholine	59-89-2

CAS, Chemical Abstracts Service

The chemical structure of some of these compounds is given in [Fig. 1.5](#)

at the 95th percentile of between 22 and 69 g/day. It is therefore likely that the difference in the per capita consumption (four to five times lower in Africa than in Europe) was mainly due to a lower number of consumers rather than to large differences in the dietary patterns of consumers ([FAO/WHO, 2015](#); [Table 1.14](#) and [Table 1.15](#)).

(iv) *Middle East and north Africa*

Intake of red meat in countries of the Middle East and north Africa was estimated in 2010 to range from 200 g/week (Afghanistan) to 700 g/week (Algeria and United Arab Emirates) ([Afshin et al., 2015](#)).

(v) *Asia*

Food consumption surveys were available from Bangladesh, China, Japan, the Philippines, the Republic of Korea, and Thailand. In Asia, the main types of red meat consumed were pork and beef ([FAO/WHO, 2015](#)). In China, the predominant red meat consumed was pork, with 63% of consumers, a mean consumption of 84 g/day, and consumption at the 95th percentile of 224 g/day for adult consumers only ([Table 1.14](#)). Based on three consecutive 24-hour recalls, a prospective study of 5000 adults from 4280 households in nine provinces showed an increase in average consumption of pork of 20% (52 vs 62 g/day per person) from 1989 to 2004

([Zhai et al., 2009](#)). For Chinese children, the mean consumption of pork was 51 g/day, and consumption of pork at the 95th percentile was 142 g/day. Beef was consumed by less than 10% of the Chinese population, with a mean consumption of 46 g/day and consumption at the 95th percentile for consumers of 130 g/day. For children, the mean consumption of beef was 32 g/day, and consumption of beef at the 95th percentile was 85 g/day. These figures were close to those reported in the Americas and Europe ([FAO/WHO, 2015](#); [Table 1.15](#)).

Similarly, in the Republic of Korea, the consumption of pork for adults was 76 g/day, and consumption of pork at the 95th percentile was 253 g/day (44% of consumers). For children, the mean consumption of pork was 30 g/day, and consumption of pork at the 95th percentile was 95 g/day. Finally, in the Philippines, for one third of the population, the mean consumption of pork for children was 75 g/day, and consumption of pork at the 95th percentile was 208 g/day (33% of consumers). On the contrary, in Japan, beef and pork were consumed by a wide range of consumers (i.e. 89% and 99% of the population, respectively). The mean consumption and consumption at the 97.5th percentile for consumers only were 53 and 83 g/day, respectively, i.e. about half of the consumption in north

**Table 1.13 Major sources of dietary *N*-nitrosamines in processed meats**

Processed meat	Concentration of nitrosamines (µg/kg)			
	NDMA	NDEA	NPYR	NPIP
Bacon fried	ND-30	ND-1	ND-200	ND-1
Cured meats	ND-4	ND-4	ND-25	ND-2
Smoked meats	ND-3	ND-7.9	ND-0.1	ND-0.1
<i>Sausages</i>				
Frankfurter	ND-84	-	-	-
Mettwurst	+	+	ND-105	ND-60
Liver sausage	ND-35	ND-25	ND-80	-
Salami	ND-80	-	-	-
Bologna	-	ND-25	ND-105	-

+, detected but not quantitated; -, not reported; ND, not detected; NDMA, *N*-nitrosodimethylamine; NDEA, *N*-nitrosodiethylamine; NPIP, *N*-nitrosopiperidine; NPYR, *N*-nitrosopyrrolidine

From: *Nitrates, nitrites and N-nitrosocompounds: A review of the occurrence in food and diet and the toxicological implications*, R. Walker, Food Additives & Contaminants, 1990, reprinted by permission of Taylor & Francis (Taylor & Francis Ltd, <http://www.tandfonline.com>) (Walker, 1990)

America or in China. In Thailand, the percentage of pork meat consumers was 89%, with a mean consumption of 23 g/day. In Bangladesh, the percentage of red meat consumers was less than 10%. The mean consumption for consumers was between 10 and 23 g/day, and the consumption at the 95th percentile was between 25 and 77 g/day (FAO/WHO, 2015; Table 1.14 and Table 1.15).

#### (iv) Oceania

The 2008/09 New Zealand Adult Nutrition Survey (University of Otago and Ministry of Health, 2011) estimated the mean consumption of beef and veal to be 180 g/day, and consumption of beef and veal at the 90th percentile to be 397 g/day for consumers only. The same survey estimated the mean consumption of lamb and mutton to be 137 g/day, and consumption of lamb and mutton at the 90th percentile to be 275 g/day. For these two food categories, the percentage of consumers was 24% for beef and veal meat, and 7% for lamb and goat meat (Parnell et al., 2012). Data on the consumption of pork, as well as the total red meat consumption, were not available for adults. For Australia, data on consumption were only available for children. They showed

a mean consumption that for consumers only ranged from 13 to 70 g/day, and a consumption at the 97.5th percentile that ranged from 83 to 257 g/day (FAO/WHO, 2015; Table 1.15).

In summary, for most countries (e.g. Australia, central and southern Europe, China, the Philippines, the Republic of Korea, and the USA), the mean consumption of red meat for consumers only was around 50–100 g/day, and high consumption was around 200–300 g/day. The percentage of meat consumers seemed to be proportional to the income or the level of development. In other words, the distribution of meat consumption was fairly similar among consumers in these countries. Therefore, analysis of per capita data only may give the wrong perception of the levels of consumption. In some countries (e.g. Japan, northern Europe, and Thailand), the consumption of red meat was low, despite a percentage of consumers of about 90%, probably due to substitution with fish and other seafoods. Finally, in less-industrialized countries for which data were available (e.g. Bangladesh, Burkina Faso, and Uganda), the percentage of consumers was below 10%, probably due to the high price of red meat. It should be noted that, in these countries, the mean and high

**Table 1.14 Worldwide consumption of red meat in adults**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, Mean (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Adult women	Bangladesh	Harvest_2007/8	Beef and other bovines meat	474	39	8.23%	0.4	0.3	0.9
Adult women	Bangladesh	Harvest_2007/8	Goat and other caprines	474	9	1.90%	0.5	0.7	2.0
Adults	Belgium	Diet_National_2004	Beef and other bovines meat	1304	449	34.43%	0.9	0.7	2.5
Adults	Belgium	Diet_National_2004	Horse and other equines	1304	16	1.23%	1.1	0.7	2.6
Adults	Belgium	Diet_National_2004	Meat from mammals other than marine mammals, NES	1304	9	0.69%	0.7	0.3	1.1
Adults	Belgium	Diet_National_2004	Pork and other porcines	1304	273	20.94%	0.9	0.5	2.3
Adults	Belgium	Diet_National_2004	Sheep and other ovines	1304	84	6.44%	0.9	0.5	2.1
General population	Brazil	Brazilian Institute of Geography and Statistics	Beef and other bovines meat	34 003	23 320	68.58%	1.4	1.2	4.4
General population	Brazil	Brazilian Institute of Geography and Statistics	Goat and other caprines	34 003	194	0.57%	1.8	1.2	4.8
General population	Brazil	Brazilian Institute of Geography and Statistics	Meat from mammals other than marine mammals, NES	34 003	2071	6.09%	1.0	0.9	3.5
General population	Brazil	Brazilian Institute of Geography and Statistics	Pork and other porcines	34 003	2577	7.58%	1.8	1.7	6.3
General population	Brazil	Brazilian Institute of Geography and Statistics	Sheep and other ovines	34 003	136	0.40%	1.5	1.1	4.8
Adult women	Burkina Faso	Harvest_2010	Beef and other bovines meat	287	7	2.44%	0.4	0.1	0.5
Adult women	Burkina Faso	Harvest_2010	Goat and other caprines	287	7	2.44%	0.7	0.5	1.5
Adult women	Burkina Faso	Harvest_2010	Meat from mammals other than marine mammals, NES	287	3	1.05%	1.7	0.5	2.2

**Table 1.14 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, Mean (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Adult women	Burkina Faso	Harvest_2010	Pork and other porcines	287	11	3.83%	0.8	0.5	1.8
Adult women	Burkina Faso	Harvest_2010	Sheep and other ovines	287	4	1.39%	0.9	0.3	1.2
General population	China	2002 China Nutrition and Health Survey	Beef and other bovines meat	65 359	5278	8.08%	0.9	0.9	3.2
General population	China	2002 China Nutrition and Health Survey	Horse and other equines	65 359	66	0.10%	2.1	4.0	10.4
General population	China	2002 China Nutrition and Health Survey	Meat from mammals other than marine mammals, NES	65 359	635	0.97%	1.2	1.2	4.6
General population	China	2002 China Nutrition and Health Survey	Pork and other porcines	65 359	41 283	63.16%	1.6	1.4	5.3
General population	China	2002 China Nutrition and Health Survey	Sheep and other ovines	65 359	3690	5.65%	1.2	1.2	4.3
Adults	Czech Republic	SISP04	Beef and other bovines meat	1666	514	30.85%	0.8	0.6	2.2
Adults	Czech Republic	SISP04	Pork and other porcines	1666	694	41.66%	1.1	0.7	2.9
Adults	Denmark	Danish_Dietary_Survey	Beef and other bovines meat	2822	2780	98.51%	0.5	0.4	1.4
Adults	Denmark	Danish_Dietary_Survey	Pork and other porcines	2822	2750	97.45%	0.6	0.5	1.8
Adults	Denmark	Danish_Dietary_Survey	Sheep and other ovines	2822	187	6.63%	0.4	0.2	1.0
Adults	Finland	FINDIET_2007	Beef and other bovines meat	1575	695	44.13%	0.6	0.5	2.0
Adults	Finland	FINDIET_2007	Pork and other porcines	1575	431	27.37%	0.7	0.6	2.4
Adults	Finland	FINDIET_2007	Sheep and other ovines	1575	62	3.94%	0.6	0.6	1.8
Adults	France	INCA2	Beef and other bovines meat	2276	2002	87.96%	0.7	0.5	1.8
Adults	France	INCA2	Horse and other equines	2276	52	2.28%	0.3	0.2	0.8
Adults	France	INCA2	Meat from mammals other than marine mammals, NES	2276	825	36.25%	0.1	0.2	0.5

**Table 1.14 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, Mean (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Adults	France	INCA2	Pork and other porcines	2276	1154	50.70%	0.3	0.3	1.1
Adults	France	INCA2	Sheep and other ovines	2276	627	27.55%	0.2	0.2	0.7
Adults	Hungary	National_Repr_Surv	Beef and other bovines meat	1074	382	35.57%	0.3	0.3	1.1
Adults	Hungary	National_Repr_Surv	Pork and other porcines	1074	860	80.07%	0.9	0.6	2.6
Adults	Hungary	National_Repr_Surv	Sheep and other ovines	1074	8	0.74%	0.5	0.2	0.8
Adults	Ireland	NSIFCS	Beef and other bovines meat	958	761	79.44%	0.7	0.6	2.0
Adults	Ireland	NSIFCS	Pork and other porcines	958	427	44.57%	0.5	0.4	1.5
Adults	Ireland	NSIFCS	Sheep and other ovines	958	361	37.68%	0.4	0.3	1.4
Adults	Italy	INRAN_SCAI_2005_06	Beef and other bovines meat	2313	1698	73.41%	0.8	0.6	2.3
Adults	Italy	INRAN_SCAI_2005_06	Goat and other caprines	2313	3	0.13%	0.6	0.2	0.8
Adults	Italy	INRAN_SCAI_2005_06	Horse and other equines	2313	57	2.46%	0.7	0.4	1.5
Adults	Italy	INRAN_SCAI_2005_06	Pork and other porcines	2313	735	31.78%	0.6	0.5	1.8
Adults	Italy	INRAN_SCAI_2005_06	Sheep and other ovines	2313	71	3.07%	0.6	0.8	1.4
General population	Japan	DSFFQ_FI	Beef and other bovines meat	2711	2406	88.75%	0.3	0.3	1.0
General population	Japan	DSFFQ_FI	Meat from mammals other than marine mammals, NES	2711	112	4.13%	0.2	0.1	0.7
General population	Japan	DSFFQ_FI	Pork and other porcines	2711	2691	99.26%	0.6	0.4	1.5
Adults	Latvia	EFSa_TEST	Beef and other bovines meat	1306	66	5.05%	0.8	0.6	2.7
Adults	Latvia	EFSa_TEST	Goat and other caprines	1306	1	0.08%	0.7		0.7
Adults	Latvia	EFSa_TEST	Meat from mammals other than marine mammals, NES	1306	20	1.53%	0.7	0.4	1.7
Adults	Latvia	EFSa_TEST	Pork and other porcines	1306	796	60.95%	1.2	0.9	3.5

**Table 1.14 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, Mean (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Adults	Latvia	EFSA_TEST	Sheep and other ovines	1306	5	0.38%	0.8	0.3	1.3
Adults	Netherlands	DNFCS_2003	Beef and other bovines meat	750	180	24.00%	1.0	0.6	2.5
Adults	Netherlands	DNFCS_2003	Horse and other equines	750	2	0.27%	0.2	0.1	0.3
Adults	Netherlands	DNFCS_2003	Pork and other porcines	750	309	41.20%	1.2	0.9	3.8
Adults	Netherlands	DNFCS_2003	Sheep and other ovines	750	30	4.00%	1.2	0.9	4.1
General population	Republic of Korea	KNHNES	Beef and other bovines meat	9391	3141	33.45%	0.9	1.2	4.1
General population	Republic of Korea	KNHNES	Pork and other porcines	9391	4124	43.91%	1.4	1.8	6.3
Adults	Spain	AESAN_FIAB	Beef and other bovines meat	981	680	69.32%	1.1	0.7	2.8
Adults	Spain	AESAN	Beef and other bovines meat	410	176	42.93%	1.2	0.8	3.2
Adults	Spain	AESAN_FIAB	Goat and other caprines	981	3	0.31%	1.1	0.2	1.3
Adults	Spain	AESAN_FIAB	Pork and other porcines	981	366	37.31%	1.0	0.7	3.0
Adults	Spain	AESAN	Pork and other porcines	410	129	31.46%	1.0	0.6	2.4
Adults	Spain	AESAN_FIAB	Sheep and other ovines	981	102	10.40%	1.0	0.5	2.3
Adults	Spain	AESAN	Sheep and other ovines	410	18	4.39%	1.1	0.9	3.8
Adults	Sweden	Riksmaten_1997_98	Beef and other bovines meat	1210	590	48.76%	0.3	0.2	0.9
Adults	Sweden	Riksmaten_1997_98	Horse and other equines	1210	8	0.66%	0.1	0.1	0.5
Adults	Sweden	Riksmaten_1997_98	Pork and other porcines	1210	699	57.77%	0.4	0.2	1.0
Adults	Sweden	Riksmaten_1997_98	Sheep and other ovines	1210	32	2.64%	0.2	0.2	1.0
General population	Thailand	FCDT	Beef and other bovines meat	16 383	7880	48.10%	0.1		
General population	Thailand	FCDT	Pork and other porcines	16 383	14 646	89.40%	0.4		
Adult women	Uganda	Harvest_2007	Beef and other bovines meat	176	8	4.55%	1.2	0.8	2.8
Adult women	Uganda	Harvest_2007	Goat and other caprines	176	2	1.14%	1.2	0.7	1.7
Adults	United Kingdom	NDNS	Beef and other bovines meat	1724	1349	78.25%	0.4	0.3	1.1

**Table 1.14 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, Mean (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Adults	United Kingdom	NDNS	Pork and other porcines	1724	535	31.03%	0.3	0.2	0.9
Adults	United Kingdom	NDNS	Sheep and other ovines	1724	434	25.17%	0.3	0.2	0.8
Adults over 16 years	USA	FCID-WWEIA data for years 2005–2010	Total red meat	31 484	23 825	75.67%	1.1		3.06*
General population	USA	FCID-WWEIA data for years 2005–2010	Total red meat	49 343	35 752	72.46%	1.2		3.59*
General population	USA	FCID-WWEIA data for years 2005–2010	Sheep meat	49 343	2518	5.10%	0.1		0.56*
General population	USA	FCID-WWEIA data for years 2005–2010	Goat meat	49 343	35	0.07%	1.9		5.8*
General population	USA	FCID-WWEIA data for years 2005–2010	Pork meat	49 343	26 256	53.21%	0.55		2.04*
General population	USA	FCID-WWEIA data for years 2005–2010	Beef meat	49 343	29 788	60.37%	0.96		3.08*

\* 95th percentile

NES, not elsewhere specified

Data on USA from [FCID \(2015\)](#): What We Eat In America – Food Commodity Intake Database 2005–10, United States Environmental Protection Agency – Office of Pesticide Programs © University of Maryland 2012 – 2016. Available from: <http://fcid.foodrisk.org/percentiles.php>

Data for other countries from [FAO/WHO \(2015\)](#): the FAO/WHO Chronic individual food consumption database – Summary statistics (CIFOcOss), © Copyright World Health Organization (WHO), 2016. All Rights Reserved. Available from: <http://www.who.int/foodsafety/databases/en/>

**Table 1.15 Worldwide consumption of red meat in children**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Adolescents	Belgium	Diet_National_2004	Beef and other bovines meat	584	175	29.97%	1.0	0.7	2.5
Adolescents	Belgium	Diet_National_2004	Horse and other equines	584	6	1.03%	1.3	0.4	1.9
Adolescents	Belgium	Diet_National_2004	Meat from mammals other than marine mammals, NES	584	11	1.88%	0.7	0.4	1.7
Adolescents	Belgium	Diet_National_2004	Pork and other porcines	584	121	20.72%	1.1	0.7	3.1
Adolescents	Belgium	Diet_National_2004	Sheep and other ovines	584	43	7.36%	1.0	0.7	2.8
Adolescents	Cyprus	Childhealth	Beef and other bovines meat	303	18	5.94%	0.6	0.2	0.9
Adolescents	Cyprus	Childhealth	Pork and other porcines	303	154	50.83%	1.1	0.6	2.8
Adolescents	Cyprus	Childhealth	Sheep and other ovines	303	12	3.96%	0.8	0.4	1.8
Adolescents	Czech Republic	SISP04	Beef and other bovines meat	298	97	32.55%	1.2	0.8	3.1
Adolescents	Czech Republic	SISP04	Pork and other porcines	298	125	41.95%	1.4	0.8	3.2
Adolescents	Denmark	Danish Dietary Survey	Beef and other bovines meat	479	478	99.79%	0.7	0.5	2.0
Adolescents	Denmark	Danish_Dietary_Survey	Pork and other porcines	479	472	98.54%	0.7	0.5	2.0
Adolescents	Denmark	Danish_Dietary_Survey	Sheep and other ovines	479	21	4.38%	0.5	0.3	1.3
Adolescents	France	INCA2	Beef and other bovines meat	973	912	93.73%	0.9	0.6	2.4
Adolescents	France	INCA2	Horse and other equines	973	21	2.16%	0.5	0.3	1.6
Adolescents	France	INCA2	Meat from mammals other than marine mammals, NES	973	424	43.58%	0.2	0.2	1.0
Adolescents	France	INCA2	Pork and other porcines	973	482	49.54%	0.4	0.3	1.2
Adolescents	France	INCA2	Sheep and other ovines	973	257	26.41%	0.3	0.3	0.8
Adolescents	Italy	INRAN_SCAI_2005_06	Beef and other bovines meat	247	204	82.59%	1.2	0.9	3.2



**Table 1.15 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Adolescents	Italy	INRAN_SCAI_2005_06	Horse and other equines	247	8	3.24%	0.8	0.3	1.4
Adolescents	Italy	INRAN_SCAI_2005_06	Pork and other porcines	247	81	32.79%	0.8	0.7	2.5
Adolescents	Italy	INRAN_SCAI_2005_06	Sheep and other ovines	247	2	0.81%	0.7	0.2	0.9
Adolescents	Latvia	EFSA_TEST	Beef and other bovines meat	470	16	3.40%	0.9	0.6	2.3
Adolescents	Latvia	EFSA_TEST	Meat from mammals other than marine mammals, NES	470	4	0.85%	0.5	0.1	0.5
Adolescents	Latvia	EFSA_TEST	Pork and other porcines	470	263	55.96%	1.4	1.1	4.1
Adolescents	Spain	AESAN_FIAB	Beef and other bovines meat	86	62	72.09%	1.4	1.0	4.5
Adolescents	Spain	NUT_INK05	Beef and other bovines meat	651	294	45.16%	1.7	1.1	4.8
Adolescents	Spain	NUT_INK05	Goat and other caprines	651	2	0.31%	1.3	0.5	1.6
Adolescents	Spain	enKid	Horse and other equines	209	2	0.96%	1.0	0.2	1.2
Adolescents	Spain	NUT_INK05	Horse and other equines	651	1	0.15%	1.3		1.3
Adolescents	Spain	enKid	Meat from mammals other than marine mammals, NES	209	69	33.01%	1.5	0.9	3.9
Adolescents	Spain	AESAN_FIAB	Pork and other porcines	86	42	48.84%	1.0	0.6	2.0
Adolescents	Spain	enKid	Pork and other porcines	209	60	28.71%	1.2	0.8	3.3
Adolescents	Spain	NUT_INK05	Pork and other porcines	651	212	32.57%	1.1	0.7	3.3
Adolescents	Spain	AESAN_FIAB	Sheep and other ovines	86	4	4.65%	1.3	0.6	2.0
Adolescents	Spain	enKid	Sheep and other ovines	209	11	5.26%	1.9	2.3	8.5
Adolescents	Spain	NUT_INK05	Sheep and other ovines	651	29	4.45%	1.3	0.7	3.8
Adolescents	Sweden	NFA	Beef and other bovines meat	1018	542	53.24%	0.5	0.4	1.7
Adolescents	Sweden	NFA	Horse and other equines	1018	9	0.88%	0.3	0.3	0.9
Adolescents	Sweden	NFA	Pork and other porcines	1018	286	28.09%	0.8	0.5	2.0
Adolescents	Sweden	NFA	Sheep and other ovines	1018	6	0.59%	0.8	0.8	2.4

**Table 1.15 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Children	Bangladesh	Harvest_2007/8	Beef and other bovines meat	555	41	7.39%	1.0	0.8	3.0
Children	Bangladesh	Harvest_2007/8	Goat and other caprines	555	12	2.16%	0.9	0.7	2.1
Children	Burkina Faso	Harvest_2010	Beef and other bovines meat	288	7	2.43%	0.9	0.6	1.5
Children	Burkina Faso	Harvest_2010	Goat and other caprines	288	6	2.08%	1.8	1.0	3.3
Children	Burkina Faso	Harvest_2010	Meat from mammals other than marine mammals, NES	288	3	1.04%	4.8	3.6	8.0
Children	Burkina Faso	Harvest_2010	Pork and other porcines	288	10	3.47%	2.2	1.6	5.7
Children	Burkina Faso	Harvest_2010	Sheep and other ovines	288	3	1.04%	1.9	0.9	2.8
Children	China	2002 China Nutrition and Health Survey	Beef and other bovines meat	2784	171	6.14%	2.0	1.7	6.7
Children	China	2002 China Nutrition and Health Survey	Horse and other equines	2784	7	0.25%	7.6	10.4	30.9
Children	China	2002 China Nutrition and Health Survey	Meat from mammals other than marine mammals, NES	2784	27	0.97%	2.6	2.6	10.6
Children	China	2002 China Nutrition and Health Survey	Pork and other porcines	2784	1703	61.17%	3.3	2.7	10.5
Children	China	2002 China Nutrition and Health Survey	Sheep and other ovines	2784	119	4.27%	2.8	2.9	10.3
Children	Japan	DSFFQ_FI	Beef and other bovines meat	71	66	92.96%	0.5	0.4	1.3
Children	Japan	DSFFQ_FI	Meat from mammals other than marine mammals, NES	71	1	1.41%	0.1		
Children	Japan	DSFFQ_FI	Pork and other porcines	71	71	100.00%	1.3	0.8	3.7
Children	Philippines	Harvest_2003	Beef and other bovines meat	1205	61	5.06%	1.3	0.8	2.8

**Table 1.15 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Children	Philippines	Harvest_2003	Meat from mammals other than marine mammals, NES	1205	22	1.83%	1.4	0.9	3.1
Children	Philippines	Harvest_2003	Pork and other porcines	1205	395	32.78%	1.4	1.5	5.2
Children	Republic of Korea	KNHNES	Beef and other bovines meat	654	255	38.99%	1.1	1.3	4.9
Children	Republic of Korea	KNHNES	Pork and other porcines	654	329	50.31%	1.9	2.0	6.5
Children (2-16 yrs)	Australia	2007 ANCNPAS	Beef and other bovines meat	4487	3898	86.87%	1.8	1.9	6.8
Children (2-16 yrs)	Australia	2007 ANCNPAS	Pork and other porcines	4487	3594	80.10%	0.9	1.2	4.1
Children (2-16 yrs)	Australia	2007 ANCNPAS	Sheep and other ovines	4487	2479	55.25%	0.6	1.1	4.0
Children (2-6 yrs)	Australia	2007 ANCNPAS	Beef and other bovines meat	1463	1226	83.80%	2.3	2.1	8.4
Children (2-6 yrs)	Australia	2007 ANCNPAS	Pork and other porcines	1463	1114	76.14%	1.3	1.4	5.1
Children (2-6 yrs)	Australia	2007 ANCNPAS	Sheep and other ovines	1463	741	50.65%	0.7	1.2	4.4
Infants	Bulgaria	NUTRICHILD	Beef and other bovines meat	860	89	10.35%	2.7	1.6	7.8
Infants	Bulgaria	NUTRICHILD	Pork and other porcines	860	9	1.05%	2.3	2.2	7.3
Infants	Bulgaria	NUTRICHILD	Sheep and other ovines	860	2	0.23%	2.4	1.0	3.1
Infants	Italy	INRAN_SCAI_2005_06	Beef and other bovines meat	16	1	6.25%	3.8		3.8
Infants	Italy	INRAN_SCAI_2005_06	Pork and other porcines	16	1	6.25%	1.0		1.0
Infants	Italy	INRAN_SCAI_2005_06	Sheep and other ovines	16	1	6.25%	1.3		1.3
Other children	Belgium	Regional_Flanders	Meat from mammals other than marine mammals, nes	625	16	2.56%	1.2	0.8	3.3

**Table 1.15 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Other children	Belgium	Regional_Flanders	Sheep and other ovines	625	10	1.60%	1.4	0.6	2.3
Other children	Belgium	Regional_Flanders	Beef and other bovines meat	625	185	29.60%	1.4	0.8	3.1
Other children	Belgium	Regional_Flanders	Horse and other equines	625	8	1.28%	1.3	0.6	2.2
Other children	Belgium	Regional_Flanders	Pork and other porcines	625	121	19.36%	1.2	0.7	3.1
Other children	Bulgaria	NUTRICHILD	Beef and other bovines meat	433	276	63.74%	2.6	1.8	7.2
Other children	Bulgaria	NUTRICHILD	Pork and other porcines	433	37	8.55%	1.9	1.2	6.2
Other children	Bulgaria	NUTRICHILD	Sheep and other ovines	433	8	1.85%	1.8	1.2	3.2
Other children	Czech Republic	SISP04	Beef and other bovines meat	389	125	32.13%	1.7	1.2	4.5
Other children	Czech Republic	SISP04	Pork and other porcines	389	121	31.11%	2.0	1.3	5.8
Other children	Denmark	Danish_Dietary_Survey	Beef and other bovines meat	490	482	98.37%	0.9	0.6	2.3
Other children	Denmark	Danish_Dietary_Survey	Pork and other porcines	490	480	97.96%	1.1	0.8	3.0
Other children	Denmark	Danish_Dietary_Survey	Sheep and other ovines	490	25	5.10%	0.6	0.3	1.3
Other children	Finland	DIPP	Beef and other bovines meat	933	634	67.95%	1.4	1.1	4.6
Other children	Finland	STRIP	Beef and other bovines meat	250	81	32.40%	0.8	0.6	2.1
Other children	Finland	DIPP	Pork and other porcines	933	373	39.98%	1.0	1.2	3.8
Other children	Finland	STRIP	Pork and other porcines	250	64	25.60%	0.8	0.6	2.4
Other children	Finland	DIPP	Sheep and other ovines	933	23	2.47%	0.6	0.4	1.5

**Table 1.15 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Other children	Finland	STRIP	Sheep and other ovines	250	4	1.60%	1.0	1.1	2.6
Other children	France	INCA2	Beef and other bovines meat	482	440	91.29%	1.5	1.0	3.9
Other children	France	INCA2	Horse and other equines	482	9	1.87%	0.7	0.2	1.1
Other children	France	INCA2	Meat from mammals other than marine mammals, NES	482	175	36.31%	0.2	0.4	1.7
Other children	France	INCA2	Pork and other porcines	482	227	47.10%	0.7	0.5	1.8
Other children	France	INCA2	Sheep and other ovines	482	130	26.97%	0.4	0.3	1.4
Other children	Greece	Regional_Crete	Beef and other bovines meat	839	24	2.86%	1.3	0.8	3.1
Other children	Greece	Regional_Crete	Goat and other caprines	839	23	2.74%	1.5	0.8	3.5
Other children	Greece	Regional_Crete	Meat from mammals other than marine mammals, NES	839	54	6.44%	1.5	0.9	3.8
Other children	Greece	Regional_Crete	Pork and other porcines	839	288	34.33%	1.7	0.9	3.9
Other children	Greece	Regional_Crete	Sheep and other ovines	839	149	17.76%	1.3	0.7	3.7
Other children	Italy	INRAN_SCAI_2005_06	Beef and other bovines meat	193	151	78.24%	2.0	1.4	6.0
Other children	Italy	INRAN_SCAI_2005_06	Horse and other equines	193	1	0.52%	1.7		1.7
Other children	Italy	INRAN_SCAI_2005_06	Pork and other porcines	193	71	36.79%	1.2	0.9	3.2
Other children	Italy	INRAN_SCAI_2005_06	Sheep and other ovines	193	4	2.07%	1.0	0.7	1.9
Other children	Latvia	EFSA_TEST	Beef and other bovines meat	189	6	3.17%	1.2	0.3	1.8

**Table 1.15 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Other children	Latvia	EFSA_TEST	Meat from mammals other than marine mammals, NES	189	2	1.06%	1.4	0.7	1.9
Other children	Latvia	EFSA_TEST	Pork and other porcines	189	105	55.56%	1.6	1.1	4.5
Other children	Netherlands	VCP_kids	Beef and other bovines meat	957	255	26.65%	1.2	1.0	3.4
Other children	Netherlands	VCP_kids	Horse and other equines	957	2	0.21%	0.3	0.1	0.4
Other children	Netherlands	VCP_kids	Pork and other porcines	957	167	17.45%	1.1	0.8	3.0
Other children	Netherlands	VCP_kids	Sheep and other ovines	957	10	1.04%	0.6	0.3	1.3
Other children	Spain	NUT_INK05	Beef and other bovines meat	399	155	38.85%	2.3	1.4	6.1
Other children	Spain	enKid	Horse and other equines	156	1	0.64%	3.9		3.9
Other children	Spain	NUT_INK05	Horse and other equines	399	2	0.50%	3.1	1.3	4.1
Other children	Spain	enKid	Meat from mammals other than marine mammals, NES	156	44	28.21%	2.4	1.4	6.4
Other children	Spain	enKid	Pork and other porcines	156	32	20.51%	1.8	1.0	4.6
Other children	Spain	NUT_INK05	Pork and other porcines	399	124	31.08%	1.5	0.9	3.7
Other children	Spain	enKid	Sheep and other ovines	156	5	3.21%	2.7	1.0	3.8
Other children	Spain	NUT_INK05	Sheep and other ovines	399	12	3.01%	1.9	1.2	4.9
Other children	Sweden	NFA	Beef and other bovines meat	1473	826	56.08%	0.6	0.5	1.9
Other children	Sweden	NFA	Horse and other equines	1473	15	1.02%	0.2	0.2	0.7

**Table 1.15 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Other children	Sweden	NFA	Pork and other porcines	1473	536	36.39%	0.9	0.7	2.7
Other children	Sweden	NFA	Sheep and other ovines	1473	15	1.02%	0.6	0.4	1.7
Toddlers	Belgium	Regional_Flanders	Beef and other bovines meat	36	12	33.33%	1.4	0.8	3.0
Toddlers	Belgium	Regional_Flanders	Meat from mammals other than marine mammals, NES	36	2	5.56%	1.8	0.7	2.3
Toddlers	Belgium	Regional_Flanders	Horse and other equines	36	1	2.78%	5.0		5.0
Toddlers	Belgium	Regional_Flanders	Pork and other porcines	36	11	30.56%	2.1	2.0	7.3
Toddlers	Bulgaria	NUTRICHILD	Beef and other bovines meat	428	229	53.50%	2.8	2.0	7.5
Toddlers	Bulgaria	NUTRICHILD	Pork and other porcines	428	26	6.07%	1.6	1.2	5.5
Toddlers	Bulgaria	NUTRICHILD	Sheep and other ovines	428	11	2.57%	1.6	1.0	4.0
Toddlers	Finland	DIPP	Beef and other bovines meat	497	406	81.69%	2.0	1.6	6.4
Toddlers	Finland	DIPP	Pork and other porcines	497	326	65.59%	1.4	1.4	4.8
Toddlers	Finland	DIPP	Sheep and other ovines	497	26	5.23%	1.0	0.8	4.5
Toddlers	Italy	INRAN_SCAI_2005_06	Beef and other bovines meat	36	20	55.56%	2.4	1.6	6.3
Toddlers	Italy	INRAN_SCAI_2005_06	Pork and other porcines	36	7	19.44%	0.6	0.2	1.1
Toddlers	Italy	INRAN_SCAI_2005_06	Sheep and other ovines	36	2	5.56%	1.6	0.7	2.1
Toddlers	Netherlands	VCP_kids	Beef and other bovines meat	322	84	26.09%	1.4	1.2	4.8
Toddlers	Netherlands	VCP_kids	Pork and other porcines	322	47	14.60%	1.2	1.1	4.0
Toddlers	Netherlands	VCP_kids	Sheep and other ovines	322	1	0.31%	1.0		1.0
Toddlers	Spain	enKid	Meat from mammals other than marine mammals, NES	17	3	17.65%	3.6	0.5	4.1
Toddlers	Spain	enKid	Sheep and other ovines	17	2	11.76%	1.1	0.5	1.4

**Table 1.15 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Children (1–3 yrs)	USA	FCID-WWEIA data for years 2005–2010	Total red meat	5338	2451	45.92%	2.4		7.04
Children (3–16 yrs)	USA	FCID-WWEIA data for years 2005–2010	Total red meat	12 521	9605	76.71%	1.7		4.98

NES, not elsewhere specified

Data on USA from [FCID \(2015\)](#): What We Eat In America - Food Commodity Intake Database 2005-10, U.S. Environmental Protection Agency - Office of Pesticide Programs © University of Maryland 2012 – 2016. Available from: <http://fcid.foodrisk.org/percentiles.php>

Data for other countries from [FAO/WHO \(2015\)](#): the FAO/WHO Chronic individual food consumption database – Summary statistics (CIFOcOs), © Copyright World Health Organization (WHO), 2016. All Rights Reserved. Available from: <http://www.who.int/foodsafety/databases/en/>



consumption for consumers were up to 90 and 150 g/day, respectively ([FAO/WHO, 2015; Table 1.14, Table 1.15](#)).

(d) *Offal consumption*

The per capita consumption of mammalian offal worldwide was generally lower than 10 g/day per person, except for Australia and European countries, where the highest levels (15 g/day per person) were reported by GEMS clusters diets ([WHO, 2013](#)). From National food consumption surveys, high mean consumption for consumers only was reported for a limited proportion of the population. For example, in Brazil, the average consumption of mammalian offal in the general population was 84 g/day per person for 3.5% of consumers ([FAO/WHO, 2015](#)). In Germany, the mean consumption of cattle offal for adults was 53 g/day per person for 0.3% of consumers. In China, the consumption of mammalian offal by the general population was 44 g/day per person for 3.5% of consumers. It should be noted that high consumers can eat up to about 260 g/day per person of mammalian offal (Brazil), and in such situations, offal was a likely substitute for other meat products ([FAO/WHO, 2015](#)).

(e) *Processed meat consumption*

The consumption of processed meat is more difficult to estimate than that of red meat, as it is a heterogeneous food group with different definitions across countries. Detailed worldwide data on processed meat consumption (g/kg bw per day) are presented in [Table 1.16](#) and [Table 1.17](#) for adults and children, respectively.

According to the per capita data collected by FAOSTAT, the total processed meat consumption was between 0 and 33 g/day ([FAO/WHO, 2015](#)). Based on the GEMS cluster diets, the total processed meat consumption ranged from less than 1 to 18 g/day ([WHO, 2013](#)).

In the NutriCoDE study, the median of mean consumption of processed meat ranged from 3.9 g/day (1.8–5.1 g/day) for the first quintile

to 34 g/day (26–76 g/day) for the fifth quintile ([Imamura et al., 2015](#)).

These levels of consumption of processed meat were consistent with those in Japan, where the percentage of consumers was about 97%, the mean consumption was 14 g/day, and the consumption at the 95th percentile was 34 g/day ([FAO/WHO, 2015; Table 1.16](#)). On the contrary, in China, the percentage of consumers of processed meat was about 2–3.8% of the total population; however, for this group, the mean consumption and the consumption at the 95th percentile were 66 and 182 g/day, respectively ([FAO/WHO, 2015; Table 1.16](#)). Based on three consecutive 24-hour recalls, a prospective study of 5000 adults from 4280 households in nine provinces showed that the average processed meat consumption increased by three-fold (5 vs 15 g/day per person) from 1989 to 2004 ([Zhai et al., 2009](#)).

Intake of processed meat in countries of the Middle East and north Africa was estimated in 2010 to range from 2.5 g/day (Palestine) to 6.7 g/day (United Arab Emirates) ([Afshin et al., 2015](#)).

In New Zealand, the mean consumption of sausages and processed meat was 110 g/day for women and 142 g/day for men. At the 90th percentile, the consumption reached 212 g/day for women and 300 g/day for men. In addition, the percentage of consumers older than 15 years was about 16% of the population ([Parnell et al., 2012](#)).

In Brazil, the percentage of consumers of processed meat was about 27% of the total population; however, for this group, the mean consumption and the consumption at the 95th percentile were 33 and 94 g/day, respectively ([FAO/WHO, 2015; Table 1.16](#)).

In the USA, detailed results were available for processed meat from game, beef, goat, and pork. Interestingly, the percentage of consumers ranged from 0.07% (processed goat meat) to 65% (processed beef meat), but the mean consumption ranged from 42 to 99 g/day, and the consumption

**Table 1.16 Worldwide consumption of processed meat in adults**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
General population	Brazil	Brazilian Institute of Geography and Statistics	Processed meat and meat products, NES	34 003	334	0.98%	0.6	0.6	2.1
General population	Brazil	Brazilian Institute of Geography and Statistics	Processed meat and meat products, NES	34 003	9047	26.61%	0.5	0.5	1.8
General population	Brazil	Brazilian Institute of Geography and Statistics	Processed meat and meat products, NES	34 003	54	0.16%	1.2	0.8	2.6
General population	Brazil	Brazilian Institute of Geography and Statistics	Processed meat and meat products, NES	34 003	14	0.04%	1.4	0.8	3.6
General population	Brazil	Brazilian Institute of Geography and Statistics	Processed meat and meat products, NES	34 003	8	0.02%	0.5	0.2	0.8
General population	Brazil	Brazilian Institute of Geography and Statistics	Processed meat and meat products, NES	34 003	24	0.07%	0.9	0.7	2.1
General population	China	2002 China Nutrition and Health Survey	Processed meat and meat products, NES	65 359	1430	2.19%	1.2	1.1	4.2
General population	China	2002 China Nutrition and Health Survey	Processed meat and meat products, NES	65 359	2483	3.80%	0.9	1.3	2.7
General population	Japan	DSFFQ_FI	Processed meat and meat products, NES	2711	2642	97.45%	0.3	0.2	0.8
General population	Japan	DSFFQ_FI	Processed meat and meat products, NES	2711	24	0.89%	0.0	0.0	
Adults	Belgium	Diet_National_2004	Processed meat and meat products, NES	1304	956	73.31%	0.8	0.7	2.7
Adults	Czech Republic	SISP04	Processed meat and meat products, NES	1666	1427	85.65%	1.2	1.0	3.9
Adults	Denmark	Danish Dietary Survey	Processed meat and meat products, NES	2822	2800	99.22%	0.4	0.3	1.3
Adults	Finland	FINDIET_2007	Processed meat and meat products, NES	1575	1188	75.43%	0.7	0.7	2.9

**Table 1.16 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Adults	France	INCA2	Processed meat and meat products, NES	2276	2167	95.21%	0.6	0.4	1.6
Adults	Hungary	National_Repr_Surv	Processed meat and meat products, NES	1074	1003	93.39%	1.1	0.8	3.0
Adults	Ireland	NSIFCS	Processed meat and meat products, NES	958	906	94.57%	0.8	0.6	2.1
Adults	Italy	INRAN_SCAI_2005_06	Processed meat and meat products, NES	2313	1921	83.05%	0.5	0.4	1.6
Adults	Latvia	EFSA_TEST	Processed meat and meat products, NES	1306	868	66.46%	0.9	0.7	2.8
Adults	Netherlands	DNFCS_2003	Processed meat and meat products, NES	750	618	82.40%	0.7	0.6	2.4
Adults	Spain	AESAN	Processed meat and meat products, NES	410	334	81.46%	0.9	0.7	2.5
Adults	Spain	AESAN_FIAB	Processed meat and meat products, NES	981	908	92.56%	0.8	0.6	2.5
Adults	Sweden	Riksmaten_1997_98	Processed meat and meat products, NES	1210	1147	94.79%	0.7	0.4	1.6
Adults	United Kingdom	NDNS	Processed meat and meat products, NES	1724	1492	86.54%	0.5	0.4	1.4

NES, not elsewhere specified

Data on USA from [FCID \(2015\)](#): What We Eat In America – Food Commodity Intake Database 2005–10, US. Environmental Protection Agency – Office of Pesticide Programs © University of Maryland 2012 – 2016. Available from: <http://fcid.foodrisk.org/percentiles.php>

Data for other countries from [FAO/WHO \(2015\)](#): the FAO/WHO Chronic individual food consumption database – Summary statistics (CIFOcOs), © Copyright World Health Organization (WHO), 2016. All Rights Reserved. Available from: <http://www.who.int/foodsafety/databases/en/>

**Table 1.17 Worldwide consumption of processed meat in children**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, Mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Adolescents	Belgium	Diet_National_2004	Processed meat and meat products, NES	584	413	70.72%	0.8	0.7	2.7
Adolescents	Cyprus	Child health	Processed meat and meat products, NES	303	183	60.40%	0.4	0.3	1.1
Adolescents	Czech Republic	SISP04	Processed meat and meat products, NES	298	274	91.95%	1.4	1.2	4.6
Adolescents	Denmark	Danish Dietary Survey	Processed meat and meat products, NES	479	477	99.58%	0.6	0.5	2.0
Adolescents	France	INCA2	Processed meat and meat products, NES	973	950	97.64%	0.7	0.5	2.2
Adolescents	Italy	INRAN_SCAI_2005_06	Processed meat and meat products, NES	247	216	87.45%	0.8	0.6	2.2
Adolescents	Latvia	EFSA_TEST	Processed meat and meat products, NES	470	333	70.85%	1.2	1.0	3.7
Adolescents	Spain	enKid	Processed meat and meat products, NES	209	190	90.91%	1.6	1.3	5.3
Adolescents	Spain	AESAN_FIAB	Processed meat and meat products, NES	86	81	94.19%	1.0	0.7	2.7
Adolescents	Spain	NUT_INK05	Processed meat and meat products, NES	651	574	88.17%	1.1	0.9	3.4
Adolescents	Sweden	NFA	Processed meat and meat products, NES	1018	918	90.18%	1.0	0.8	2.5
Children	China	2002 China Nutrition and Health Survey	Processed meat and meat products, NES	2784	78	2.80%	2.2	1.5	6.8
Children	China	2002 China Nutrition and Health Survey	Processed meat and meat products, NES	2784	78	2.80%	2.2	6.3	8.9
Children	Japan	DSFFQ_FI	Processed meat and meat products, NES	71	71	100.00%	0.8	0.6	2.6
Infants	Bulgaria	NUTRICHILD	Processed meat and meat products, NES	860	33	3.84%	2.1	1.3	6.3
Infants	Italy	INRAN_SCAI_2005_06	Processed meat and meat products, NES	16	1	6.25%	1.5		1.5

**Table 1.17 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, Mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Other children	Belgium	Regional Flanders	Processed meat and meat products, NES	625	468	74.88%	1.5	1.3	4.7
Other children	Bulgaria	NUTRICHILD	Processed meat and meat products, NES	433	261	60.28%	1.9	1.6	5.7
Other children	Czech Republic	SISP04	Processed meat and meat products, NES	389	314	80.72%	1.6	1.4	5.5
Other children	Denmark	Danish Dietary Survey	Processed meat and meat products, NES	490	488	99.59%	1.3	0.9	4.0
Other children	Finland	STRIP	Processed meat and meat products, NES	250	218	87.20%	1.3	1.0	3.9
Other children	Finland	DIPP	Processed meat and meat products, NES	933	825	88.42%	1.8	1.6	6.1
Other children	France	INCA2	Processed meat and meat products, NES	482	465	96.47%	1.3	0.9	3.5
Other children	Greece	Regional Crete	Processed meat and meat products, NES	839	327	38.97%	0.5	0.5	1.8
Other children	Italy	INRAN_SCAI_2005_06	Processed meat and meat products, NES	193	157	81.35%	1.2	1.0	4.0
Other children	Latvia	EFSA_TEST	Processed meat and meat products, NES	189	113	59.79%	1.8	1.6	6.4
Other children	Netherlands	VCP kids	Processed meat and meat products, NES	957	748	78.16%	1.6	1.2	4.5
Other children	Spain	enKid	Processed meat and meat products, NES	156	138	88.46%	2.2	1.5	6.5
Other children	Spain	NUT_INK05	Processed meat and meat products, NES	399	357	89.47%	1.7	1.2	4.5
Other children	Sweden	NFA	Processed meat and meat products, NES	1473	1379	93.62%	1.5	1.1	4.3
Toddlers	Belgium	Regional Flanders	Processed meat and meat products, NES	36	24	66.67%	1.9	1.2	5.7
Toddlers	Bulgaria	NUTRICHILD	Processed meat and meat products, NES	428	164	38.32%	2.0	1.5	5.2

**Table 1.17 (continued)**

Age class	Country	Survey	Meat type	No. of subjects	No. of consumers	Percentage of consumers	Consumers, Mean consumption (g/bw per day)	Consumers, STD (g/bw per day)	Consumers, P975 (g/bw per day)
Toddlers	Finland	DIPP	Processed meat and meat products, NES	497	142	28.57%	1.5	1.6	5.5
Toddlers	Italy	INRAN_SCAI_2005_06	Processed meat and meat products, NES	36	22	61.11%	1.6	1.2	5.6
Toddlers	Netherlands	VCP kids	Processed meat and meat products, NES	322	254	78.88%	1.8	1.6	6.5
Toddlers	Spain	enKid	Processed meat and meat products, NES	17	13	76.47%	2.7	1.8	6.9

NES, not elsewhere specified

Data on USA from [FCID \(2015\)](#): What We Eat In America – Food Commodity Intake Database 2005–10, United States Environmental Protection Agency – Office of Pesticide Programs

© University of Maryland 2012 – 2016. Available from: <http://fcid.foodrisk.org/percentiles.php>

Data for other countries from [FAO/WHO \(2015\)](#): the FAO/WHO Chronic individual food consumption database – Summary statistics (CIFOCoss), © Copyright World Health Organization (WHO), 2016. All Rights Reserved. Available from: <http://www.who.int/foodsafety/databases/en/>

at the 95th percentile ranged from 152 to 309 g/day ([FCID, 2015](#)).

In Europe, the mean consumption of processed meat for adults was between about 10 and 80 g/day. The consumption at the 95th percentile was up to 200 g/day ([EFSA, 2011](#)). In the EPIC cohort, the lowest consumption of processed meat was found in Greece, with 11 g/day for women and 19 g/day for men. The highest consumption of processed meat was found in Norway for women (48 g/day) and in Germany for men (89 g/day) ([Linseisen et al., 2006](#)).

[The Working Group noted that despite the weaknesses of the data set, it seemed that in certain countries the consumption of processed meat is similar to the consumption of red meat for consumers only. However, the percentage of consumers of processed meat seemed to be much smaller, leading to a per capita consumption four to five times lower than that of red meat.]

(f) *Dietary exposure to chemicals in meat*

(i) *Chemicals in the environment*

Several chemicals classified as carcinogens by the International Agency for Research on Cancer (IARC) are present in the environment and can contaminate meat through air, water, or animal feed. They can be generated either from industrial activities or from microorganisms ([IARC, 2010a, b, 2012a, b, 2016](#)).

*Dioxin and dioxin-like compounds:* The Joint FAO/WHO Expert Committee on Food Additives (JECFA) assessed dioxins and related compounds in 2002. The dietary exposure estimate, expressed as toxic equivalency factors for PCDDs and PCDFs based on national data, ranged from 33 to 42 pg/kg bw per month and from 81 to 100 pg/kg bw per month at the 50th and 90th percentiles, respectively. For coplanar PCBs, the dietary exposure estimate ranged from 9 to 47 pg/kg bw per month and from 25 to

130 pg/kg bw per month at the 50th and 90th percentiles, respectively. The contribution from meat was estimated to range from 6% in Asia to 23% in north America for PCDDs and PCDFs, and from 4% in Asia to 55% in north America for dioxin-like PCBs ([JECFA, 2002](#)).

*Brominated flame retardants (BFRs):* Food consumption, especially fish and meat product consumption, is a major route of human contamination ([Lyche et al., 2015](#)). For example, higher levels of PBDEs in humans were found in studies in the USA where fish were most highly contaminated (median, 616 pg/g), followed by meat (median, 190 pg/g). However, unlike many European countries where fish consumption predominates, dietary intake of PBDEs in the USA is mostly from meat consumption ([Schechter et al., 2008](#)).

*Heavy metals:* The heavy metals cadmium, arsenic, and lead have been classified as carcinogens by IARC ([IARC, 2012a](#)). For the EU, the European Food Safety Authority (EFSA) has estimated that average weekly dietary exposure to cadmium was 2.04 µg/kg bw, and at the 95th percentile, weekly dietary exposure to cadmium was 3.66 µg/kg bw. Food consumed in larger quantities had the greatest impact on dietary exposure to cadmium. This was true for the broad food categories of grains and grain products (26.9%). Meat and edible offal were estimated to contribute 7.7% of the total dietary exposure ([EFSA, 2012](#)). In 2010, JECFA estimated that for adults, the mean dietary exposure to cadmium was 2.2–12 µg/kg bw per month, and high-level dietary exposure to cadmium was 6.9–12.1 µg/kg bw per month. For children aged 6 months to 12 years, the mean dietary exposure to cadmium was 3.9–20.6 µg/kg bw per month. Meat was not part of the food groups that contributed significantly (40–85%) to the total dietary

exposure to cadmium (i.e. rice, wheat, vegetables, and molluscs) ([JECFA, 2013](#)).

Dietary exposure to inorganic arsenic was last evaluated by JECFA in 2011. The occurrence of total arsenic in meat ranged from 0.004 to 0.78 mg/kg, and meat was not a major contributor to dietary exposure to inorganic arsenic ([JECFA, 2011](#)).

Lead was last evaluated by JECFA in 2011. Mean dietary exposure to lead ranged from 0.02 to 3 µg/kg bw per day for adults, and from 0.03 to 9 µg/kg bw per day for children. The contribution of meat and meat products, including offal, was estimated to be 9% of the total dietary exposure to lead ([JECFA, 2013](#)).

*Mycotoxins*: EFSA concluded that carry-over of aflatoxin, deoxynivalenol, zearalenone, and fumonisin to products of animal origin was very low ([EFSA, 2004a, b, c, 2005c](#); [Kan & Meijer, 2007](#)). Accumulation of ochratoxin A occurred predominantly in the blood, liver, and kidney. Muscle, milk, and eggs contained much lower levels of this mycotoxin ([EFSA, 2004d](#)).

## (ii) Chemicals from cooking practices

*Heterocyclic aromatic amines (HAAs)*: No international dietary exposure assessment was available for HAAs; however, in the EPIC study, dietary exposure to HAAs was estimated in the Heidelberg cohort (Germany) using a detailed dietary questionnaire that assessed meat consumption, cooking methods, and degree of browning of the respective food items. Results based on total meat consumption (including poultry meat) showed a total median exposure to HAAs of 30.6 ng/day (13–71.3 ng/day) ([Rohrmann et al., 2007](#)). Other studies' results showed a significantly lower dietary exposure to HAAs for Europe (6.1 ng/kg bw per day) ([Zimmerli et al., 2001](#)) and the USA (11.0–19.9 ng/kg bw per day) ([Keating & Bogen, 2004](#)).

*Polycyclic aromatic hydrocarbons (PAHs)*: In 2006, JECFA estimated the dietary exposure to PAHs in 18 countries, including Australia, Brazil, New Zealand, and the United Kingdom. Estimated intake of BaP ranged from < 1 to 2.0 µg/day and from 0.0001 to 0.005 µg/kg bw per day. For the other nine PAHs, intake ranged from less than 1 to ~12 µg/day and from 0.0001 to 0.015 µg/kg bw per day ([WHO, 2006](#)). Generally, despite high concentrations of PAHs, meat and barbecued foods were not major contributors to PAH exposure; however, in the USA, grilled and barbecued meat was estimated to contribute to 21% of the intake of BaP ([WHO, 2006](#)). Cereals, vegetal oil, animal fat, and vegetal fat contributed up to 60% to the whole food intake of PAHs, as they are major contributors by weight to the total diet ([Dennis et al., 1983](#)).

*Nitrosamines*: The main sources of NOCs in the diet are nitrite-preserved meat products ([Tricker, 1997](#); [Haorah et al., 2001](#)). [Haorah et al. \(2001\)](#) reported a mean concentration of 5.5 µmol/kg of NOCs in frankfurters, but only 0.5 µmol/kg of NOCs in fresh meat.

*Acrylamide*: Acrylamide may occur in meat during cooking ([Tareke et al., 2002](#)). However, meat has been estimated to be a minor contributor, between 0.2% and 2% of total dietary intake ([WHO, 2006](#)).

## 1.4 Exposure assessment and biological markers

### 1.4.1 Questionnaires

A description of the epidemiological studies included in this *Monograph*, in terms of their study design, is provided in Section 2. A review of dietary assessment methodologies used in the epidemiological studies is beyond the scope of this *Monograph* (e.g. [Thompson & Subar, 2013](#)).



The majority of the studies used food frequency questionnaires (FFQs) to assess individual meat intake (including red meat and processed meat). FFQs are typically used in epidemiological studies to measure usual dietary intake in individuals for several reasons. First, FFQs are a feasible approach in case–control studies, where usual diet must be ascertained retrospectively (often from the distant past). Second, in large prospective cohort studies, FFQs can be distributed by mail or online to a large number of participants; are self-administered (typically); may be optically scanned, computer-assisted, or web-based; and are analysed using precoded foods/food groups and portion sizes.

The FFQ approach asks respondents to report their usual frequency of consumption for each food from a list of foods during a specific period of time (several months or a year). FFQs are generally used for ranking subjects according to food or nutrient intake, rather than for estimating absolute levels of intake. In addition, they are widely used in case–control and cohort studies to assess an association between dietary intake and disease risk ([Kushi, 1994](#); [Beaton, 1994](#); [Sempos et al., 1999](#)).

The ability to quantify total dietary intake depends on the number of food items listed in the FFQ, on the level of detail collected within the questionnaire, on whether portion sizes for the foods/food groups are included, and on the timeframe of intake or reference period used. For red meat and processed meat specifically, the classifications used to define red meat and processed meat as a food category also influence the calculation of total dietary intake ([Block et al., 1986](#); [Rimm et al., 1992](#)).

Although food lists included in FFQs vary based on the purpose of the study and the study population, the appropriateness of the food lists is crucial. The full variability of an individual's diet, which includes many foods and mixed dishes, cannot be captured by a finite food list. [Ollberding et al. \(2012\)](#), for example, identified

a food list for their FFQ using 3-day measured food records that could capture 85% or more of the intake of key nutrients and also food items traditionally consumed by the populations represented in the Multiethnic Cohort Study.

Many FFQs have been developed and adapted to suit different research questions and populations. In the USA, for example, several questionnaires are commonly used (and are cited in this *Monograph*), including:

*Health Habits and History Questionnaire (HHHQ) or Block questionnaire* ([Block et al., 1986, 1990](#); [Sobell et al., 1989](#)): This is a semiquantitative food frequency questionnaire (SQFFQ) originally developed by the National Cancer Institute (NCI). The SQFFQ collects portion size information; however, portion sizes are specified as standardized portions or by choosing from a range of portions sizes (e.g. small, medium, or large). The original Block FFQ has been modified, and is continually updated by researchers to suit their research questions and populations.

*Harvard FFQ or Willett questionnaire* ([Caan et al., 1998](#); [McCann et al., 1999](#)): This FFQ was developed at Harvard University. Standard portion size defaults are included as part of the food items listed, rather than as a separate listing.

*NCI Diet History Questionnaire (DHQ)*: The DHQ was designed with an emphasis on cognitive ease of use for respondents ([Subar et al., 1995, 2001](#)). It is an SQFFQ, which uses an embedded question approach, that was developed by NCI.

Definitions of red meat and processed meat as a food category varied across the studies included in this *Monograph*. Red meat was commonly defined as beef, pork, lamb, or a combination thereof, and processed meat was generally defined as meat made largely from pork, beef, or poultry that undergoes methods of preservation,

such as curing, smoking, or drying ([Santarelli et al., 2008](#)). While many studies explicitly defined these classifications ([Tiemersma et al., 2002](#); [Ferrucci et al., 2009](#); [Cross et al., 2010](#)), other studies provided either no description or an unclear description of these classifications ([Kato et al., 1997](#); [Järvinen et al., 2001](#)). The level of detail collected by the epidemiological studies, in terms of meat intake, varied widely. Most studies reported the association between categories of meat intake labelled “red meat” or “processed meat” and cancer risk; however, several studies reported results for individual red meat items (e.g. beef or pork) ([Brink et al., 2005](#); [Norat et al., 2005](#); [Sato et al., 2006](#); [Takachi et al., 2011](#); [Egeberg et al., 2013](#)) and/or processed meat items (e.g., hot dogs or bacon). The studies that included detailed information on the intake of specific processed meat items were superior to those that combined generic items into one food group or one-line items (e.g. “processed meat”), as the amount of nitrate, nitrite, and haem iron in processed foods can vary dramatically.

#### (a) Portion size

Scientists have used many methods to improve assessment of portion size in the studies included in this report. For example, [Pietinen et al. \(1999\)](#) included a portion size booklet of 122 photographs of foods, each with three to five different portion sizes. In the Canadian National Breast Screening Study (CNBSS), [Kabat et al. \(2007\)](#) also included photographs of portion sizes to improve portion size assessment. In the Finnish Mobile Clinic Health Examination Survey, [Järvinen et al. \(2001\)](#) used plastic food models and real foods to help estimate portion sizes for their interviewer-assisted FFQ. [Dixon et al. \(2004\)](#) also used three-dimensional food models, plastic cups, and spoons to help participants identify usual serving sizes in the Kaiser Permanente Medical Care Program in northern California, USA.

#### (b) Validation and calibration

The relative validity of an FFQ provides information on how well the instrument is measuring what it is intended to measure. This is completed by comparing intake assessed using an FFQ with intake assessed using a reference method (which is deemed to be superior) in the same individuals (e.g. an interviewer-led dietary history or multiple 24-hour recalls). FFQs may often be validated for their ability to assess total energy intake in comparison with the doubly labelled water technique ([Hill & Davies, 2001](#)). The superior method is often prohibitive for use in large epidemiological studies due to participant burden, or overall cost of administering and coding the instrument. Calibration studies are used to calibrate an FFQ to a reference method using a regression model. Many of the studies included in this *Monograph* used various statistical methods, employing measurement error models and energy adjustment to assess the validity of the FFQs and to adjust estimates of the relative risks for disease outcomes ([Bingham & Day, 1997](#); [Kipnis et al., 1997](#); [Carroll et al., 1998](#); [Hu et al., 1999](#)). For example, in the National Institutes of Health – American Association of Retired Persons (NIH-AARP) Diet and Health Study, the FFQ used was calibrated against two non-consecutive 24-hour dietary recalls ([Cross et al., 2010](#)). In the EPIC study, investigators used a computerized 24-hour dietary recall method to calibrate dietary measurements across countries and to correct for systematic over- or underestimation of dietary intake ([Norat et al., 2005](#); [Pala et al., 2009](#)). [Tiemersma et al. \(2002\)](#) validated their short SQFFQ using a dietary history method, which is a robust, interviewer-administered dietary assessment method. Each study included in this *Monograph* was examined to determine whether the FFQ used to assess red meat and processed meat exposure had been validated (see Section 2).

(i) *Heterocyclic aromatic amines*

Assessment of dietary HAA intake is challenging, as HAA concentrations vary greatly according to cooking technique, temperature, cooking time, and meat type ([Sinha et al., 1995](#); [Knize et al., 1998](#); [Sinha et al., 1998a, b](#)). Epidemiological studies have tried to overcome these difficulties using surrogate markers of HAA intake, such as method of cooking, surface browning, total cooking time, and gravy intake.

To estimate the intake of these cooked meat mutagens, a detailed meat-cooking module was developed. The meat-cooking module was a modified version of the 1992, 100-item, self-administered HHHQ to assess usual dietary intake over the past year ([Block et al., 1986](#)). An interviewer-administered questionnaire on meat-cooking practices was also used to assess the consumption of 23 meat, poultry, and fish items using a matrix similar to the 100-item HHHQ. The questionnaire collected information on cooking methods; embedded questions assessed how well the meat was cooked. Portion size was estimated as small, medium, or large, relative to the standard portion size indicated for each food listed in the questionnaire and meat-cooking module. A mutagen database, called Computerized Heterocyclic Amines Resource for Research in Epidemiology of Disease (CHARRED), developed by NCI/NIH ([NCI, 2017](#)) was used to estimate the intake of mutagenic compounds from cooked meats. The CHARRED database estimates the HAA content of commonly consumed meats, based on detailed information about the meat-cooking methods used and meat-doneness level. The relative validity of this meat-cooking module has been measured using multiple food diaries (three of four non-consecutive day diaries completed over a 3-month period) as the reference method ([Cantwell et al., 2004](#)). Dietary intake of the three most abundant HAAs was considered: MeIQx, 2-amino-3,4,8-trimethylimidazo[4,5-f]quinoxaline (4,8-DiMeIQx), and PhIP. Crude

correlation coefficients of HAA intake, assessed using the FFQ and food diaries, were 0.43 (95% confidence interval, CI, 0.30–0.55) for MeIQx intake and 0.22 (95% CI, 0.07–0.36) for PhIP intake. Deattenuated correlations were 0.60 (95% CI, 0.49–0.69) and 0.36 (95% CI, 0.22–0.49), respectively ([Cantwell et al., 2004](#)). This meat-cooking module has been used in the Prostate, Lung, Colorectal and Ovarian (PLCO) Cancer Screening Trial ([Cross et al., 2005](#)), Nurses' Health Study (NHS) ([Wu et al., 2010](#)), Health Professionals Follow-Up Study (HPFS), and other studies ([Cantwell et al., 2004](#)).

(ii) *Nitrate and nitrite*

Dietary assessment of nitrate and nitrite intake was reported by the NIH-AARP Diet and Health Study ([Dellavalle et al., 2013](#)). The baseline questionnaire included an FFQ that asked participants about their frequency of consumption and portion sizes of 124 food items over the past 12 months. Intake of each item was assessed using 10 predefined categories, ranging from “never” to “≥ 2 times per day” for foods, “never” to “≥ 6 times per day” for beverages, and three portion size categories. The FFQ was developed and validated by NCI using two 24-hour recalls in a subset of the cohort ([Thompson et al., 2008](#)). Concentrations of nitrate and nitrite for each food item were estimated from the existing body of scientific literature, as previously described ([Ward et al., 2003, 2006](#); [Kilfoy et al., 2011](#)). Daily intake of nitrate and nitrite was calculated by multiplying the frequency of consumption by the portion size and the nitrate and nitrite content of each food item, and then summing across all food items. Nitrate and nitrite intake from animal sources and plant sources was calculated separately. In addition to examining nitrate and nitrite intake from all animal sources, intake from processed meat sources was examined separately, as were animal sources excluding processed meat (this primarily included intake

from fresh meats, eggs, yogurt, cheese, and other dairy products).

The nitrate and nitrite content of over 3000 foods was determined by conducting a review of the literature, focusing on Canadian and USA foods, and by calculating the means of the published values weighted by the number of samples analysed ([Ward et al., 2003, 2006](#); [Kilfoy et al., 2011](#)). The nitrate and nitrite values for foods constituting an FFQ line item were combined by weighting the food-specific values by sex-specific intake amounts from the 1994–1996 Continuing Survey of Food Intakes by Individuals (CSFII) ([Subar et al., 2000](#)). For example, the nitrate content of a line item was calculated using the weighted average of the nitrate content of the included foods, where the weights were determined by intake amounts from the CSFII, based on age group and sex. Daily intake of nitrate and nitrite was calculated by multiplying the frequency of consumption of each line item by its nitrate or nitrite content and summing over line items. In addition to calculating nitrate and nitrite intake from all foods, nitrite intake from plant, animal, and processed meat sources was calculated separately.

#### (c) *Heterogeneity across studies*

There was substantial heterogeneity across the studies included in this *Monograph* due to a variety of factors, such as different methods of dietary assessment and/or measurement, definitions (e.g. food groups and serving sizes), analytical categorizations (e.g. servings/week and g/day), exposure contrasts (e.g. analytical cut-points and intake level comparisons), and degrees of adjustment for potential confounding factors. Each cohort study included in this *Monograph* is described in Section 2. The strengths and limitations of the questionnaires used in studies included in this *Monograph* are outlined below.

#### (d) *Cohort studies*

A major strength of cohort studies in nutritional epidemiology is their ability to demonstrate a temporal relationship between dietary exposure and cancer risk, as all dietary assessments are completed before diagnoses. This limits difficulties with recall bias and reverse causation.

[Wei et al. \(2004\)](#) used a validated, self-administered, 61-item SQFFQ at baseline in 87 733 women from the NHS and a validated, self-administered, 131-item SQFFQ in 46 632 men from the HPFS. The study had several strengths. For example, the FFQs used were extensively validated and tested for reproducibility using data collected from a subgroup of participants who completed two FFQs (1 year apart) and two 1-week diet records (6 months apart during the intervening year). The association between baseline meat intake and cancer risk was assessed in this study, and red meat intake was clearly defined as the consumption of beef, pork, or lamb as a main dish. In addition, in this combined cohort of women and men, risk estimates were adjusted using a multivariate model that included important confounders (age; family history; body mass index, BMI; physical activity; beef, pork, or lamb as a main dish; processed meat; alcohol; calcium; folate; height; smoking pack-years before aged 30 years; history of endoscopy; and sex). A limitation of this study was the quantification of red meat in servings per day only (i.e. not in g/day).

In the Physicians' Health Study (PHS), [Chen et al. \(1998\)](#) used a nested case-control design to assess the relationship between red meat intake and colorectal cancer by *N*-acetyltransferase (NAT) genotype. The study included 212 men who were recruited as part of the Physicians' Health Study and were subsequently diagnosed with colorectal cancer or rectal cancer during 13 years of follow-up and were genotyped via baseline blood sample, along with 221 controls. At baseline, participants completed an abbreviated, self-administered FFQ, which inquired

about usual consumption of red meat (beef, pork, or lamb as a main dish, as a mixed dish or sandwich, and as hot dogs), chicken, and fish. The abbreviated FFQ used in this study was not validated, but an expanded form of this FFQ was validated among other male health professionals. There were some limitations to this study, as the use of an abbreviated FFQ (with fewer food items listed) prevented the adjustment of risk estimates for total energy intake. In addition, dietary intake of processed meat was included with red meat intake, and meat intake was assessed as servings per day only (i.e. not in g/day).

Dietary assessment in the NIH-AARP Diet and Health Study was described in detail by [Cross et al. \(2010\)](#). The study included approximately half a million women and men, each of whom completed a validated, self-administered, 124-item FFQ at baseline. Approximately 6 months later, cancer-free participants were mailed a risk factor questionnaire, which detailed information on meat intake and cooking preferences. Meat cooking method (grilled/barbecued, pan-fried, microwaved, and broiled) and doneness level (well done/very well done and medium/rare) were used in conjunction with the CHARRED database to estimate the intake of several HAAs. The FFQ assessed the usual frequency of consumption and portion size information of foods and drinks over the past 12 months. All types of beef, pork, and lamb were considered red meat, including bacon, beef, cold cuts, ham, hamburger, hot dogs, liver, pork, sausage, and steak. Processed meat included bacon, cold cuts (red and white meat), ham, luncheon meats (red and white meat), poultry sausage, red meat sausage, and standard hot dogs and low-fat hot dogs made from poultry. Meats added to complex food mixtures, such as pizza, chilli, lasagne, and stew, contributed to the relevant meat type. There were many notable strengths to this study. Several of these strengths were related to the FFQ, which not only contained detailed questions pertaining to the components

of meat, but also was calibrated within this study population using two non-consecutive 24-hour dietary recalls. However, there was overlap in the definitions of red meat and processed meat, as some processed meat items were classified as red meat.

In the EPIC study ([Norat et al., 2005](#)), dietary intake over the 12 months before enrolment was measured by country-specific, validated dietary questionnaires (88–266 food items, depending on the country), which were self-administered in most countries; in Malmö, Sweden, a questionnaire combined with a food record was used. A second dietary measurement was taken from an 8% random sample of the cohort (36 994 participants) using a computerized 24-hour dietary recall method to calibrate dietary measurements across countries and to correct for systematic over- or underestimation of dietary intake. The major strengths of this study were the large variability in dietary intake across the population and the use of a computerized 24-hour dietary recall method to calibrate dietary measurements across countries.

In the Alpha-Tocopherol, Beta-Carotene Cancer Prevention (ATBC) Study, researchers used a self-administered, modified dietary history method to capture usual dietary intake 12 months before recruitment. The dietary history method included 276 food items and a portion size booklet of 122 photographs of foods, each with 3–5 different portion sizes. Red meat intake was defined as intake of beef, pork, or lamb ([Pietinen et al., 1999](#)). A major strength of this study was the use of a detailed questionnaire.

In the Multiethnic Cohort Study, [Ollberding et al. \(2012\)](#) assessed diet using a validated quantitative FFQ, which included a list of foods identified from 3-day measured food records, to capture 85% or more of the intake of key nutrients and food items traditionally consumed by the populations represented in the cohort. The definition of meat intake was clearly defined as total meat, red meat, and processed meat. Risk

estimates were adjusted for important potential confounders, including energy intake. This study had many strengths, including a large variability in diet due to the inclusion of multiple ethnicities and the use of an extensive dietary questionnaire.

The PLCO Cancer Screening Trial ([Ferrucci et al., 2009](#)) used an NCI DHQ to assess usual intake (both frequency and portion size) of 124 food items over the past year. The definition of red meat (g/day) included bacon, beef, cheeseburger, cold cuts, ham, hamburger, hot dogs, liver, pork, sausage, veal, venison, and red meat from mixed dishes. Processed meat included bacon, cold cuts, ham, hot dogs, and sausage. However, a limitation of this study was the clear overlap in the definitions of red meat and processed meat, as some processed meat items were classified as red meat.

[Flood et al. \(2003\)](#) used a 62-item NCI Block FFQ in the Breast Cancer Detection Demonstration Project (BCDDP) to assess red meat intake in the previous year. A limitation of this study was the combined estimate of exposure to meat, which included pork, beef, hamburger, processed meat, and liver, so risk estimates for red meat alone or processed meat alone were not possible. As the cohort was generated based on a screening programme, participants may have changed their dietary habits before baseline, and recorded intake may therefore not have accurately reflected long-term intake.

[Singh & Fraser \(1998\)](#) assessed dietary intake in the Adventists Health Study cohort using a self-administered, mailed, 55-item SQFFQ. The SQFFQ included just six questions regarding red meat intake, defined as current intake of beef or pork. A limitation of this study was the relatively short dietary questionnaire in a low-risk population, with low red meat consumption.

In the New York University Women's Health Study, [Kato et al. \(1997\)](#) assessed red meat intake using a 70-item FFQ, which was slightly modified from the questionnaire designed by Block and coworkers ([Block et al., 1986](#)). However, the

FFQ was not very extensive, and there were no quantitative data on red meat intake provided. It is also unclear whether intake of red meat included processed meat.

A study by [Tiemersma et al. \(2002\)](#) in the Netherlands examined the association between meat intake and cancer risk using a nested case-control design. A strength of this study was the use of an SQFFQ, which was validated for use through comparison with a dietary history method. A limitation of this study, however, was that a major source of meat in the population (i.e. a mixture of pork and beef) was not captured by the FFQ.

In the Shanghai Women's Health Study, [Lee et al. \(2009\)](#) assessed dietary intake at baseline using a validated quantitative FFQ, which included 19 food items/groups of animal origin. A major strength of this study was that the FFQ was administered by interview.

In the prospective cohort study of 37 112 residents of Melbourne, Australia, [English et al. \(2004\)](#) assessed dietary intake using a 124-item FFQ. They also provided a clear definition of what they included in terms of fresh red meat (veal, beef, lamb, pork, rabbit, or other game). A limitation of this study, however, was that portion size was not measured.

In the Iowa Women's Health Study, ([Lee et al., 2005](#)) assessed usual dietary intake over the past year using a validated, 127-item, self-administered SQFFQ virtually identical to the questionnaire used in the 1984 survey of the NHS ([Bostick et al., 1994](#)). Red meat was defined as beef, pork, or lamb as a main dish. This study had several strengths, including assessment of reliability and accuracy of the FFQ used, which was comparable to what was observed in the NHS. The extensive FFQ allowed for multivariable adjustment, including age, total energy intake, height, parity, total vitamin E intake, interaction term vitamin E \*age, and vitamin A supplement use.

*(e) Case-control studies*

A description of the case-control studies included in this *Monograph* is provided in Section 2. Case-control studies investigating the association between meat intake and cancer risk are limited, as they assess dietary intake after cancer has been diagnosed, which can lead to recall bias. In addition, patients often change their dietary intake due to the presence of a solid tumour to avoid pain or reflux, for example. As a result, investigators usually ask cases included in their studies to recall dietary intake in the period before diagnosis of cancer to capture usual diet before diagnosis. As a result, case-control studies are limited due to the measurement error associated with dietary intake due to memory recall. [Tavani et al. \(2000\)](#), for example, assessed total red meat intake (beef, veal, and pork) per week 2 years before diagnosis, while [Chiu et al. \(2003\)](#) assessed dietary intake 5 years before diagnosis in a case-control study in Shanghai.

In the North Carolina Colon Cancer Study, [Butler et al. \(2003\)](#) assessed usual diet in the year before diagnosis for patients, or the year before the date of selection for controls, using a 150-item FFQ, which was a modified version of the Block questionnaire. However, no information regarding validation was provided. Red meat intake was calculated as the sum of hamburger, steak, pork chop, sausage, and bacon intake.

*(f) Conclusion*

As outlined in this section, the questionnaires used in the cohort and case-control studies varied in several ways, including in the methods of dietary assessment and/or measurement, the use of validated/calibrated questionnaires, the definitions of meat and processed meat as food groups, the inclusion of serving or portion sizes, the ability to assess intake (i.e. in g/day), and the degree of adjustment for potential confounding factors.

*1.4.2 Biological markers*

Despite more than 35 years of research, no long-term validated biomarkers of exposure have been employed in molecular epidemiology studies to assess the role of genotoxins in cooked or processed meat and cancer risk. Additionally, other than HAAs, the biomarkers of PAHs or NOCs are not specific to meat, as they may also measure environmental or endogenous exposure.

[The Working Group noted that short-lived biomarkers, including urinary metabolites and DNA adducts of meat-related genotoxins, exist; however, they cannot be used as biomarkers of exposure in epidemiological studies, and do not belong in this section (see Section 4 for details).]

The accumulation of PhIP in hair may represent the first long-term biomarker of HAA exposure in cooked meats, although this biomarker is a measure of the unmetabolized chemical and not the biologically effective dose. Harmonization of the method across laboratories is required for validation and implementation in epidemiological studies.

Recent studies of omnivores have used metabolomics to identify constituents of meat in plasma and urine to measure meat consumption. Metabolomics is still a developing technology. It employs liquid chromatography-mass spectrometry (LC-MS)-based methods to identify the constituents or chemicals present in cooked or processed meat, and may provide reliable assessment of dietary habits and patterns of meat consumption in the future.

Data on the most promising biomarkers of exposure for red meat and processed meat consumption in epidemiological studies are summarized in Section 1.4.2(a) and (b).

*(a) Hair biomarkers*

Gas chromatography-mass spectrometry (GC-MS) with negative ion chemical ionization or liquid chromatography-tandem mass spectrometry (LC-MS/MS) with triple quadrupole-mass spectrometry (TQ-MS) instruments have been employed to measure PhIP in the hair of subjects in European countries ([Alexander et al., 2002](#)), Japan ([Kobayashi et al., 2005](#); [Kobayashi et al., 2007](#); [Iwasaki et al., 2014](#)), and the USA ([Bessette et al., 2009](#); [Turesky et al., 2013](#)). PhIP was identified in the hair of omnivores, but not in the hair of vegetarians ([Bessette et al., 2009](#)). The binding of PhIP to hair is strongly driven by melanin content, and binding levels of PhIP in hair should be normalized to melanin content ([Bessette et al., 2009](#); [Turesky et al., 2013](#)). [Turesky et al. \(2013\)](#) observed that, after being fed a semicontrolled diet, levels of PhIP in the hair of volunteers increased in an exposure-dependent manner. Levels of PhIP in hair were stable over time, varying in two meat eaters by less than 24% over a 6-month interval ([Turesky et al., 2013](#)). In another study, levels of PhIP in the hair of Japanese subjects were correlated with grilled/stir-fried meat intake, but not with grilled/stir-fried fish intake ([Kobayashi et al., 2005](#)). Levels of PhIP were further correlated with dietary HAA intake, according to an FFQ ([Kobayashi et al., 2007](#)). Since the binding of PhIP to hair is largely influenced by pigmentation, the biomonitoring of PhIP in an older population with predominantly white hair may be difficult. Moreover, because the growth cycles of individual hair follicles are asynchronous across the scalp, hair samples should be consistently collected from the same area of the scalp for comparison of PhIP levels in the hair of individuals ([Bessette et al., 2009](#); [Turesky et al., 2013](#)). Despite these limitations, biomonitoring of PhIP levels in hair is the first biomarker for assessing long-term exposure to this cooked meat carcinogen. Other HAAs bind

less efficiently to hair, and exposure cannot be assessed with hair ([Bessette et al., 2009](#); [Iwasaki et al., 2014](#)).

There are reports on the measurement by GC-NICI/MS of hydroxylated PAHs, including naphthalene and pyrene, in the hair of subjects ([Schummer et al., 2009](#); [Appenzeller et al., 2012](#); [Appenzeller & Tsatsakis, 2012](#)). Using hair to assess exposure to PAHs through meat consumption is challenging because of the multiple sources of exposure to PAHs and the low levels of hydroxylated PAHs in hair.

*(b) Urinary and plasma biomarkers*

Targeted approaches have been used to measure different procarcinogens, their metabolites, and DNA adducts in urine. More recently, untargeted metabolomics approaches have been used to understand dietary patterns of meat consumption, to strengthen self-administered FFQs.

*(i) Metabolomics: nutrients and secondary or indirect biomarkers*

Plasma and urine from subjects on different diets have been characterized by proton nuclear magnetic resonance, GC-MS, and LC-MS techniques, and hundreds of chemicals have been identified ([Puiggròs et al., 2011](#); [Hedrick et al., 2012](#); [Scalbert et al., 2014](#)).

Correlations were observed among chemical biomarkers of red meat, shellfish, fish, other food components, multivitamins, and diets, in plasma ([Guertin et al., 2014](#)). Several biomarkers in urine correlated to meat intake included: creatine, creatinine, carnitine, carnosine, ophidine, 1-methylhistidine, and 1-methylhistidine and 3-methylhistidine ([Dragsted, 2010](#); [Puiggròs et al., 2011](#)).

In a urinary metabolomic study employing LC-MS, 3-indoleacetyl-glucuronide, a microbiome metabolite of tryptophan, which is found at high concentrations in animal protein, was identified, possibly reflecting differences in the



protein sources between the diets ([Andersen et al., 2014](#)). Indole propionate was also identified as a potential biomarker of red meat in plasma ([Guertin et al., 2014](#)). Indoles are metabolites of tryptophan that are largely produced by the bacterial flora; however, they are not specific to meat, as they are also found in high amounts in soya and eggs ([Guertin et al., 2014](#)).

(ii) *Urinary 1-methylhistidine, 3-methylhistidine, creatinine, and taurine*

There were marked differences between the proton nuclear magnetic resonance spectra of high-red meat, low-red meat, and vegetarian diets, which included elevated urinary levels of creatinine, taurine, carnitine, trimethylamine-*N*-oxide, and methylhistidine in the high-red meat group. However, the spectral changes differentiating the low-red meat and vegetarian groups were subtle. The urinary metabolite trimethylamine-*N*-oxide, a product formed from carnitine by the bacterial microbiota, was associated with meat intake, but it is also a biomarker of fish intake, and may confound the interpretation of meat consumption patterns ([Stella et al., 2006](#)).

In another controlled meat-feeding study, the urinary excretion of creatinine, taurine, 1-methylhistidine, and 3-methylhistidine was investigated in individuals who consumed various amounts of red meat: vegetarian (0 g/day), low red meat (60 g/day), medium red meat (120 g/day), and high red meat (420 g/day) ([Cross et al., 2011](#)). All components demonstrated a significant dose-response relationship, increasing as red meat intake increased ( $P_{\text{trend}} < 0.0001$ ). There were significant differences in the mean levels of 1-methylhistidine and 3-methylhistidine across the four dietary intake groups ( $P < 0.01$  and  $P < 0.05$ , respectively). However, taurine and creatinine levels in the vegetarian and low-red meat intake groups could not be distinguished ( $P = 0.95$  and  $P = 0.88$ , respectively). 3-Methylhistidine and creatinine are

formed during muscle catabolism, thus lack specificity for meat intake. 1-Methylhistidine has also been found in the urine of subjects on a fish diet ([Lloyd et al., 2011](#)). Another study reported that the mean urinary levels of 1-methylhistidine and 3-methylhistidine did not differ among 131 colorectal adenoma and control subjects ( $P = 0.72$ ) ([Cross et al., 2014](#)). Thus, methylhistidine may not be a good indicator of meat processing conditions, and the levels of methylhistidine present in meat may not correlate to the levels of procarcinogens formed in cooked or processed meat.

To date, there are no chemical markers or metabolites of meat constituents that can provide information on the methods of meat processing and cooking that produce carcinogens.

## 1.5 Regulations and guidelines

In many countries, the production of red meat and processed meat is subject to stringent regulations. These regulations are primarily intended to prevent infectious diseases and minimize contamination of the meat products. Under the auspices of WHO and FAO, the Codex Alimentarius was established to provide international food standards, guidelines, and codes of practice to protect and promote safety, quality, and fairness in the international food trade ([Codex Alimentarius, 2015](#)). The scope of standards issued by the Codex Alimentarius is illustrative of standards and regulatory measures typically issued on a national basis for the maintenance of food safety in relation to meat products ([Table 1.18](#)).

An exhaustive list of all regional and national food authorities is not provided here, but a summary of those operating in Europe and the USA is provided.

EFSA ([EFSA, 2015](#)) is the EU risk assessment authority for food and feed safety. For red meat and processed meat, relevant EFSA panels or units include animal health and welfare, biological

**Table 1.18 Examples of meat-related food safety standards issued by Codex Alimentarius**

Reference	Standard	Committee	Last modified
CAC/GL 14-1991	Guide for the Microbiological Quality of Spices and Herbs Used in Processed Meat and Poultry Products	CCPMPP	1991
CAC/GL 78-2011	Guidelines for the Control of <i>Campylobacter</i> and <i>Salmonella</i> in Chicken Meat	CCFH	2011
CAC/GL 85-2014	Guidelines for the Control of <i>Taenia saginata</i> in Meat of Domestic Cattle	CCFH	2014
CAC/GL 86-2015	Guidelines for the Control of <i>Trichinella</i> spp. in Meat of Suidae	CCFH	2015
CAC/GL 87-2016	Guidelines for the Control of Nontyphoidal <i>Salmonella</i> spp. in Beef and Pork Meat	CCFH	2016
CAC/RCP 58-2005	Code of Hygienic Practice for Meat	CCMPH	2005
CODEX STAN 89-1981	Standard for Luncheon Meat	CCPMPP	2015
CODEX STAN 98-1981	Standard for Cooked Cured Chopped Meat	CCPMPP	2015

CCFH, Codex Committee on Food Hygiene; CCMPH, Codex Committee on Meat Hygiene; CCPMPP, Codex Committee on Processed Meat and Poultry Products

From [Codex Alimentarius \(2016a\)](#)

monitoring, contaminants, and assessment and methodological support.

In the USA, the relevant statutory authority for safety in relation to meat products is the United States Department of Agriculture ([Department of Agriculture, 2015](#)). The United States Food and Drug Administration (FDA) is responsible for regulating chemicals authorized in meat. A range of guidance documents and regulations are issued by this administration ([FDA, 2015](#)).

### 1.5.1 Prevention of infectious disease

The broad issues addressed by food safety regulations have been summarized by [Henson & Caswell \(1999\)](#), and include new potential food-borne risks, such as bovine spongiform encephalopathy and genetically modified organisms, as well as recognized risks posed by well-characterized bacteria. The scientific rationale for food safety regulations involves risk assessment, management, and communication.

For meat products, the regulations aim to decrease contamination by microbial pathogens (e.g. *Listeria monocytogenes*, *Escherichia coli*, and *Salmonella*) by minimizing cross-contamination

of other foods and water with enteric pathogens of animal origin ([Sofos, 2008](#)).

Many countries approach food safety, specifically in relation to meat production, through compliance with hazard analysis and critical control point (HACCP)-based regulations; HACCP is a safety and quality management tool ([Hudson et al., 1996](#)).

### 1.5.2 Prevention of contamination

#### (a) Red meat

Red meat may contain residues from veterinary drugs. These compounds are generally regulated at the national level, but 67 of them are regulated by international standards (i.e. maximum residue limits, MRLs) established by the [Codex Alimentarius \(2015\)](#). There is currently no international monitoring of the frequency of use of these chemicals.

Red meat is usually free of additives. However, in certain circumstances, colours are used for certification stamps on the surfaces of fresh cuts of meat, and are indicated in the food category system with a notation for “stamping, marking or branding the product” ([Codex Alimentarius, 2016a](#)).

Red meat may also contain chemicals present in the environment or used in the production of feed-like pesticide residues. When there is sufficient scientific information available about a chemical, the Joint FAO/WHO Expert Committee defines its acceptable daily intake (ADI), which is the amount of chemical, expressed based on body weight, that can be ingested over a lifetime without appreciable health risks. From the ADI, the Codex Alimentarius Commission establishes an MRL per kilogram of food that is recommended as being legally acceptable. The Codex Alimentarius Commission does not establish an MRL for a chemical if dietary exposure is above the ADI. Furthermore, no MRL is established if a chemical is assessed to be a genotoxic carcinogen in humans ([Codex Alimentarius, 2015](#)). MRLs have been established by the Codex Alimentarius for several pesticide residues possibly occurring in meat ([Codex Alimentarius, 2016b](#)). Most of these limits were established at the limit of detection of the analytical method.

Other chemical contaminants present in the environment, such as heavy metals or persistent organic pollutants, may also occur in red meat. Some of these contaminants are regulated internationally by the Codex Alimentarius. WHO/GEMS has collected national monitoring data on 145 environmental contaminants ([WHO, 2015b](#)). Moreover, the Codex Alimentarius has adopted codes of practice to reduce food and feed contamination by lead ([Codex Alimentarius, 2004](#)), by dioxin and dioxin-like PCBs ([Codex Alimentarius, 2006](#)), and by PAHs ([Codex Alimentarius, 2009](#)).

#### (b) Processed meat

National regulations are in place for processed meat in many countries around the world, e.g. in the USA ([Office of the Federal Register, 2015](#)). In Europe, the European Parliament and the Council of the EU define a “meat product” in Annex I to Regulation (EC) No 853/2004. The annex states that “meat products” means processed products

resulting from the processing of meat or from the further processing of such processed products, so that the cut surface shows that the product no longer has the characteristics of fresh meat” ([European Commission, 2004](#)).

At the international level, there is currently no active committee of the Codex Alimentarius to deal with meat (abolished in 1971) or processed meat (abolished in 1990), and the international standards for meat products are established by horizontal committees (e.g. committees for food additives, contaminants, or pesticide residues). In addition to the chemicals possibly present in meat in general, processed meat may contain food additives. However, many of these food additives, such as nitrites (80 mg/kg), colouring agents such as erythrosine (30 mg/kg), and antioxidants including butylated hydroxytoluene (100 mg/kg) are regulated by international standards established by the [Codex Alimentarius \(2016a\)](#).

## References

- Afshin A, Micha R, Khatibzadeh S, Fahimi S, Shi P, Powles J et al.; 2010 Global Burden of Diseases, Injuries, and Risk Factors Study: NUTRItition and ChrOnic Diseases Expert Group (NUTRICODE), and Metabolic Risk Factors of ChrOnic Diseases Collaborating Group (2015). The impact of dietary habits and metabolic risk factors on cardiovascular and diabetes mortality in countries of the Middle East and North Africa in 2010: a comparative risk assessment analysis. *BMJ Open*, 5(5):e006385. doi:[10.1136/bmjopen-2014-006385](#) PMID:[25995236](#)
- AFSSA (2005). Avis de l'Agence française de sécurité sanitaire des aliments relatif à l'évaluation de l'exposition de la population française aux dioxines, furanes et PCB de type dioxine. Saisine No.2005-SA-0372. Maisons-Alfort, France: l'Agence française de sécurité sanitaire des aliments. [French]
- Alaejos MS, Afonso AM (2011). Factors that affect the content of heterocyclic aromatic amines in foods. *Comp Rev Food Sci Food Safe*, 10(2):52–108. doi:[10.1111/j.1541-4337.2010.00141.x](#)
- Alexander J, Reistad R, Hegstad S, Frandsen H, Ingebrigtsen K, Paulsen JE et al. (2002). Biomarkers of exposure to heterocyclic amines: approaches to

- improve the exposure assessment. *Food Chem Toxicol*, 40(8):1131–7. doi:[10.1016/S0278-6915\(02\)00053-4](https://doi.org/10.1016/S0278-6915(02)00053-4) PMID:[12067575](https://pubmed.ncbi.nlm.nih.gov/12067575/)
- Alomirah H, Al-Zenki S, Al-Hooti S, Zaghloul S, Sawaya W, Ahmed N et al. (2011). Concentrations and dietary exposure to polycyclic aromatic hydrocarbons (PAHs) from grilled and smoked foods. *Food Contr*, 22(12):2028–35. doi:[10.1016/j.foodcont.2011.05.024](https://doi.org/10.1016/j.foodcont.2011.05.024)
- American Institute for Cancer Research/World Cancer Research Fund (1997). Food, nutrition, and the prevention of cancer: a global perspective. Washington (DC), USA: American Institute for Cancer Research.
- Andersen MB, Rinnan Å, Manach C, Poulsen SK, Pujos-Guillot E, Larsen TM et al. (2014). Untargeted metabolomics as a screening tool for estimating compliance to a dietary pattern. *J Proteome Res*, 13(3):1405–18. doi:[10.1021/pr400964s](https://doi.org/10.1021/pr400964s) PMID:[24444418](https://pubmed.ncbi.nlm.nih.gov/24444418/)
- ANSES (2011). Etude de l'alimentation totale française 2 (EAT 2). Avis de l'ANSES, rapport d'expertise. Résidus de pesticides, additifs, acrylamide, hydrocarbures aromatiques polycycliques. Maisons-Alfort, France: Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail. [French]
- ANSES (2012). Avis de l'Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail relatif aux analyses de retardateurs de flamme bromés (RFB) à mettre en œuvre dans le cadre des prochains plans de surveillance. Avis de l'Anses - Saisine No. 2010-SA-0225. Maisons-Alfort, France: Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail. [French]
- Appenzeller BM, Mathon C, Schummer C, Alkerwi A, Lair ML (2012). Simultaneous determination of nicotine and PAH metabolites in human hair specimen: a potential methodology to assess tobacco smoke contribution in PAH exposure. *Toxicol Lett*, 210(2):211–9. doi:[10.1016/j.toxlet.2011.11.022](https://doi.org/10.1016/j.toxlet.2011.11.022) PMID:[22155355](https://pubmed.ncbi.nlm.nih.gov/22155355/)
- Appenzeller BM, Tsatsakis AM (2012). Hair analysis for biomonitoring of environmental and occupational exposure to organic pollutants: state of the art, critical review and future needs. *Toxicol Lett*, 210(2):119–40. doi:[10.1016/j.toxlet.2011.10.021](https://doi.org/10.1016/j.toxlet.2011.10.021) PMID:[22079616](https://pubmed.ncbi.nlm.nih.gov/22079616/)
- Asgar MA, Fazilah A, Huda N, Bhat R, Karim AA (2010). Nonmeat protein alternatives as meat extenders and meat analogs. *Comp Rev Food Sci Food Safe*, 9(5):513–29. doi:[10.1111/j.1541-4337.2010.00124.x](https://doi.org/10.1111/j.1541-4337.2010.00124.x)
- Awadt WA, Ghareeb K, Bohm J (2012). Occurrence, health risks and methods of analysis for aflatoxins and ochratoxin A. *J Vet. Anim Sci*, 2(1):1–10.
- Back YM, Lee JH, Shin HS, Lee KG (2009). Analysis of heterocyclic amines and beta-carbolines by liquid chromatography–mass spectrometry in cooked meats commonly consumed in Korea. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess*, 26(3):298–305. doi:[10.1080/02652030802526834](https://doi.org/10.1080/02652030802526834) PMID:[19680902](https://pubmed.ncbi.nlm.nih.gov/19680902/)
- Balogh Z, Gray JI, Gomaa EA, Booren AM (2000). Formation and inhibition of heterocyclic aromatic amines in fried ground beef patties. *Food Chem Toxicol*, 38(5):395–401. doi:[10.1016/S0278-6915\(00\)00010-7](https://doi.org/10.1016/S0278-6915(00)00010-7) PMID:[10762724](https://pubmed.ncbi.nlm.nih.gov/10762724/)
- Barnes K, Collins T, Dion S, Reynolds H, Riess H, Stanzyk A et al. (2012). Importance of cattle biodiversity and its influence on the nutrient composition of beef. *Anim Front*, 2(4):54–60. doi:[10.2527/af.2012-0062](https://doi.org/10.2527/af.2012-0062)
- Bax ML, Aubry L, Ferreira C, Daudin JD, Gatellier P, Rémond D et al. (2012). Cooking temperature is a key determinant of in vitro meat protein digestion rate: investigation of underlying mechanisms. *J Agric Food Chem*, 60(10):2569–76. doi:[10.1021/jf205280y](https://doi.org/10.1021/jf205280y) PMID:[22335241](https://pubmed.ncbi.nlm.nih.gov/22335241/)
- Bax ML, Buffière C, Hafnaoui N, Gaudichon C, Savary-Auzeloux I, Dardevet D et al. (2013). Effects of meat cooking, and of ingested amount, on protein digestion speed and entry of residual proteins into the colon: a study in minipigs. *PLoS One*, 8(4):e61252 doi:[10.1371/journal.pone.0061252](https://doi.org/10.1371/journal.pone.0061252) PMID:[23593443](https://pubmed.ncbi.nlm.nih.gov/23593443/)
- Beaton GH (1994). Approaches to analysis of dietary data: relationship between planned analyses and choice of methodology. *Am J Clin Nutr*, 59(1):Suppl: 253S–61S. PMID:[8279436](https://pubmed.ncbi.nlm.nih.gov/8279436/)
- Bender A (1992). Meat and meat products in human nutrition in developing countries. Food & Nutrition Paper (Vol. 53). Rome, Italy: Food and Agriculture Organization of the United Nations. Available from: <http://www.fao.org/docrep/T0562E/T0562E00.HTM>
- Bessette EE, Yasa I, Dunbar D, Wilkens LR, Le Marchand L, Turesky RJ (2009). Biomonitoring of carcinogenic heterocyclic aromatic amines in hair: a validation study. *Chem Res Toxicol*, 22(8):1454–63. doi:[10.1021/tx900155f](https://doi.org/10.1021/tx900155f) PMID:[19588936](https://pubmed.ncbi.nlm.nih.gov/19588936/)
- Bingham SA, Day NE (1997). Using biochemical markers to assess the validity of prospective dietary assessment methods and the effect of energy adjustment. *Am J Clin Nutr*, 65(4):Suppl: 1130S–7S. PMID:[9094909](https://pubmed.ncbi.nlm.nih.gov/9094909/)
- Block G, Hartman AM, Dresser CM, Carroll MD, Gannon J, Gardner L (1986). A data-based approach to diet questionnaire design and testing. *Am J Epidemiol*, 124(3):453–69. PMID:[3740045](https://pubmed.ncbi.nlm.nih.gov/3740045/)
- Block G, Woods M, Potosky A, Clifford C (1990). Validation of a self-administered diet history questionnaire using multiple diet records. *J Clin Epidemiol*, 43(12):1327–35. doi:[10.1016/0895-4356\(90\)90099-B](https://doi.org/10.1016/0895-4356(90)90099-B) PMID:[2254769](https://pubmed.ncbi.nlm.nih.gov/2254769/)
- Bostick RM, Potter JD, Kushi LH, Sellers TA, Steinmetz KA, McKenzie DR et al. (1994). Sugar, meat, and fat intake, and non-dietary risk factors for colon cancer incidence in Iowa women (United States). *Cancer Causes Control*, 5(1):38–52. doi:[10.1007/BF01830725](https://doi.org/10.1007/BF01830725) PMID:[8123778](https://pubmed.ncbi.nlm.nih.gov/8123778/)
- Braekevelt E, Lau BP, Tague B, Popovic S, Tittlemier SA (2011). Effect of cooking on concentrations of  $\beta$ -estradiol and metabolites in model matrices and beef.

- J Agric Food Chem*, 59(3):915–20. doi:[10.1021/jf103064q](https://doi.org/10.1021/jf103064q) PMID:[21218831](https://pubmed.ncbi.nlm.nih.gov/21218831/)
- Brink M, Weijnenberg MP, de Goeij AF, Roemen GM, Lentjes MH, de Bruïne AP et al. (2005). Meat consumption and K-ras mutations in sporadic colon and rectal cancer in The Netherlands Cohort Study. *Br J Cancer*, 92(7):1310–20. doi:[10.1038/sj.bjc.6602491](https://doi.org/10.1038/sj.bjc.6602491) PMID:[15812479](https://pubmed.ncbi.nlm.nih.gov/15812479/)
- Butler LM, Sinha R, Millikan RC, Martin CF, Newman B, Gammon MD et al. (2003). Heterocyclic amines, meat intake, and association with colon cancer in a population-based study. *Am J Epidemiol*, 157(5):434–45. doi:[10.1093/aje/kwf221](https://doi.org/10.1093/aje/kwf221) PMID:[12615608](https://pubmed.ncbi.nlm.nih.gov/12615608/)
- Caan BJ, Slattery ML, Potter J, Quesenberry CPJ Jr, Coates AO, Schaffer DM (1998). Comparison of the Block and the Willett self-administered semiquantitative food frequency questionnaires with an interviewer-administered dietary history. *Am J Epidemiol*, 148(12):1137–47. doi:[10.1093/oxfordjournals.aje.a009598](https://doi.org/10.1093/oxfordjournals.aje.a009598) PMID:[9867257](https://pubmed.ncbi.nlm.nih.gov/9867257/)
- Cantwell M, Mittl B, Curtin J, Carroll R, Potischman N, Caporaso N et al. (2004). Relative validity of a food frequency questionnaire with a meat-cooking and heterocyclic amine module. *Cancer Epidemiol Biomarkers Prev*, 13(2):293–8. doi:[10.1158/1055-9965.EPI-270-2](https://doi.org/10.1158/1055-9965.EPI-270-2) PMID:[14973110](https://pubmed.ncbi.nlm.nih.gov/14973110/)
- Carroll RJ, Freedman LS, Kipnis V, Li L (1998). A new class of measurement-error models, with applications to dietary data. *Can J Stat*, 26(3):467–77. doi:[10.2307/3315770](https://doi.org/10.2307/3315770)
- Chattopadhyay MK (2014). Use of antibiotics as feed additives: a burning question. *Front Microbiol*, 5(334):334 PMID:[25071747](https://pubmed.ncbi.nlm.nih.gov/25071747/)
- Chen BH, Lin YS (1997). Formation of polycyclic aromatic hydrocarbons during processing of duck meat. *J Agric Food Chem*, 45(4):1394–403. doi:[10.1021/jf9606363](https://doi.org/10.1021/jf9606363)
- Chen G, Smith JS (2015). Determination of advanced glycation endproducts in cooked meat products. *Food Chem*, 168:190–5. doi:[10.1016/j.foodchem.2014.06.081](https://doi.org/10.1016/j.foodchem.2014.06.081) PMID:[25172699](https://pubmed.ncbi.nlm.nih.gov/25172699/)
- Chen J, Stampfer MJ, Hough HL, Garcia-Closas M, Willett WC, Hennekens CH et al. (1998). A prospective study of N-acetyltransferase genotype, red meat intake, and risk of colorectal cancer. *Cancer Res*, 58(15):3307–11. PMID:[9699660](https://pubmed.ncbi.nlm.nih.gov/9699660/)
- Chiu BC, Ji BT, Dai Q, Gridley G, McLaughlin JK, Gao YT et al. (2003). Dietary factors and risk of colon cancer in Shanghai, China. *Cancer Epidemiol Biomarkers Prev*, 12(3):201–8. PMID:[12646508](https://pubmed.ncbi.nlm.nih.gov/12646508/)
- Codex Alimentarius (2004). Code of practice for the prevention and reduction of lead contamination in foods (CAC/RCP 56-2004). Rome, Italy: World Health Organization/Food and Agriculture Organization of the United Nations. Available from: [http://www.codexalimentarius.org/download/standards/10099/CXP\\_056e.pdf](http://www.codexalimentarius.org/download/standards/10099/CXP_056e.pdf)
- Codex Alimentarius (2006). Code of practice for the reduction of dioxin and dioxin-like PCB contamination in foods and feeds (CAC/RCP 62-2006). Rome, Italy: World Health Organization/Food and Agriculture Organization of the United Nations. Available from: [http://www.codexalimentarius.org/download/standards/10693/CXP\\_062e.pdf](http://www.codexalimentarius.org/download/standards/10693/CXP_062e.pdf)
- Codex Alimentarius (2009). Code of practice for the reduction of contamination of food with polycyclic aromatic hydrocarbons (PAH) from smoking and direct drying processes (CAC/RCP 68-2009). Rome, Italy: World Health Organization/Food and Agriculture Organization of the United Nations. Available from: [http://www.codexalimentarius.org/download/standards/11257/CXP\\_068e.pdf](http://www.codexalimentarius.org/download/standards/11257/CXP_068e.pdf)
- Codex Alimentarius (2015). International food standards. Maximum residue limits (MRLs) and risk management recommendations (RMRs) for residues of veterinary drugs in foods. CAC/MRL 2-2015. Updated at the 38th Session of the Codex Alimentarius Commission (July 2015). Rome, Italy: World Health Organization/Food and Agriculture Organization of the United Nations. Available from: [http://fao.org/input/download/standards/45/MRL2\\_2015e.pdf](http://fao.org/input/download/standards/45/MRL2_2015e.pdf)
- Codex Alimentarius (2016a). General standard for food additives. Codex STAN 192-1995. Adopted in 1995. Revised in 2016. Rome, Italy: World Health Organization/Food and Agriculture Organization of the United Nations. Available from: [www.fao.org/gsaonline/docs/CXS\\_192e.pdf](http://www.fao.org/gsaonline/docs/CXS_192e.pdf)
- Codex Alimentarius (2016b). Pesticide residues in food and feed. Codex pesticides residues in food online database. Rome, Italy: World Health Organization/Food and Agriculture Organization of the United Nations. Available from: <http://www.fao.org/fao-who-codexalimentarius/codex-texts/dbs/pestres/en/>
- Cross AJ, Ferrucci LM, Risch A, Graubard BI, Ward MH, Park Y et al. (2010). A large prospective study of meat consumption and colorectal cancer risk: an investigation of potential mechanisms underlying this association. *Cancer Res*, 70(6):2406–14. doi:[10.1158/0008-5472.CAN-09-3929](https://doi.org/10.1158/0008-5472.CAN-09-3929) PMID:[20215514](https://pubmed.ncbi.nlm.nih.gov/20215514/)
- Cross AJ, Major JM, Rothman N, Sinha R (2014). Urinary 1-methylhistidine and 3-methylhistidine, meat intake, and colorectal adenoma risk. *Eur J Cancer Prev*, 23(5):385–90. doi:[10.1097/CEJ.0000000000000027](https://doi.org/10.1097/CEJ.0000000000000027) PMID:[24681531](https://pubmed.ncbi.nlm.nih.gov/24681531/)
- Cross AJ, Major JM, Sinha R (2011). Urinary biomarkers of meat consumption. *Cancer Epidemiol Biomarkers Prev*, 20(6):1107–11. doi:[10.1158/1055-9965.EPI-11-0048](https://doi.org/10.1158/1055-9965.EPI-11-0048) PMID:[21527577](https://pubmed.ncbi.nlm.nih.gov/21527577/)
- Cross AJ, Peters U, Kirsh VA, Andriole GL, Reding D, Hayes RB et al. (2005). A prospective study of meat and meat mutagens and prostate cancer risk. *Cancer Res*, 65(24):11779–84. doi:[10.1158/0008-5472.CAN-05-2191](https://doi.org/10.1158/0008-5472.CAN-05-2191) PMID:[16357191](https://pubmed.ncbi.nlm.nih.gov/16357191/)

- De Mey E, De Klerck K, De Maere H, Dewulf L, Derdelinckx G, Peeters MC et al. (2014). The occurrence of N-nitrosamines, residual nitrite and biogenic amines in commercial dry fermented sausages and evaluation of their occasional relation. *Meat Sci*, 96(2 Pt A):821–8. doi:[10.1016/j.meatsci.2013.09.010](https://doi.org/10.1016/j.meatsci.2013.09.010) PMID:[24200576](https://pubmed.ncbi.nlm.nih.gov/24200576/)
- De Mey E, De Maere H, Paelinck H, Fraeye I (2015). Volatile N-nitrosamines in meat products: potential precursors, influence of processing and mitigation strategies. *Crit Rev Food Sci Nutr*, 57(13):2909–2923 doi:[10.1080/10408398.2015.1078769](https://doi.org/10.1080/10408398.2015.1078769) PMID:[26528731](https://pubmed.ncbi.nlm.nih.gov/26528731/)
- Dellavalle CT, Daniel CR, Aschebrook-Kilfoy B, Hollenbeck AR, Cross AJ, Sinha R et al. (2013). Dietary intake of nitrate and nitrite and risk of renal cell carcinoma in the NIH-AARP Diet and Health Study. *Br J Cancer*, 108(1):205–12. doi:[10.1038/bjc.2012.522](https://doi.org/10.1038/bjc.2012.522) PMID:[23169285](https://pubmed.ncbi.nlm.nih.gov/23169285/)
- Dennis MJ, Massey RC, McWeeny DJ, Knowles ME, Watson D (1983). Analysis of polycyclic aromatic hydrocarbons in UK total diets. *Food Chem Toxicol*, 21(5):569–74. doi:[10.1016/0278-6915\(83\)90142-4](https://doi.org/10.1016/0278-6915(83)90142-4) PMID:[6686183](https://pubmed.ncbi.nlm.nih.gov/6686183/)
- Department of Agriculture (2015). Federal Meat Inspection Act. Washington (DC), USA: United States Department of Agriculture, Food Safety and Inspection Service. Available from: [http://www.fsis.usda.gov/wps/portal/](http://www.fsis.usda.gov/wps/portal/fsis/topics/rulemaking/federal-meat-inspection-act)
- Dixon LB, Balder HF, Virtanen MJ, Rashidkhani B, Männistö S, Krogh V et al. (2004). Dietary patterns associated with colon and rectal cancer: results from the Dietary Patterns and Cancer (DIETSCAN) Project. *Am J Clin Nutr*, 80(4):1003–11. PMID:[15447912](https://pubmed.ncbi.nlm.nih.gov/15447912/)
- Dragsted LO (2010). Biomarkers of meat intake and the application of nutrigenomics. *Meat Sci*, 84(2):301–7. doi:[10.1016/j.meatsci.2009.08.028](https://doi.org/10.1016/j.meatsci.2009.08.028) PMID:[20374789](https://pubmed.ncbi.nlm.nih.gov/20374789/)
- EFSA (2003). Opinion of the Scientific Panel on Biological Hazards on the request from the Commission related to the effects of nitrites/nitrates on the microbiological safety of meat products. *EFSA J*, 14:1–31.
- EFSA (2004a). Opinion of the Scientific Panel on Contaminants in the Food Chain on a request from the Commission related to aflatoxin B1 as undesirable substance in animal feed. *EFSA J*, 39:1–27.
- EFSA (2004b). Opinion of the Scientific Panel on Contaminants in the Food Chain on a request from the Commission related to deoxynivalenol (DON) as undesirable substance in animal feed. *EFSA J*, 73:1–41.
- EFSA (2004c). Opinion of the Scientific Panel on Contaminants in the Food Chain on a request from the Commission related to zearalenone as undesirable substance in animal feed. *EFSA J*, 89:1–35.
- EFSA (2004d). Opinion of the Scientific Panel on Contaminants in the Food Chain on a request from the Commission related to ochratoxin A (OTA) as undesirable substance in animal feed. *EFSA J*, 101:1–36.
- EFSA (2005c). Opinion of the Scientific Panel on Contaminants in the Food Chain on a request from the Commission related to fumonisins as undesirable substances in animal feed. *EFSA J*, 235:1–32.
- EFSA (2007). Opinion of the scientific panel on contaminations in the food chain on a request from the European Commission related to hormone residues in bovine meat and meat products. Parma, Italy: European Food Safety Authority. *EFSA J*. 510:1–62.
- EFSA (2008). Polycyclic aromatic hydrocarbons in food. Scientific Opinion of the Panel on Contaminants in the Food Chain. *EFSA J*, 724:1–114.
- EFSA (2011). The EFSA Comprehensive European Food Consumption Database [online database]. Parma, Italy: European Food Safety Authority. Available from: <http://www.efsa.europa.eu/en/food-consumption/comprehensive-database>, accessed September 2015
- EFSA (2012). Cadmium dietary exposure in the European population. *EFSA J*, 10(1):2551. <http://dx.doi.org/10.2903/j.efsa.2012.2551>. Available from: [www.efsa.europa.eu/efsajournal](http://www.efsa.europa.eu/efsajournal) doi:[10.2903/j.efsa.2012.2551](https://doi.org/10.2903/j.efsa.2012.2551)
- EFSA (2015). European Food Safety Authority [website]. Parma, Italy: European Food Safety Authority. Available from: <http://www.efsa.europa.eu/>
- Egeberg R, Olsen A, Christensen J, Halkjær J, Jakobsen MU, Overvad K et al. (2013). Associations between red meat and risks for colon and rectal cancer depend on the type of red meat consumed. *J Nutr*, 143(4):464–72. doi:[10.3945/jn.112.168799](https://doi.org/10.3945/jn.112.168799) PMID:[23427329](https://pubmed.ncbi.nlm.nih.gov/23427329/)
- Engel E, Ratel J, Bouhlef J, Planche C, Meurillon M (2015). Novel approaches to improving the chemical safety of the meat chain towards toxicants. *Meat Sci*, 109:75–85. doi:[10.1016/j.meatsci.2015.05.016](https://doi.org/10.1016/j.meatsci.2015.05.016) PMID:[26043665](https://pubmed.ncbi.nlm.nih.gov/26043665/)
- English DR, MacInnis RJ, Hodge AM, Hopper JL, Haydon AM, Giles GG (2004). Red meat, chicken, and fish consumption and risk of colorectal cancer. *Cancer Epidemiol Biomarkers Prev*, 13(9):1509–14. PMID:[15342453](https://pubmed.ncbi.nlm.nih.gov/15342453/)
- EPA (2010). An exposure assessment of polybrominated diphenyl ethers. Report No. EPA/600/R-08/086F. Washington (DC), USA: National Center for Environmental Assessment, Office of Research and Development, United States Environmental Protection Agency.
- European Commission (2003). Regulation (EC) No 1831/2003 of the European Parliament and of the Council of 22 September 2003 on additives for use in animal nutrition. *Official Journal of the European Union L*, 268:29–43.
- European Commission (2004). Regulation (EC) No. 853/2004 of the European Parliament and of the Council of 29 April 2004 laying down specific hygiene rules for food of animal origin. Available from: <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2004:226:0022:0082:EN:PDF>
- FAO (2015). Food Balance. FAOSTAT [online database]. Food and Agriculture Organization of the United Nations Statistics Division. Rome, Italy: Food and

- Agriculture Organization of the United Nations. Available from: [http://faostat3.fao.org/browse/FB/\\*E](http://faostat3.fao.org/browse/FB/*E), accessed 9 July 2015
- FAO/WHO (2015). The FAO/WHO Chronic individual food consumption database – summary statistics (CIFOCS). Food and Agriculture Organization of the United Nations/World Health Organization. Available from: <http://www.who.int/foodsafety/databases/en/>
- FCID (2015). What we eat in America - Food commodity intake database 2005-10. Commodity consumption calculator. United States Environmental Protection Agency, Office of Pesticide Programs © University of Maryland 2012-2016. Available from: <http://fcid.foodrisk.org/percentiles.php>
- FDA (2005). Subject: Illegal residues in meat, poultry, seafood, and other animal derived foods. Chapter: Post-approval monitoring of animal drugs, feeds and devices In: Compliance program guidance manual. United States Food and Drug Administration.
- FDA (2008). Subject: Mycotoxins in domestic and imported food. Chapter 07: Molecular biology and natural toxins. In: Compliance program guidance manual. United States Food and Drug Administration.
- FDA (2015). Guidance and regulation. Washington (DC), USA: United States Department of Agriculture, Food and Drug Administration. Available from: <http://www.fda.gov/Food/GuidanceRegulation>
- Felton JS, Knize MG, Roper M, Fultz E, Shen NH, Turteltaub KW (1992). Chemical analysis, prevention, and low-level dosimetry of heterocyclic amines from cooked food. *Cancer Res*, 52(7):Suppl: 2103s–7s. PMID:1544148
- Ferrucci LM, Cross AJ, Graubard BI, Brinton LA, McCarty CA, Ziegler RG et al. (2009). Intake of meat, meat mutagens, and iron and the risk of breast cancer in the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial. *Br J Cancer*, 101(1):178–84. doi:10.1038/sj.bjc.6605118 PMID:19513076
- Flood A, Velie EM, Sinha R, Chatterjee N, Lacey JV Jr, Schairer C et al. (2003). Meat, fat, and their subtypes as risk factors for colorectal cancer in a prospective cohort of women. *Am J Epidemiol*, 158(1):59–68. doi:10.1093/aje/kwg099 PMID:12835287
- Fung T, Hu FB, Fuchs C, Giovannucci E, Hunter DJ, Stampfer MJ et al. (2003). Major dietary patterns and the risk of colorectal cancer in women. *Arch Intern Med*, 163(3):309–14. doi:10.1001/archinte.163.3.309 PMID:12578511
- Gibis M, Weiss J (2010). Inhibitory effect of marinades with hibiscus extract on formation of heterocyclic aromatic amines and sensory quality of fried beef patties. *Meat Sci*, 85(4):735–42. doi:10.1016/j.meatsci.2010.03.034 PMID:20418021
- Gibis M, Weiss J (2012). Antioxidant capacity and inhibitory effect of grape seed and rosemary extract in marinades on the formation of heterocyclic amines in fried beef patties. *Food Chem*, 134(2):766–74. doi:10.1016/j.foodchem.2012.02.179 PMID:23107689
- Gille D, Schmid A (2015). Vitamin B12 in meat and dairy products. *Nutr Rev*, 73(2):106–15. doi:10.1093/nutrit/nuu011 PMID:26024497
- Giri A, Khummueng W, Mercier F, Kondjoyan N, Tournayre P, Meurillon M et al. (2015). Relevance of two-dimensional gas chromatography and high resolution olfactometry for the parallel determination of heat-induced toxicants and odorants in cooked food. *J Chromatogr A*, 1388:217–26. doi:10.1016/j.chroma.2015.01.045 PMID:25728653
- Givens DI (2005). The role of animal nutrition in improving the nutritive value of animal-derived foods in relation to chronic disease. *Proc Nutr Soc*, 64(3):395–402. doi:10.1079/PNS2005448 PMID:16048674
- Goldberg T, Cai W, Peppas M, Dardaine V, Baliga BS, Uribarri J et al. (2004). Advanced glycoxidation end products in commonly consumed foods. *J Am Diet Assoc*, 104(8):1287–91. doi:10.1016/j.jada.2004.05.214 PMID:15281050
- Guerrero Legarreta I (2010). Canned products and pâté. In: Toldra F, editor. Handbook of meat processing. Blackwell Publishing. Part 2, Chapter 19; 337–349. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/9780813820897.ch19.summary>
- Guertin KA, Moore SC, Sampson JN, Huang WY, Xiao Q, Stolzenberg-Solomon RZ et al. (2014). Metabolomics in nutritional epidemiology: identifying metabolites associated with diet and quantifying their potential to uncover diet-disease relations in populations. *Am J Clin Nutr*, 100(1):208–17. doi:10.3945/ajcn.113.078758 PMID:24740205
- Haorah J, Zhou L, Wang X, Xu G, Mirvish SS (2001). Determination of total N-nitroso compounds and their precursors in frankfurters, fresh meat, dried salted fish, sauces, tobacco, and tobacco smoke particulates. *J Agric Food Chem*, 49(12):6068–78. doi:10.1021/jf010602h PMID:11743810
- Hedlund M, Padler-Karavani V, Varki NM, Varki A (2008). Evidence for a human-specific mechanism for diet and antibody-mediated inflammation in carcinoma progression. *Proc Natl Acad Sci U S A*, 105(48):18936–41. doi:10.1073/pnas.0803943105 PMID:19017806
- Hedrick VE, Dietrich AM, Estabrooks PA, Savla J, Serrano E, Davy BM (2012). Dietary biomarkers: advances, limitations and future directions. *Nutr J*, 11(1):109 doi:10.1186/1475-2891-11-109 PMID:23237668
- Heinz G, Hautzinger P (2007). Meat processing technology for small- to medium-scale producers. Rome, Italy: Food and Agriculture Organization of the United Nations. Available from: <http://www.fao.org/documents/card/fr/c/fb92d00f-7ff3-593a-a77c-7b19003b2554/>
- Henson S, Caswell J (1999). Food safety regulation: an overview of contemporary issues. *Food Policy*, 24(6):589–603. doi:10.1016/S0306-9192(99)00072-X

- Heshmati A. (2015). Impact of cooking procedures on antibacterial drug residues in foods: a review. *Journal of Food Quality & Hazards Control*, 2:33–37.
- Hill RJ, Davies PS (2001). The validity of self-reported energy intake as determined using the doubly labelled water technique. *Br J Nutr*, 85(4):415–30. doi:[10.1079/BJN2000281](https://doi.org/10.1079/BJN2000281) PMID:[11348556](https://pubmed.ncbi.nlm.nih.gov/11348556/)
- Honikel KO (2008). The use and control of nitrate and nitrite for the processing of meat products. *Meat Sci*, 78(1–2):68–76. doi:[10.1016/j.meatsci.2007.05.030](https://doi.org/10.1016/j.meatsci.2007.05.030) PMID:[22062097](https://pubmed.ncbi.nlm.nih.gov/22062097/)
- Honikel KO (2010). Curing. Chapter 6. In: Toldra F, editor. *Handbook of meat processing*. Blackwell Publishing; 125–41. doi:[10.1002/9780813820897.ch6](https://doi.org/10.1002/9780813820897.ch6)
- Hori T, Nakagawa R, Tobiishi K, Iida T, Tsutsumi T, Sasaki K et al. (2005). Effects of cooking on concentrations of polychlorinated dibenzo-p-dioxins and related compounds in fish and meat. *J Agric Food Chem*, 53(22):8820–8. doi:[10.1021/jf050978l](https://doi.org/10.1021/jf050978l) PMID:[16248590](https://pubmed.ncbi.nlm.nih.gov/16248590/)
- Hu FB (2002). Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol*, 13(1):3–9. doi:[10.1097/00041433-200202000-00002](https://doi.org/10.1097/00041433-200202000-00002) PMID:[11790957](https://pubmed.ncbi.nlm.nih.gov/11790957/)
- Hu FB, Stampfer MJ, Rimm E, Ascherio A, Rosner BA, Spiegelman D et al. (1999). Dietary fat and coronary heart disease: a comparison of approaches for adjusting for total energy intake and modeling repeated dietary measurements. *Am J Epidemiol*, 149(6):531–40. doi:[10.1093/oxfordjournals.aje.a009849](https://doi.org/10.1093/oxfordjournals.aje.a009849) PMID:[10084242](https://pubmed.ncbi.nlm.nih.gov/10084242/)
- Hudson WR, Mead GC, Hinton MH (1996). Relevance of abattoir hygiene assessment to microbial contamination of British beef carcasses. *Vet Rec*, 139(24):587–9. PMID:[8981733](https://pubmed.ncbi.nlm.nih.gov/8981733/)
- Hull GLJ, Woodside JV, Ames JM, Cuskelly GJ (2012). N<sup>ε</sup>-(Carboxymethyl)lysine content of foods commonly consumed in a Western style diet. *Food Chem*, 131(1):170–4. doi:[10.1016/j.foodchem.2011.08.055](https://doi.org/10.1016/j.foodchem.2011.08.055) PMID:[23017409](https://pubmed.ncbi.nlm.nih.gov/23017409/)
- Hygreeva D, Pandey MC, Radhakrishna K (2014). Potential applications of plant based derivatives as fat replacers, antioxidants and antimicrobials in fresh and processed meat products. *Meat Sci*, 98(1):47–57. doi:[10.1016/j.meatsci.2014.04.006](https://doi.org/10.1016/j.meatsci.2014.04.006) PMID:[24845336](https://pubmed.ncbi.nlm.nih.gov/24845336/)
- IARC (1993). Some naturally occurring substances: food items and constituents, heterocyclic aromatic amines and mycotoxins. IARC Monogr Eval Carcinog Risks Hum, 56:1–599. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol56/index.php>
- IARC (2010a). Some non-heterocyclic polycyclic aromatic hydrocarbons and some related exposures. IARC Monogr Eval Carcinog Risks Hum, 92:1–853. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol92/index.php> PMID:[21141735](https://pubmed.ncbi.nlm.nih.gov/21141735/)
- IARC (2010b). Ingested nitrate and nitrite, and cyanobacterial peptide toxins. IARC Monogr Eval Carcinog Risks Hum, 94:1–448. PMID:21141240. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol94/index.php>
- IARC (2012a). Arsenic, metals, fibres, and dusts. IARC Monogr Eval Carcinog Risks Hum, 100C:1–499. PMID:23189751. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol100C/index.php>
- IARC (2012b). Chemical agents and related occupations. IARC Monogr Eval Carcinog Risks Hum, 100F:1–599. PMID:23189753. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol100F/index.php>
- IARC (2016). Polychlorinated biphenyls and polybrominated biphenyls. IARC Monogr Eval Carcinog Risks Hum, 107:1–502. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol107/index.php>
- Imamura F, Micha R, Khatibzadeh S, Fahimi S, Shi P, Powles J et al. ; Global Burden of Diseases Nutrition and Chronic Diseases Expert Group (NutriCoDE) (2015). Dietary quality among men and women in 187 countries in 1990 and 2010: a systematic assessment. *Lancet Glob Health*, 3(3):e132–42. doi:[10.1016/S2214-109X\(14\)70381-X](https://doi.org/10.1016/S2214-109X(14)70381-X) PMID:[25701991](https://pubmed.ncbi.nlm.nih.gov/25701991/)
- IPCS (2009). Principles and methods for the risk assessment of chemicals in food. Environmental Health Criteria 240. Geneva, Switzerland: International Programme on Chemical Safety, Food and Agriculture Organization of the United Nations/World Health Organization. Available from: <http://www.who.int/foodsafety/publications/chemical-food/en/>
- Isam T, Kadim IT, Mahgoub O (2007). Postharvest handling of red meat. In: Rahman MS editor. *Handbook of food preservation*. New York, USA: Marcel Dekker; 173.
- Iwasaki M, Mukai T, Takachi R, Ishihara J, Totsuka Y, Tsugane S (2014). Validity of a self-administered food frequency questionnaire in the estimation of heterocyclic aromatic amines. *Cancer Causes Control*, 25(8):1015–28. doi:[10.1007/s10552-014-0401-7](https://doi.org/10.1007/s10552-014-0401-7) PMID:[24890804](https://pubmed.ncbi.nlm.nih.gov/24890804/)
- Järvinen R, Knekt P, Hakulinen T, Rissanen H, Heliövaara M (2001). Dietary fat, cholesterol and colorectal cancer in a prospective study. *Br J Cancer*, 85(3):357–61. doi:[10.1054/bjoc.2001.1906](https://doi.org/10.1054/bjoc.2001.1906) PMID:[11487265](https://pubmed.ncbi.nlm.nih.gov/11487265/)
- JECFA (2000). Toxicological evaluation of certain veterinary drug residues in food: Estradiol-17 $\alpha$ , progesterone, and testosterone. Prepared by the Fifty-second meeting of the Joint FAO/WHO Expert Committee on Food Additives. Geneva, Switzerland: World Health Organization. WHO Food Additives Series: 43. Available from: <http://www.inchem.org/documents/jecfa/jecmono/v43jec05.htm>
- JECFA (2002). Polychlorinated dibenzodioxins (PCDDs), polychlorinated dibenzofurans (PCDFs), and coplanar polychlorinated biphenyls (PCBs). *Evaluations of the Joint FAO/WHO Expert Committee on Food Additives*. Geneva, Switzerland: World Health Organization. Available from: <http://apps.who>



- [int/food-additives-contaminants-jecfa-database/chemical.aspx?chemID=2753](http://int/food-additives-contaminants-jecfa-database/chemical.aspx?chemID=2753)
- JECFA (2011). Acrylamide. In: Safety evaluation of certain contaminants in food. Prepared by the Seventy-second meeting of the Joint FAO/WHO Expert Committee on Food Additives. WHO Food Additives Series: 63, FAO JECFA Monographs 8.
- JECFA (2013). Cadmium. Evaluations of the Joint FAO/WHO Committee on Food Additives. Geneva, Switzerland: World Health Organization. Available from: <http://apps.who.int/food-additives-contaminants-jecfa-database/chemical.aspx?chemID=1376>
- Jeong S-H, Kang D, Lim M-W, Kang CS, Sung HJ (2010). Risk assessment of growth hormones and antimicrobial residues in meat. *Toxicol Res*, 26(4):301–13. doi:[10.5487/TR.2010.26.4.301](https://doi.org/10.5487/TR.2010.26.4.301) PMID:[24278538](https://pubmed.ncbi.nlm.nih.gov/24278538/)
- Kabat GC, Miller AB, Jain M, Rohan TE (2007). A cohort study of dietary iron and heme iron intake and risk of colorectal cancer in women. *Br J Cancer*, 97(1):118–22. doi:[10.1038/sj.bjc.6603837](https://doi.org/10.1038/sj.bjc.6603837) PMID:[17551493](https://pubmed.ncbi.nlm.nih.gov/17551493/)
- Kan CA, Meijer GAL (2007). The risk of contamination of food with toxic substances present in animal feed. *Anim Feed Sci Technol*, 133(1–2):84–108. doi:[10.1016/j.anifeeds.2006.08.005](https://doi.org/10.1016/j.anifeeds.2006.08.005)
- Kato I, Akhmedkhanov A, Koenig K, Toniolo PG, Shore RE, Riboli E (1997). Prospective study of diet and female colorectal cancer: the New York University Women's Health Study. *Nutr Cancer*, 28(3):276–81. doi:[10.1080/01635589709514588](https://doi.org/10.1080/01635589709514588) PMID:[9343837](https://pubmed.ncbi.nlm.nih.gov/9343837/)
- Kazerouni N, Sinha R, Hsu C-H, Greenberg A, Rothman N (2001). Analysis of 200 food items for benzo[a]pyrene and estimation of its intake in an epidemiologic study. *Food Chem Toxicol*, 39(5):423–36. doi:[10.1016/S0278-6915\(00\)00158-7](https://doi.org/10.1016/S0278-6915(00)00158-7) PMID:[11313108](https://pubmed.ncbi.nlm.nih.gov/11313108/)
- Keating GA, Bogen KT (2004). Estimates of heterocyclic amine intake in the US population. *J Chromatogr B Analyt Technol Biomed Life Sci*, 802(1):127–33. doi:[10.1016/j.jchromb.2003.10.047](https://doi.org/10.1016/j.jchromb.2003.10.047) PMID:[15036004](https://pubmed.ncbi.nlm.nih.gov/15036004/)
- Kesse E, Clavel-Chapelon F, Boutron-Ruault MC (2006). Dietary patterns and risk of colorectal tumors: a cohort of French women of the National Education System (E3N). *Am J Epidemiol*, 164(11):1085–93. doi:[10.1093/aje/kwj324](https://doi.org/10.1093/aje/kwj324) PMID:[16990408](https://pubmed.ncbi.nlm.nih.gov/16990408/)
- Khan MR, Busquets R, Santos FJ, Puignou L (2008). New method for the analysis of heterocyclic amines in meat extracts using pressurised liquid extraction and liquid chromatography–tandem mass spectrometry. *J Chromatogr A*, 1194(2):155–60. doi:[10.1016/j.chroma.2008.04.058](https://doi.org/10.1016/j.chroma.2008.04.058) PMID:[18490022](https://pubmed.ncbi.nlm.nih.gov/18490022/)
- Kilfoy BA, Zhang Y, Park Y, Holford TR, Schatzkin A, Hollenbeck A et al. (2011). Dietary nitrate and nitrite and the risk of thyroid cancer in the NIH-AARP Diet and Health Study. *Int J Cancer*, 129(1):160–72. doi:[10.1002/ijc.25650](https://doi.org/10.1002/ijc.25650) PMID:[20824705](https://pubmed.ncbi.nlm.nih.gov/20824705/)
- Kipnis V, Freedman LS, Brown CC, Hartman AM, Schatzkin A, Wacholder S (1997). Effect of measurement error on energy-adjustment models in nutritional epidemiology. *Am J Epidemiol*, 146(10):842–55. doi:[10.1093/oxfordjournals.aje.a009202](https://doi.org/10.1093/oxfordjournals.aje.a009202) PMID:[9384205](https://pubmed.ncbi.nlm.nih.gov/9384205/)
- Knize MG, Brown ED, Salmon CP, Levander OA, Felton JS, Rothman N (1998). Heterocyclic amine content in restaurant-cooked hamburgers, steaks, ribs and chicken. *J Food Ag. Chem.*, 46(11):4648–51. doi:[10.1021/jf980639a](https://doi.org/10.1021/jf980639a)
- Kobayashi M, Hanaoka T, Hashimoto H, Tsugane S (2005). 2-Amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) level in human hair as biomarkers for dietary grilled/stir-fried meat and fish intake. *Mutat Res*, 588(2):136–42. doi:[10.1016/j.mrgentox.2005.09.008](https://doi.org/10.1016/j.mrgentox.2005.09.008) PMID:[16289877](https://pubmed.ncbi.nlm.nih.gov/16289877/)
- Kobayashi M, Hanaoka T, Tsugane S (2007). Validity of a self-administered food frequency questionnaire in the assessment of heterocyclic amine intake using 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) levels in hair. *Mutat Res*, 630(1–2):14–9. doi:[10.1016/j.mrgentox.2007.02.003](https://doi.org/10.1016/j.mrgentox.2007.02.003) PMID:[17392018](https://pubmed.ncbi.nlm.nih.gov/17392018/)
- Krause R, Knoll K, Henle T (2003). Studies on the formation of furosine and pyridosine during acid hydrolysis of different Amadori products of lysine. *Eur Food Res Technol*, 216(4):277–83. doi:[10.1007/s00217-002-0649-0](https://doi.org/10.1007/s00217-002-0649-0)
- Kushi LH (1994). Gaps in epidemiologic research methods: design considerations for studies that use food-frequency questionnaires. *Am J Clin Nutr*, 59(1):Suppl:180S–4S. PMID:[8279420](https://pubmed.ncbi.nlm.nih.gov/8279420/)
- Larsen JC (2006). Risk assessments of polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans, and dioxin-like polychlorinated biphenyls in food. *Mol Nutr Food Res*, 50(10):885–96. doi:[10.1002/mnfr.200500247](https://doi.org/10.1002/mnfr.200500247) PMID:[17009211](https://pubmed.ncbi.nlm.nih.gov/17009211/)
- Lawrie RA (1998). *Lawrie's meat science*. Cambridge, England: Woodhead Publishing Ltd.
- Lawrie RA, Ledward DA (2006). Chapter 5: The conversion of muscle to meat. In: Lawrie RA, editor. *Lawrie's meat science*. Cambridge, England: Woodhead Publishing Ltd.
- Ledesma E, Rendueles M, Díaz M (2014). Benzo(a)pyrene penetration on a smoked meat product during smoking time. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess*, 31(10):1688–98. doi:[10.1080/19440049.2014.949875](https://doi.org/10.1080/19440049.2014.949875) PMID:[25078362](https://pubmed.ncbi.nlm.nih.gov/25078362/)
- Lee DH, Anderson KE, Folsom AR, Jacobs DR Jr (2005). Heme iron, zinc and upper digestive tract cancer: the Iowa Women's Health Study. *Int J Cancer*, 117(4):643–7. doi:[10.1002/ijc.21215](https://doi.org/10.1002/ijc.21215) PMID:[15929082](https://pubmed.ncbi.nlm.nih.gov/15929082/)
- Lee SA, Shu XO, Yang G, Li H, Gao YT, Zheng W (2009). Animal origin foods and colorectal cancer risk: a report from the Shanghai Women's Health Study. *Nutr Cancer*, 61(2):194–205. doi:[10.1080/01635580802419780](https://doi.org/10.1080/01635580802419780) PMID:[19235035](https://pubmed.ncbi.nlm.nih.gov/19235035/)

- Linseisen J, Kesse E, Slimani N, Bueno-De-Mesquita HB, Ocké MC, Skeie G et al. (2002). Meat consumption in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohorts: results from 24-hour dietary recalls. *Public Health Nutr*, 5(6B):6B: 1243–58. doi:[10.1079/PHN2002402](https://doi.org/10.1079/PHN2002402) PMID:[12639230](https://pubmed.ncbi.nlm.nih.gov/12639230/)
- Linseisen J, Rohrmann S, Norat T, Gonzalez CA, Dorransoro Iraeta M, Morote Gómez P et al. (2006). Dietary intake of different types and characteristics of processed meat which might be associated with cancer risk - results from the 24-hour diet recalls in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Public Health Nutr*, 9(4):449–64. doi:[10.1079/PHN2005861](https://doi.org/10.1079/PHN2005861) PMID:[16870017](https://pubmed.ncbi.nlm.nih.gov/16870017/)
- Lloyd AJ, Favé G, Beckmann M, Lin W, Tailliant K, Xie L et al. (2011). Use of mass spectrometry fingerprinting to identify urinary metabolites after consumption of specific foods. *Am J Clin Nutr*, 94(4):981–91. doi:[10.3945/ajcn.111.017921](https://doi.org/10.3945/ajcn.111.017921) PMID:[21865330](https://pubmed.ncbi.nlm.nih.gov/21865330/)
- Lombardi-Boccia G, Martinez-Dominguez B, Aguzzi A (2002). Total heme and non-heme iron in raw and cooked meats. *J Food Sci*, 67(5):1738–41. doi:[10.1111/j.1365-2621.2002.tb08715.x](https://doi.org/10.1111/j.1365-2621.2002.tb08715.x)
- Lorenzo JM, Crecente S, Franco D, Sarriés MV, Gómez M (2014). The effect of livestock production system and concentrate level on carcass traits and meat quality of foals slaughtered at 18 months of age. *Animal*, 8(3):494–503. doi:[10.1017/S175173111300236X](https://doi.org/10.1017/S175173111300236X) PMID:[24398030](https://pubmed.ncbi.nlm.nih.gov/24398030/)
- Lorenzo JM, Fuciños C, Purriños L, Franco D (2010). Intramuscular fatty acid composition of “Galician Mountain” foals breed: effect of sex, slaughtered age and livestock production system. *Meat Sci*, 86(3):825–31. doi:[10.1016/j.meatsci.2010.07.004](https://doi.org/10.1016/j.meatsci.2010.07.004) PMID:[20675062](https://pubmed.ncbi.nlm.nih.gov/20675062/)
- Lyche JL, Rosseland C, Berge G, Polder A (2015). Human health risk associated with brominated flame-retardants (BFRs). *Environ Int*, 74:170–80. doi:[10.1016/j.envint.2014.09.006](https://doi.org/10.1016/j.envint.2014.09.006) PMID:[25454234](https://pubmed.ncbi.nlm.nih.gov/25454234/)
- Manabe S, Izumikawa S, Asakuno K, Wada O, Kanai Y (1991). Detection of carcinogenic amino-alpha-carbolines and amino-gamma-carbolines in diesel-exhaust particles. *Environ Pollut*, 70(3):255–65. doi:[10.1016/0269-7491\(91\)90013-M](https://doi.org/10.1016/0269-7491(91)90013-M) PMID:[15092136](https://pubmed.ncbi.nlm.nih.gov/15092136/)
- Marroquín-Cardona AG, Johnson NM, Phillips TD, Hayes AW (2014). Mycotoxins in a changing global environment – a review. *Food Chem Toxicol*, 69:220–30. doi:[10.1016/j.fct.2014.04.025](https://doi.org/10.1016/j.fct.2014.04.025) PMID:[24769018](https://pubmed.ncbi.nlm.nih.gov/24769018/)
- McCann SE, Marshall JR, Trevisan M, Russell M, Muti P, Markovic N et al. (1999). Recent alcohol intake as estimated by the health habits and history questionnaire, the Harvard semiquantitative food frequency questionnaire, and a more detailed alcohol intake questionnaire. *Am J Epidemiol*, 150(4):334–40. doi:[10.1093/oxfordjournals.aje.a010012](https://doi.org/10.1093/oxfordjournals.aje.a010012) PMID:[10453809](https://pubmed.ncbi.nlm.nih.gov/10453809/)
- Mottier P, Parisod V, Turesky RJ (2000). Quantitative determination of polycyclic aromatic hydrocarbons in barbecued meat sausages by gas chromatography coupled to mass spectrometry. *J Agric Food Chem*, 48(4):1160–6. doi:[10.1021/jf991205y](https://doi.org/10.1021/jf991205y) PMID:[10775366](https://pubmed.ncbi.nlm.nih.gov/10775366/)
- Murkovic M (2004). Formation of heterocyclic aromatic amines in model systems. *J Chromatogr B Analyt Technol Biomed Life Sci*, 802(1):3–10. doi:[10.1016/j.jchromb.2003.09.026](https://doi.org/10.1016/j.jchromb.2003.09.026) PMID:[15035991](https://pubmed.ncbi.nlm.nih.gov/15035991/)
- Nachman KE, Smith TJ (2015). Hormone use in food animal production: assessing potential dietary exposures and breast cancer risk. *Curr Environ Health Rep*, 2(1):1–14. doi:[10.1007/s40572-014-0042-8](https://doi.org/10.1007/s40572-014-0042-8) PMID:[26231238](https://pubmed.ncbi.nlm.nih.gov/26231238/)
- NCI (2017). CHARRED: Computerized Heterocyclic Amines Resource for Research in Epidemiology of Disease. Bethesda (MD), USA: Division of Cancer Epidemiology and Genetics, National Cancer Institute. Available from: <http://dceg.cancer.gov/tools/design/charred>
- Ni W, McNaughton L, LeMaster DM, Sinha R, Turesky RJ (2008). Quantitation of 13 heterocyclic aromatic amines in cooked beef, pork, and chicken by liquid chromatography–electrospray ionization/tandem mass spectrometry. *J Agric Food Chem*, 56(1):68–78. doi:[10.1021/jf072461a](https://doi.org/10.1021/jf072461a) PMID:[18069786](https://pubmed.ncbi.nlm.nih.gov/18069786/)
- Norat T, Bingham S, Ferrari P, Slimani N, Jenab M, Mazuir M et al. (2005). Meat, fish, and colorectal cancer risk: the European Prospective Investigation into Cancer and Nutrition. *J Natl Cancer Inst*, 97(12):906–16. doi:[10.1093/jnci/dji164](https://doi.org/10.1093/jnci/dji164) PMID:[15956652](https://pubmed.ncbi.nlm.nih.gov/15956652/)
- Ockermann HW, Basu L (2010). Chapters 9 and 10: Fermented meat products: production and consumption. In: Toldrá F, editor. *Handbook of meat processing*. Columbus (OH), USA: The Ohio State University.
- Office of the Federal Register (2015). Part 319—definitions and standards of identity or composition. Title 9 - Animals and animal products (9 CFR Part 319). United States Code of Federal Regulations. Washington (DC), USA: United States Government Publishing Office. Available from: <http://www.gpo.gov/fdsys/pkg/CFR-2015-title9-vol2/xml/CFR-2015-title9-vol2-part319.xml>
- Olatunji OS, Fatoki OS, Opeolu BO, Ximba BJ (2014). Determination of polycyclic aromatic hydrocarbons [PAHs] in processed meat products using gas chromatography – flame ionization detector. *Food Chem*, 156:296–300. doi:[10.1016/j.foodchem.2014.01.120](https://doi.org/10.1016/j.foodchem.2014.01.120) PMID:[24629971](https://pubmed.ncbi.nlm.nih.gov/24629971/)
- Ollberding NJ, Wilkens LR, Henderson BE, Kolonel LN, Le Marchand L (2012). Meat consumption, heterocyclic amines and colorectal cancer risk: the Multiethnic Cohort Study. *Int J Cancer*, 131(7):E1125–33. doi:[10.1002/ijc.27546](https://doi.org/10.1002/ijc.27546) PMID:[22438055](https://pubmed.ncbi.nlm.nih.gov/22438055/)
- Pala V, Krogh V, Berrino F, Sieri S, Grioni S, Tjønneland A et al. (2009). Meat, eggs, dairy products, and risk of breast cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohort. *Am J Clin*

- Nutr*, 90(3):602–12. doi:[10.3945/ajcn.2008.27173](https://doi.org/10.3945/ajcn.2008.27173) PMID:[19491385](https://pubmed.ncbi.nlm.nih.gov/19491385/)
- Paris A, Andre F, Antignac JP, Le Bizet B, Bonneau M, Briant C et al. (2006). Hormones et promoteurs de croissance en productions animales: de la physiologie à l'évaluation du risque. *INRA Prod Anim*, 19(3):151–240. [French]
- Parnell WR, Blakey CW, Smith C (2012). Secondary analysis of Adult Nutrition Survey 2008/09 for intake of Beef and Lamb for the New Zealand population and for consumers. Dunedin, New Zealand: LINZ® Nutrition and Activity Research Unit, University of Otago, Report No.: 2012.138.
- Pearson AM, Gillett TA (1996). Curing. Chapter 3. In: Pearson AM, Gillett TA, editors. *Processed meats*, 3rd Edition. Maryland, USA: Aspen Publishers, Inc; 53–78.
- Perelló G, Martí-Cid R, Castell V, Llobet JM, Domingo JL (2010). Influence of various cooking processes on the concentrations of PCDD/PCDFs, PCBs and PCDEs in foods. *Food Contr*, 21(2):178–85. doi:[10.1016/j.foodcont.2009.05.003](https://doi.org/10.1016/j.foodcont.2009.05.003)
- Perelló G, Martí-Cid R, Llobet JM, Domingo JL (2008). Effects of various cooking processes on the concentrations of arsenic, cadmium, mercury, and lead in foods. *J Agric Food Chem*, 56(23):11262–9. doi:[10.1021/jf802411q](https://doi.org/10.1021/jf802411q) PMID:[18986150](https://pubmed.ncbi.nlm.nih.gov/18986150/)
- Pietinen P, Malila N, Virtanen M, Hartman TJ, Tangrea JA, Albanes D et al. (1999). Diet and risk of colorectal cancer in a cohort of Finnish men. *Cancer Causes Control*, 10(5):387–96. doi:[10.1023/A:1008962219408](https://doi.org/10.1023/A:1008962219408) PMID:[10530608](https://pubmed.ncbi.nlm.nih.gov/10530608/)
- Preussmann R, Stewart BW (1984). N-Nitroso carcinogens. In: Searle CE, editor. *Chemical carcinogens*. Washington (DC), USA: American Chemical Society; 643–828.
- Pryer JA, Nichols R, Elliott P, Thakrar B, Brunner E, Marmot M (2001). Dietary patterns among a national random sample of British adults. *J Epidemiol Community Health*, 55(1):29–37. doi:[10.1136/jech.55.1.29](https://doi.org/10.1136/jech.55.1.29) PMID:[11112948](https://pubmed.ncbi.nlm.nih.gov/11112948/)
- Puiggròs F, Solà R, Bladé C, Salvadó MJ, Arola L (2011). Nutritional biomarkers and foodomic methodologies for qualitative and quantitative analysis of bioactive ingredients in dietary intervention studies. *J Chromatogr A*, 1218(42):7399–414. doi:[10.1016/j.chroma.2011.08.051](https://doi.org/10.1016/j.chroma.2011.08.051) PMID:[21917262](https://pubmed.ncbi.nlm.nih.gov/21917262/)
- Purchas RW, Busboom JR (2005). The effect of production system and age on levels+ of iron, taurine, carnosine, coenzyme Q(10), and creatine in beef muscles and liver. *Meat Sci*, 70(4):589–96. doi:[10.1016/j.meatsci.2005.02.008](https://doi.org/10.1016/j.meatsci.2005.02.008) PMID:[22063884](https://pubmed.ncbi.nlm.nih.gov/22063884/)
- Purchas RW, Busboom JR, Wilkinson BH (2006). Changes in the forms of iron and in concentrations of taurine, carnosine, coenzyme Q(10), and creatine in beef longissimus muscle with cooking and simulated stomach and duodenal digestion. *Meat Sci*, 74(3):443–9. doi:[10.1016/j.meatsci.2006.03.015](https://doi.org/10.1016/j.meatsci.2006.03.015) PMID:[22063048](https://pubmed.ncbi.nlm.nih.gov/22063048/)
- Rahman MS (2007). Food preservation overview. In: Rahman MS, editor. *Handbook of food preservation*. New York, USA: Marcel Dekker; 635–66.
- Rimm EB, Giovannucci EL, Stampfer MJ, Colditz GA, Litin LB, Willett WC (1992). Reproducibility and validity of an expanded self-administered semiquantitative food frequency questionnaire among male health professionals. *Am J Epidemiol*, 135(10):1114–26, discussion 1127–36. PMID:[1632423](https://pubmed.ncbi.nlm.nih.gov/1632423/)
- Rohrmann S, Linseisen J, Becker N, Norat T, Sinha R, Skeie G et al. ; European Prospective Investigation into Cancer and Nutrition (2002). Cooking of meat and fish in Europe – results from the European Prospective Investigation into Cancer and Nutrition (EPIC). *Eur J Clin Nutr*, 56(12):1216–30. doi:[10.1038/sj.ejcn.1601494](https://doi.org/10.1038/sj.ejcn.1601494) PMID:[12494307](https://pubmed.ncbi.nlm.nih.gov/12494307/)
- Rohrmann S, Zoller D, Hermann S, Linseisen J (2007). Intake of heterocyclic aromatic amines from meat in the European Prospective Investigation into Cancer and Nutrition (EPIC)-Heidelberg cohort. *Br J Nutr*, 98(6):1112–5. doi:[10.1017/S000711450778145X](https://doi.org/10.1017/S000711450778145X) PMID:[18309547](https://pubmed.ncbi.nlm.nih.gov/18309547/)
- Roseiro LC, Gomes A, Santos C (2011). Influence of processing in the prevalence of polycyclic aromatic hydrocarbons in a Portuguese traditional meat product. *Food Chem Toxicol*, 49(6):1340–5. doi:[10.1016/j.fct.2011.03.017](https://doi.org/10.1016/j.fct.2011.03.017) PMID:[21419819](https://pubmed.ncbi.nlm.nih.gov/21419819/)
- Rywotycycki R (2007). The effect of baking of various kinds of raw meat from different animal species and meat with functional additives on nitrosamine contamination level. *Food Chem*, 101(2):540–8. doi:[10.1016/j.foodchem.2006.02.012](https://doi.org/10.1016/j.foodchem.2006.02.012)
- Samraj AN, Pearce OMT, Läubli H, Crittenden AN, Bergfeld AK, Banda K et al. (2015). A red meat-derived glycan promotes inflammation and cancer progression. *Proc Natl Acad Sci U S A*, 112(2):542–7. doi:[10.1073/pnas.1417508112](https://doi.org/10.1073/pnas.1417508112) PMID:[25548184](https://pubmed.ncbi.nlm.nih.gov/25548184/)
- Santarelli RL, Pierre F, Corpet DE (2008). Processed meat and colorectal cancer: a review of epidemiologic and experimental evidence. *Nutr Cancer*, 60(2):131–44. doi:[10.1080/01635580701684872](https://doi.org/10.1080/01635580701684872) PMID:[18444144](https://pubmed.ncbi.nlm.nih.gov/18444144/)
- Santé-Lhoutellier V, Astruc T, Marinova P, Greve E, Gatellier P (2008). Effect of meat cooking on physicochemical state and in vitro digestibility of myofibrillar proteins. *J Agric Food Chem*, 56(4):1488–94. doi:[10.1021/jf072999g](https://doi.org/10.1021/jf072999g) PMID:[18237130](https://pubmed.ncbi.nlm.nih.gov/18237130/)
- Sato Y, Nakaya N, Kuriyama S, Nishino Y, Tsubono Y, Tsuji I (2006). Meat consumption and risk of colorectal cancer in Japan: the Miyagi Cohort Study. *Eur J Cancer Prev*, 15(3):211–8. doi:[10.1097/01.cej.0000197455.87356.05](https://doi.org/10.1097/01.cej.0000197455.87356.05) PMID:[16679863](https://pubmed.ncbi.nlm.nih.gov/16679863/)
- Scalbert A, Brennan L, Manach C, Andres-Lacueva C, Dragsted LO, Draper J et al. (2014). The food metabolome: a window over dietary exposure. *Am J Clin*

- Nutr*, 99(6):1286–308. doi:[10.3945/ajcn.113.076133](https://doi.org/10.3945/ajcn.113.076133) PMID:[24760973](https://pubmed.ncbi.nlm.nih.gov/24760973/)
- Schechter A, Harris TR, Shah N, Musumba A, Pöpke O (2008). Brominated flame retardants in US food. *Mol Nutr Food Res*, 52(2):266–72. doi:[10.1002/mnfr.200700166](https://doi.org/10.1002/mnfr.200700166) PMID:[18040989](https://pubmed.ncbi.nlm.nih.gov/18040989/)
- Schonfeldt HC, Hall NG (2011). Determining iron bio-availability with a constant heme iron value. *J Food Compos Anal*, 24(4–5):738–40. doi:[10.1016/j.jfca.2011.01.002](https://doi.org/10.1016/j.jfca.2011.01.002)
- Schroeder H (2010). Les hydrocarbures aromatiques polycycliques présentent-ils un risque de neurotoxicité développementale? ANSES Bulletin veille scientifique santé environnement travail; 11:83–8. Available from: <https://www.anses.fr/fr/system/files/BVS-mg-011-SCHROEDER.pdf> [French]
- Schummer C, Appenzeller BM, Millet M, Wennig R (2009). Determination of hydroxylated metabolites of polycyclic aromatic hydrocarbons in human hair by gas chromatography–negative chemical ionization mass spectrometry. *J Chromatogr A*, 1216(32):6012–9. doi:[10.1016/j.chroma.2009.05.068](https://doi.org/10.1016/j.chroma.2009.05.068) PMID:[19577242](https://pubmed.ncbi.nlm.nih.gov/19577242/)
- Sempos CT, Liu K, Ernst ND (1999). Food and nutrient exposures: what to consider when evaluating epidemiologic evidence. *Am J Clin Nutr*, 69(6):1330S–8S. PMID:[10357757](https://pubmed.ncbi.nlm.nih.gov/10357757/)
- Shin H-S, Strasburg GM, Gray JI (2002). A model system study of the inhibition of heterocyclic aromatic amine formation by organosulfur compounds. *J Agric Food Chem*, 50(26):7684–90. doi:[10.1021/jf025707e](https://doi.org/10.1021/jf025707e) PMID:[12475289](https://pubmed.ncbi.nlm.nih.gov/12475289/)
- Sikorski ZE, Kalakowski E (2010). Smoking. In: Toldra F, editor. Handbook of meat processing. Oxford, UK: Wiley-Blackwell. doi:[10.1002/9780813820897.ch12](https://doi.org/10.1002/9780813820897.ch12)
- Sindelar JJ, Milkowski AL (2012). Human safety controversies surrounding nitrate and nitrite in the diet. *Nitric Oxide*, 26(4):259–66. doi:[10.1016/j.niox.2012.03.011](https://doi.org/10.1016/j.niox.2012.03.011) PMID:[22487433](https://pubmed.ncbi.nlm.nih.gov/22487433/)
- Sindelar JJ, Sebranek JG, Bacus JN (2010). Uncured, natural, and organic processed meat products. Meat processing technology series. Champaign (IL), USA: American Meat Science Association; 1–42. Available from <http://www.lulu.com/shop/jeffrey-sindelar/uncured-natural-and-organic-processed-meat-products/paperback/product-12200306.html>
- Singh PN, Fraser GE (1998). Dietary risk factors for colon cancer in a low-risk population. *Am J Epidemiol*, 148(8):761–74. doi:[10.1093/oxfordjournals.aje.a009697](https://doi.org/10.1093/oxfordjournals.aje.a009697) PMID:[9786231](https://pubmed.ncbi.nlm.nih.gov/9786231/)
- Sinha R, Knize MG, Salmon CP, Brown ED, Rhodes D, Felton JS et al. (1998a). Heterocyclic amine content of pork products cooked by different methods and to varying degrees of doneness. *Food Chem Toxicol*, 36(4):289–97. doi:[10.1016/S0278-6915\(97\)00159-2](https://doi.org/10.1016/S0278-6915(97)00159-2) PMID:[9651045](https://pubmed.ncbi.nlm.nih.gov/9651045/)
- Sinha R, Rothman N, Brown ED, Salmon CP, Knize MG, Swanson CA et al. (1995). High concentrations of the carcinogen 2-amino-1-methyl-6-phenylimidazo-[4,5-b]pyridine (PhIP) occur in chicken but are dependent on the cooking method. *Cancer Res*, 55(20):4516–9. PMID:[7553619](https://pubmed.ncbi.nlm.nih.gov/7553619/)
- Sinha R, Rothman N, Salmon CP, Knize MG, Brown ED, Swanson CA et al. (1998b). Heterocyclic amine content in beef cooked by different methods to varying degrees of doneness and gravy made from meat drippings. *Food Chem Toxicol*, 36(4):279–87. doi:[10.1016/S0278-6915\(97\)00162-2](https://doi.org/10.1016/S0278-6915(97)00162-2) PMID:[9651044](https://pubmed.ncbi.nlm.nih.gov/9651044/)
- Skog K, Solyakov A, Jägerstad MI (2000). Effects of heating conditions and additives on the formation of heterocyclic amines with reference to amino-carbolines in a meat juice model system. *Food Chem Toxicol*, 68(3):299–308. doi:[10.1016/S0308-8146\(99\)00195-8](https://doi.org/10.1016/S0308-8146(99)00195-8)
- Skog KI, Johansson MA, Jägerstad MI (1998). Carcinogenic heterocyclic amines in model systems and cooked foods: a review on formation, occurrence and intake. *Food Chem Toxicol*, 36(9–10):879–96. doi:[10.1016/S0278-6915\(98\)00061-1](https://doi.org/10.1016/S0278-6915(98)00061-1) PMID:[9737435](https://pubmed.ncbi.nlm.nih.gov/9737435/)
- Sobell J, Block G, Koslowe P, Tobin J, Andres R (1989). Validation of a retrospective questionnaire assessing diet 10–15 years ago. *Am J Epidemiol*, 130(1):173–87. PMID:[2741904](https://pubmed.ncbi.nlm.nih.gov/2741904/)
- Sofos JN (2008). Challenges to meat safety in the 21st century. *Meat Sci*, 78(1–2):3–13. doi:[10.1016/j.meatsci.2007.07.027](https://doi.org/10.1016/j.meatsci.2007.07.027) PMID:[22062090](https://pubmed.ncbi.nlm.nih.gov/22062090/)
- Stadler RH, Lineback DR (2009). Process-induced food toxicants, occurrence, formation, mitigation and health risks. Hoboken (NJ), USA: John Wiley & Sons.
- Stella C, Beckwith-Hall B, Cloarec O, Holmes E, Lindon JC, Powell J et al. (2006). Susceptibility of human metabolic phenotypes to dietary modulation. *J Proteome Res*, 5(10):2780–8. doi:[10.1021/pr060265y](https://doi.org/10.1021/pr060265y) PMID:[17022649](https://pubmed.ncbi.nlm.nih.gov/17022649/)
- Subar AF, Midthune D, Kulldorff M, Brown CC, Thompson FE, Kipnis V et al. (2000). Evaluation of alternative approaches to assign nutrient values to food groups in food frequency questionnaires. *Am J Epidemiol*, 152(3):279–86. doi:[10.1093/aje/152.3.279](https://doi.org/10.1093/aje/152.3.279) PMID:[10933275](https://pubmed.ncbi.nlm.nih.gov/10933275/)
- Subar AF, Thompson FE, Smith AF, Jobe JB, Ziegler RG, Potischman N et al. (1995). Improving food frequency questionnaires: a qualitative approach using cognitive interviewing. *J Am Diet Assoc*, 95(7):781–8, quiz 789–90. doi:[10.1016/S0002-8223\(95\)00217-0](https://doi.org/10.1016/S0002-8223(95)00217-0) PMID:[7797809](https://pubmed.ncbi.nlm.nih.gov/7797809/)
- Subar AF, Ziegler RG, Thompson FE, Johnson CC, Weissfeld JL, Reding D et al.; Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial Investigators (2001). Is shorter always better? Relative importance of questionnaire length and cognitive ease on response rates and data quality for two dietary questionnaires. *Am J Epidemiol*, 153(4):404–9. doi:[10.1093/aje/153.4.404](https://doi.org/10.1093/aje/153.4.404) PMID:[11207159](https://pubmed.ncbi.nlm.nih.gov/11207159/)

- Sugimura T, Wakabayashi K, Nakagama H, Nagao M (2004). Heterocyclic amines: mutagens/carcinogens produced during cooking of meat and fish. *Cancer Sci*, 95(4):290–9. doi:[10.1111/j.1349-7006.2004.tb03205.x](https://doi.org/10.1111/j.1349-7006.2004.tb03205.x) PMID:[15072585](https://pubmed.ncbi.nlm.nih.gov/15072585/)
- Sun XD, Holley RA (2011). Factors influencing gel formation by myofibrillar proteins in muscle foods. *Comp Rev Food Sci Food Safe*, 10(1):33–51. doi:[10.1111/j.1541-4337.2010.00137.x](https://doi.org/10.1111/j.1541-4337.2010.00137.x)
- Sy MM, Feinberg M, Verger P, Barré T, Cléménçon S, Crépet A (2013). New approach for the assessment of cluster diets. *Food Chem Toxicol*, 52:180–7. doi:[10.1016/j.fct.2012.11.005](https://doi.org/10.1016/j.fct.2012.11.005) PMID:[23182740](https://pubmed.ncbi.nlm.nih.gov/23182740/)
- Takachi R, Tsubono Y, Baba K, Inoue M, Sasazuki S, Iwasaki M et al.; Japan Public Health Center-Based Prospective Study Group (2011). Red meat intake may increase the risk of colon cancer in Japanese, a population with relatively low red meat consumption. *Asia Pac J Clin Nutr*, 20(4):603–12. PMID:[22094846](https://pubmed.ncbi.nlm.nih.gov/22094846/)
- Tangvoranuntakul P, Gagneux P, Diaz S, Bardor M, Varki N, Varki A et al. (2003). Human uptake and incorporation of an immunogenic nonhuman dietary sialic acid. *Proc Natl Acad Sci U S A*, 100(21):12045–50. doi:[10.1073/pnas.2131556100](https://doi.org/10.1073/pnas.2131556100) PMID:[14523234](https://pubmed.ncbi.nlm.nih.gov/14523234/)
- Tareke E, Rydberg P, Karlsson P, Eriksson S, Törnqvist M (2002). Analysis of acrylamide, a carcinogen formed in heated foodstuffs. *J Agric Food Chem*, 50(17):4998–5006. doi:[10.1021/jf020302f](https://doi.org/10.1021/jf020302f) PMID:[12166997](https://pubmed.ncbi.nlm.nih.gov/12166997/)
- Tavani A, La Vecchia C, Gallus S, Lagiou P, Trichopoulos D, Levi F et al. (2000). Red meat intake and cancer risk: a study in Italy. *Int J Cancer*, 86(3):425–8. doi:[10.1002/\(SICI\)1097-0215\(20000501\)86:3<425::AID-IJC19>3.0.CO;2-S](https://doi.org/10.1002/(SICI)1097-0215(20000501)86:3<425::AID-IJC19>3.0.CO;2-S) PMID:[10760833](https://pubmed.ncbi.nlm.nih.gov/10760833/)
- Thompson FE, Kipnis V, Midthune D, Freedman LS, Carroll RJ, Subar AF et al. (2008). Performance of a food-frequency questionnaire in the US NIH-AARP (National Institutes of Health-American Association of Retired Persons) Diet and Health Study. *Public Health Nutr*, 11(2):183–95. doi:[10.1017/S1368980007000419](https://doi.org/10.1017/S1368980007000419) PMID:[17610761](https://pubmed.ncbi.nlm.nih.gov/17610761/)
- Thompson FE, Subar AF (2013). Dietary assessment methodology. In: Coulston AM, Boushey CJ, Ferruzzi MG, editors. *Nutrition in the prevention and treatment of disease*. 3rd ed. Elsevier. doi:[10.1016/B978-0-12-391884-0.00001-9](https://doi.org/10.1016/B978-0-12-391884-0.00001-9)
- Tiemersma EW, Kampman E, Bueno de Mesquita HB, Bunschoten A, van Schothorst EM, Kok FJ et al. (2002). Meat consumption, cigarette smoking, and genetic susceptibility in the etiology of colorectal cancer: results from a Dutch prospective study. *Cancer Causes Control*, 13(4):383–93. doi:[10.1023/A:1015236701054](https://doi.org/10.1023/A:1015236701054) PMID:[12074508](https://pubmed.ncbi.nlm.nih.gov/12074508/)
- Toldrá F (2010). *Handbook of meat processing*. Oxford, UK: Wiley-Blackwell. doi:[10.1002/9780813820897](https://doi.org/10.1002/9780813820897)
- Tran NL, Barraji L, Smith K, Javier A, Burke TA (2004). Combining food frequency and survey data to quantify long-term dietary exposure: a methyl mercury case study. *Risk Anal*, 24(1):19–30. doi:[10.1111/j.0272-4332.2004.00408.x](https://doi.org/10.1111/j.0272-4332.2004.00408.x) PMID:[15027997](https://pubmed.ncbi.nlm.nih.gov/15027997/)
- Tricker AR (1997). N-Nitroso compounds and man: sources of exposure, endogenous formation and occurrence in body fluids. *Eur J Cancer Prev*, 6(3):226–68. doi:[10.1097/00008469-199706000-00003](https://doi.org/10.1097/00008469-199706000-00003) PMID:[9306073](https://pubmed.ncbi.nlm.nih.gov/9306073/)
- Turesky RJ, Liu L, Gu D, Yonemori KM, White KK, Wilkens LR et al. (2013). Biomonitoring the cooked meat carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine in hair: impact of exposure, hair pigmentation, and cytochrome P450 1A2 phenotype. *Cancer Epidemiol Biomarkers Prev*, 22(3):356–64. doi:[10.1158/1055-9965.EPI-12-1206](https://doi.org/10.1158/1055-9965.EPI-12-1206) PMID:[23329727](https://pubmed.ncbi.nlm.nih.gov/23329727/)
- University of Otago and Ministry of Health (2011). Methodology report for the 2008/09 New Zealand Adult Nutrition Survey. Wellington, New Zealand: Ministry of Health. Available from: <http://www.health.govt.nz/system/files/documents/publications/methodology-report.pdf>, accessed August 2015
- Uribarri J, Woodruff S, Goodman S, Cai W, Chen X, Pyzik R et al. (2010). Advanced glycation end products in foods and a practical guide to their reduction in the diet. *J Am Diet Assoc*, 110(6):911–16.e12. doi:[10.1016/j.jada.2010.03.018](https://doi.org/10.1016/j.jada.2010.03.018) PMID:[20497781](https://pubmed.ncbi.nlm.nih.gov/20497781/)
- Vitaglione P, Fogliano V (2004). Use of antioxidants to minimize the human health risk associated to mutagenic/carcinogenic heterocyclic amines in food. *J Chromatogr B Analyt Technol Biomed Life Sci*, 802(1):189–99. doi:[10.1016/j.jchromb.2003.09.029](https://doi.org/10.1016/j.jchromb.2003.09.029) PMID:[15036011](https://pubmed.ncbi.nlm.nih.gov/15036011/)
- Walker R (1990). Nitrates, nitrites and N-nitrosocompounds: a review of the occurrence in food and diet and the toxicological implications. *Food Addit Contam*, 7(6):717–68. doi:[10.1080/02652039009373938](https://doi.org/10.1080/02652039009373938) PMID:[2079111](https://pubmed.ncbi.nlm.nih.gov/2079111/)
- Ward MH, Cantor KP, Riley D, Merkle S, Lynch CF (2003). Nitrate in public water supplies and risk of bladder cancer. *Epidemiology*, 14(2):183–90. doi:[10.1097/01.EDE.0000050664.28048.DF](https://doi.org/10.1097/01.EDE.0000050664.28048.DF) PMID:[12606884](https://pubmed.ncbi.nlm.nih.gov/12606884/)
- Ward MH, Cerhan JR, Colt JS, Hartge P (2006). Risk of non-Hodgkin lymphoma and nitrate and nitrite from drinking water and diet. *Epidemiology*, 17(4):375–82. doi:[10.1097/01.ede.0000219675.79395.9f](https://doi.org/10.1097/01.ede.0000219675.79395.9f) PMID:[16699473](https://pubmed.ncbi.nlm.nih.gov/16699473/)
- Wei EK, Giovannucci E, Wu K, Rosner B, Fuchs CS, Willett WC et al. (2004). Comparison of risk factors for colon and rectal cancer. *Int J Cancer*, 108(3):433–42. doi:[10.1002/ijc.11540](https://doi.org/10.1002/ijc.11540) PMID:[14648711](https://pubmed.ncbi.nlm.nih.gov/14648711/)
- Weiss J, Gibis M, Schuh V, Salminen H (2010). Advances in ingredient and processing systems for meat and meat products. *Meat Sci*, 86(1):196–213. doi:[10.1016/j.meatsci.2010.05.008](https://doi.org/10.1016/j.meatsci.2010.05.008) PMID:[20619800](https://pubmed.ncbi.nlm.nih.gov/20619800/)
- WHO (2006). Evaluation of certain food contaminants. Sixty-fourth report of the Joint FAO/WHO Expert

- Committee on Food Additives. WHO Technical Report Series No. 930. Geneva, Switzerland: World Health Organization.
- WHO (2013). GEMS/Food cluster diets - 2012. A part of the FOSCOLLAB platform for food safety data and information. Geneva, Switzerland: World Health Organization. Available from: [https://extranet.who.int/sree/Reports?op=vs&path=/WHO\\_HQ\\_Reports/G7/PROD/EXT/GEMS\\_cluster\\_diets\\_2012&userid=G7\\_ro&password=inetsoft123](https://extranet.who.int/sree/Reports?op=vs&path=/WHO_HQ_Reports/G7/PROD/EXT/GEMS_cluster_diets_2012&userid=G7_ro&password=inetsoft123)
- WHO (2015a). WHO databases on food safety [online databases]. Geneva, Switzerland: World Health Organization. Available from: <http://www.who.int/foodsafety/databases/en/>, accessed 9 July 2015
- WHO (2015b). GEMS/Food contaminants [online database]. Geneva, Switzerland: Global Environment Monitoring System - Food Contamination Monitoring and Assessment Programme, World Health Organization. Available from: <https://extranet.who.int/gemsfood/Search.aspx>
- Williams P (2007). Nutritional composition of red meat. *Nutr Diet*, 64(s4): S11311–9. doi:[10.1111/j.1747-0080.2007.00197.x](https://doi.org/10.1111/j.1747-0080.2007.00197.x)
- Wood JD, Enser M (1997). Factors influencing fatty acids in meat and the role of antioxidants in improving meat quality. *Br J Nutr*, 78(01):Suppl 1: S49–60. doi:[10.1079/BJN19970134](https://doi.org/10.1079/BJN19970134) PMID:[9292774](https://pubmed.ncbi.nlm.nih.gov/9292774/)
- Wood JD, Enser M, Fisher AV, Nute GR, Richardson RI, Sheard PR (1999). Manipulating meat quality and composition. *Proc Nutr Soc*, 58(2):363–70. doi:[10.1017/S0029665199000488](https://doi.org/10.1017/S0029665199000488) PMID:[10466178](https://pubmed.ncbi.nlm.nih.gov/10466178/)
- Wu K, Sinha R, Holmes MD, Giovannucci E, Willett W, Cho E (2010). Meat mutagens and breast cancer in postmenopausal women – a cohort analysis. *Cancer Epidemiol Biomarkers Prev*, 19(5):1301–10. doi:[10.1158/1055-9965.EPI-10-0002](https://doi.org/10.1158/1055-9965.EPI-10-0002) PMID:[20447922](https://pubmed.ncbi.nlm.nih.gov/20447922/)
- Yaylayan VA, Locas CP, Wnorowski A, O'Brien J (2004). The role of creatine in the generation of N-methylacrylamide: a new toxicant in cooked meat. *J Agric Food Chem*, 52(17):5559–65. doi:[10.1021/jf049421g](https://doi.org/10.1021/jf049421g) PMID:[15315400](https://pubmed.ncbi.nlm.nih.gov/15315400/)
- Zeitoun MM, Ahmed SM (2011). Effect of cooking method on the residues of natural sex steroid hormones in local and imported meats and meat products in Al-Qassim region. *J Agril Vet Sci*, 4(2):83–92.
- Zetlaoui M, Feinberg M, Verger P, Cléménçon S (2011). Extraction of food consumption systems by nonnegative matrix factorization (NMF) for the assessment of food choices. *Biometrics*, 67(4):1647–58. doi:[10.1111/j.1541-0420.2011.01588.x](https://doi.org/10.1111/j.1541-0420.2011.01588.x) PMID:[21418050](https://pubmed.ncbi.nlm.nih.gov/21418050/)
- Zhai F, Wang H, Du S, He Y, Wang Z, Ge K et al. (2009). Prospective study on nutrition transition in China. *Nutr Rev*, 67:Suppl 1: S56–61. doi:[10.1111/j.1753-4887.2009.00160.x](https://doi.org/10.1111/j.1753-4887.2009.00160.x) PMID:[19453679](https://pubmed.ncbi.nlm.nih.gov/19453679/)
- Zimmerli B, Rhyh P, Zoller O, Schlatter J (2001). Occurrence of heterocyclic aromatic amines in the Swiss diet: analytical method, exposure estimation and risk assessment. *Food Addit Contam*, 18(6):533–51. doi:[10.1080/02652030119545](https://doi.org/10.1080/02652030119545) PMID:[11407752](https://pubmed.ncbi.nlm.nih.gov/11407752/)
- Zukál E, Incze K (2010). Drying. Part 1, Chapter 11. In: Toldrá F, editor. Handbook of meat processing. Oxford, UK: Wiley-Blackwell; 219–229. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/9780813820897.ch11/summary>