

Betel Quid and Areca Nut

References to Supplementary Web Tables, Section 2

- Ahmad MS, Ali SA, Ali AS, Chaubey KK (2006). Epidemiological and etiological study of oral submucous fibrosis among gutkha chewers of Patna, Bihar, India. *J Indian Soc Pedod Prev Dent*, 24:84–89. [PMID:16823233](#)
- Anwar S, Williams SA, Scott-Smith J *et al.* (2005). A comparison of attitudes and practices of *gutka* users and non-users in Chitrakoot, India. A pilot. *Prim Dent Care*, 12:5–10. [PMID:15703153](#)
- Ariyawardana A, Athukorala AD, Arulanandam A (2006). Effect of betel chewing, tobacco smoking and alcohol consumption on oral submucous fibrosis: a case-control study in Sri Lanka. *J Oral Pathol Med*, 35:197–201. [PMID:16519765](#)
- Ariyawardana A, Sitheequ MA, Ranasinghe AW *et al.* (2007). Prevalence of oral cancer and pre-cancer and associated risk factors among tea estate workers in the central Sri Lanka. *J Oral Pathol Med*, 36:581–587. [PMID:17944750](#)
- Balaram P, Sridhar H, Rajkumar T *et al.* (2002). Oral cancer in southern India: the influence of smoking, drinking, paan-chewing and oral hygiene. *Int J Cancer*, 98:440–445. [PMID:11920597](#)
- Boonyaphiphat P, Thongsuksai P, Sriplung H, Puttawibul P (2002). Lifestyle habits and genetic susceptibility and the risk of esophageal cancer in the Thai population. *Cancer Lett*, 186:193–199. [doi:10.1016/S0304-3835\(02\)00354-3](#) [PMID:12213289](#)
- Chang HC, Kung YY, Hsieh CF *et al.* (2008). Biological risk factors relevant to chronic disease in three ethnic groups in Taiwan: results from Li-Shin Outreaching Neighborhood Screening (LIONS A1). *Ethn Dis*, 18:228–234. [PMID:18507279](#)
- Chang HY, Chiou CJ, Lin MC *et al.* (2005). A population study of the self-care behaviors and their associated factors of diabetes in Taiwan: results from the 2001 National Health Interview Survey in Taiwan. *Prev Med*, 40:344–348. [doi:10.1016/j.ypmed.2004.06.012](#) [PMID:15533549](#)
- Changrani J, Gany FM, Cruz G *et al.* (2006). Paan and *Gutka* Use in the United States: A Pilot Study in Bangladeshi and Indian-Gujarati Immigrants in New York City. *J Immigr Refug Stud*, 4:99–110. [PMID:17492057](#)
- Chatrchaiwiwatana S (2007). Factors affecting tooth loss among rural Khon Kaen adults: analysis of two data sets. *Public Health*, 121:106–112. [PMID:17005217](#)
- Chaturvedi P, Chaturvedi U, Sanyal B (2002). Prevalence of tobacco consumption in schoolchildren in rural India—an epidemic of tobaccogenic cancers looming ahead in the Third World. *J Cancer Educ*, 17:6. [PMID:12000110](#)
- Chaudhry K, Prabhakar AK, Prabhakaran PS *et al.* (2001) *Prevalence of Tobacco Use in Karnataka and Uttar Pradesh in India.*, New Delhi: Indian Council of Medical Research and WHO South East Asian Regional Office.
- Chelleng PK, Narain K, Das HK *et al.* (2000). Risk factors for cancer nasopharynx: a case-control study from Nagaland, India. *Natl Med J India*, 13:6–8. [PMID:10743368](#)
- Chen PC, Kuo C, Pan CC, Chou MY (2002). Risk of oral cancer associated with human papillomavirus infection, betel quid chewing, and cigarette smoking in Taiwan—an integrated molecular and epidemiological study of 58 cases. *J Oral Pathol Med*, 31:317–322. [PMID:12190813](#)
- Chen PC, Pan CC, Kuo C, Lin CP (2006). Risk of oral nonmalignant lesions associated with human papillomavirus infection, betel quid chewing, and cigarette smoking in Taiwan: an integrated molecular and epidemiologic study. *Arch Pathol Lab Med*, 130:57–61. [PMID:16390239](#)
- Chitra S, Ashok L, Anand L *et al.* (2004). Risk factors for esophageal cancer in Coimbatore, southern India: a hospital-based case-control study. *Indian J Gastroenterol*, 23:19–21. [PMID:15106710](#)
- Chou FH, Yang YH, Kuo SH *et al.* (2009). Relationships among smoking, drinking, betel quid chewing and pregnancy-related nausea and vomiting in Taiwanese aboriginal women. *Kaohsiung J Med Sci*, 25:62–69. [PMID:19321408](#)
- Chu NF, Wu DM, Shen MH, Lin YS (2006). Prevalence of adverse behaviors among young military conscripts in Taiwan. *Mil Med*, 171:301–305. [PMID:16673743](#)

- Chung CH, Yang YH, Wang TY *et al.* (2005). Oral precancerous disorders associated with areca quid chewing, smoking, and alcohol drinking in southern Taiwan. *J Oral Pathol Med*, 34:460–466. [PMID:16091112](#)
- Dikshit RP, Kanhere S (2000). Tobacco habits and risk of lung, oropharyngeal and oral cavity cancer: a population-based case-control study in Bhopal, India. *Int J Epidemiol*, 29:609–614. [doi:10.1093/ije/29.4.609](#) [PMID:10922335](#)
- Dongre A, Deshmukh P, Murali N, Garg B (2008). Tobacco consumption among adolescents in rural Wardha: where and how tobacco control should focus its attention? *Indian J Cancer*, 45:100–106. [PMID:19018113](#)
- Eswar N (2002). Oral health status among the tobacco and betel nut chewers in the Kishore Ganj District of Bangladesh. A statistical study. *Indian J Dent Res*, 13:167–171. [PMID:12765097](#)
- Gajalakshmi V, Asma S, Warren CW (2004). Tobacco survey among youth in South India. *Asian Pac J Cancer Prev*, 5:273–278. [PMID:15373706](#)
- Gajalakshmi V, Peto R, Kanimozhi VC *et al.* (2007). Cohort Profile: the Chennai prospective study of mortality among 500,000 adults in Tamil Nadu, South India. *Int J Epidemiol*, 36:1190–1195. [doi:10.1093/ije/dym091](#) [PMID:17510070](#)
- Hildesheim A, Anderson LM, Chen CJ *et al.* (1997). CYP2E1 genetic polymorphisms and risk of nasopharyngeal carcinoma in Taiwan. *J Natl Cancer Inst*, 89:1207–1212. [doi:10.1093/jnci/89.16.1207](#) [PMID:9274915](#)
- Huang HL, Lee CH, Yen YY *et al.* (2009). School-level contextual factors associated with betel quid chewing among schoolchildren in Taiwan. *Community Dent Oral Epidemiol*, 37:58–67. [PMID:19191820](#)
- IARC (2004). Betel-quid and areca-nut chewing and some areca-nut derived nitrosamines. *IARC Monogr Eval Carcinog Risks Hum*, 85:1–334. [PMID:15635762](#)
- Jacob BJ, Straif K, Thomas G *et al.* (2004). Betel quid without tobacco as a risk factor for oral precancers. *Oral Oncol*, 40:697–704. [PMID:15172639](#)
- Jayakody AA, Viner RM, Haines MM *et al.* (2006). Illicit and traditional drug use among ethnic minority adolescents in East London. *Public Health*, 120:329–338. [PMID:16543028](#)
- Jayalekshmi PA, Gangadharan P, Akiba S *et al.* (2009). Tobacco chewing and female oral cavity cancer risk in Karunagappally cohort, India. *Br J Cancer*, 100:848–852. [doi:10.1038/sj.bjc.6604907](#) [PMID:19259098](#)
- Kaduri P, Kitua H, Mbatia J *et al.* (2008). Smokeless tobacco use among adolescents in Ilala Municipality, Tanzania. *Tanzania J Health Res*, 10:28–33. [PMID:18680962](#)
- Kapil U, Goindi G, Singh V *et al.* (2005a). Consumption of tobacco, alcohol and betel leaf amongst school children in Delhi. *Indian J Pediatr*, 72:993. [PMID:16391460](#)
- Kapil U, Singh P, Bahadur S *et al.* (2005b). Assessment of risk factors in laryngeal cancer in India: a case-control study. *Asian Pac J Cancer Prev*, 6:202–207. [PMID:16101334](#)
- Kaur S, Singh S (2002). Cause for concern in Punjab villages. High levels of Gutka intake among students. *Lifeline*, 7:3–4.
- Ko YC, Huang YL, Lee CH *et al.* (1995). Betel quid chewing, cigarette smoking and alcohol consumption related to oral cancer in Taiwan. *J Oral Pathol Med*, 24:450–453. [PMID:8600280](#)
- Lai CS, Shieh TY, Yang YH *et al.* (2006). Factors associated with quitting areca (betel) quid chewing. *Community Dent Oral Epidemiol*, 34:467–474. [doi:10.1111/j.1600-0528.2006.00305.x](#) [PMID:17092276](#)
- Lan TY, Chang WC, Tsai YJ *et al.* (2007). Areca nut chewing and mortality in an elderly cohort study. *Am J Epidemiol*, 165:677–683. [doi:10.1093/aje/kwk056](#) [PMID:17204513](#)
- Lee CH, Lee JM, Wu DC *et al.* (2005b). Independent and combined effects of alcohol intake, tobacco smoking and betel quid chewing on the risk of esophageal cancer in Taiwan. *Int J Cancer*, 113:475–482. [doi:10.1002/ijc.20619](#) [PMID:15455377](#)
- Lee CH, Wu DC, Lee JM *et al.* (2007). Anatomical subsite discrepancy in relation to the impact of the consumption of alcohol, tobacco and betel quid on esophageal cancer. *Int J Cancer*, 120:1755–1762. [doi:10.1002/ijc.22324](#) [PMID:17230518](#)

- Lee KW, Kuo WR, Tsai SM *et al.* (2005a). Different impact from betel quid, alcohol and cigarette: risk factors for pharyngeal and laryngeal cancer. *Int J Cancer*, 117:831–836. [doi:10.1002/ijc.21237](https://doi.org/10.1002/ijc.21237) [PMID:15957167](https://pubmed.ncbi.nlm.nih.gov/15957167/)
- Lin CF, Wang JD, Chen PH *et al.* (2006). Predictors of betel quid chewing behavior and cessation patterns in Taiwan aborigines. *BMC Public Health*, 6:271. [doi:10.1186/1471-2458-6-271](https://doi.org/10.1186/1471-2458-6-271) [PMID:17081309](https://pubmed.ncbi.nlm.nih.gov/17081309/)
- Lin WY, Pi-Sunyer FX, Liu CS *et al.* (2009). Betel nut chewing is strongly associated with general and central obesity in Chinese male middle-aged adults. *Obesity (Silver Spring)*, 17:1247–1254. [PMID:19247275](https://pubmed.ncbi.nlm.nih.gov/19247275/)
- Lin YS, Chu NF, Wu DM, Shen MH (2004). Prevalence and factors associated with the consumption of betel-nut among military conscripts in Taiwan. *Eur J Epidemiol*, 19:343–351. [doi:10.1023/B:EJEP.0000024695.87008.ff](https://doi.org/10.1023/B:EJEP.0000024695.87008.ff) [PMID:15180105](https://pubmed.ncbi.nlm.nih.gov/15180105/)
- Lu CT, Yen YY, Ho CS *et al.* (1996). A case-control study of oral cancer in Changhua County, Taiwan. *J Oral Pathol Med*, 25:245–248. [PMID:8835822](https://pubmed.ncbi.nlm.nih.gov/8835822/)
- Mazahir S, Malik R, Maqsood M *et al.* (2006). Socio-demographic correlates of betel, areca and smokeless tobacco use as a high risk behavior for head and neck cancers in a squatter settlement of Karachi, Pakistan. *Subst Abuse Treat Prev Policy*, 1:10. [PMID:16722535](https://pubmed.ncbi.nlm.nih.gov/16722535/)
- Merchant A, Husain SS, Hosain M *et al.* (2000). Paan without tobacco: an independent risk factor for oral cancer. *Int J Cancer*, 86:128–131. [PMID:10728606](https://pubmed.ncbi.nlm.nih.gov/10728606/)
- Muwonge R, Ramadas K, Sankila R *et al.* (2008). Role of tobacco smoking, chewing and alcohol drinking in the risk of oral cancer in Trivandrum, India: a nested case-control design using incident cancer cases. *Oral Oncol*, 44:446–454. [PMID:17933578](https://pubmed.ncbi.nlm.nih.gov/17933578/)
- Nandakumar A, Thimmasetty KT, Sreeramareddy NM *et al.* (1990). A population-based case-control investigation on cancers of the oral cavity in Bangalore, India. *Br J Cancer*, 62:847–851. [PMID:2245179](https://pubmed.ncbi.nlm.nih.gov/2245179/)
- Nichter M, Nichter M, Van Sickle D (2004). Popular perceptions of tobacco products and patterns of use among male college students in India. *Soc Sci Med*, 59:415–431. [PMID:15110430](https://pubmed.ncbi.nlm.nih.gov/15110430/)
- Nisar N, Qadri MH, Fatima K, Perveen S (2007). A community based study about knowledge and practices regarding tobacco consumption and passive smoking in Gadap Town, Karachi. *J Pak Med Assoc*, 57:186–188. [PMID:17489526](https://pubmed.ncbi.nlm.nih.gov/17489526/)
- Núñez-de la Mora A, Jesmin F, Bentley GR (2007). Betel nut use among first and second generation Bangladeshi women in London, UK. *J Immigr Minor Health*, 9:299–306. [PMID:17431783](https://pubmed.ncbi.nlm.nih.gov/17431783/)
- Oakley E, Demaine L, Warnakulasuriya S (2005). Areca (betel) nut chewing habit among high-school children in the Commonwealth of the Northern Mariana Islands (Micronesia). *Bull World Health Organ*, 83:656–660. [PMID:16211156](https://pubmed.ncbi.nlm.nih.gov/16211156/)
- Paudel D (2003). Tobacco use among adolescent students in secondary schools of Pokhara - Summary Report. Thesis for Master's Degree in Public Health, Institute of Medicine Tribhuvan University, Nepal.
- Rajan G, Ramesh S, Sankaralingam S (2007). Areca nut use in rural Tamil Nadu: a growing threat. *Indian J Med Sci*, 61:332–337. [doi:10.4103/0019-5359.32680](https://doi.org/10.4103/0019-5359.32680) [PMID:17558096](https://pubmed.ncbi.nlm.nih.gov/17558096/)
- Rao DN, Desai PB (1998). Risk assessment of tobacco, alcohol and diet in cancers of base tongue and oral tongue—a case control study. *Indian J Cancer*, 35:65–72. [PMID:9849026](https://pubmed.ncbi.nlm.nih.gov/9849026/)
- Rao DN, Ganesh B, Rao RS, Desai PB (1994). Risk assessment of tobacco, alcohol and diet in oral cancer—a case-control study. *Int J Cancer*, 58:469–473. [PMID:8056441](https://pubmed.ncbi.nlm.nih.gov/8056441/)
- Rozi S, Akhtar S (2007). Prevalence and predictors of smokeless tobacco use among high-school males in Karachi, Pakistan. *East Mediterr Health J*, 13:916–924. [PMID:17955775](https://pubmed.ncbi.nlm.nih.gov/17955775/)
- Sankaranarayanan R, Duffy SW, Day NE *et al.* (1989b). A case-control investigation of cancer of the oral tongue and the floor of the mouth in southern India. *Int J Cancer*, 44:617–621. [PMID:2793234](https://pubmed.ncbi.nlm.nih.gov/2793234/)
- Sankaranarayanan R, Duffy SW, Padmakumary G *et al.* (1990a). Risk factors for cancer of the buccal and labial mucosa in Kerala, southern India. *J Epidemiol Community Health*, 44:286–292. [PMID:2277249](https://pubmed.ncbi.nlm.nih.gov/2277249/)
- Sankaranarayanan R, Duffy SW, Nair MK *et al.* (1990b). Tobacco and alcohol as risk factors in cancer of the larynx in Kerala, India. *Int J Cancer*, 45:879–882. [doi:10.1002/ijc.2910450517](https://doi.org/10.1002/ijc.2910450517) [PMID:2335391](https://pubmed.ncbi.nlm.nih.gov/2335391/)
- Sankaranarayanan R, Duffy SW, Padmakumary G *et al.* (1989a). Tobacco chewing, alcohol and nasal snuff in cancer of the gingiva in Kerala, India. *Br J Cancer*, 60:638–643. [PMID:2803939](https://pubmed.ncbi.nlm.nih.gov/2803939/)

- Sapkota A, Gajalakshmi V, Jetly DH *et al.* (2007). Smokeless tobacco and increased risk of hypopharyngeal and laryngeal cancers: a multicentric case-control study from India. *Int J Cancer*, 121:1793–1798. [doi:10.1002/ijc.22832](https://doi.org/10.1002/ijc.22832) PMID:17583577
- Shah SM, Merchant AT, Luby SP, Chotani RA (2002). Addicted schoolchildren: prevalence and characteristics of areca nut chewers among primary school children in Karachi, Pakistan. *J Paediatr Child Health*, 38:507–510. [doi:10.1046/j.1440-1754.2002.00040.x](https://doi.org/10.1046/j.1440-1754.2002.00040.x) PMID:12354270
- Sinha DN, Gupta PC (2005). Tobacco use among students in Orissa and Uttar Pradesh. *Indian Pediatr*, 42:846–847. PMID:16141500
- Sinha DN, Gupta PC, Pednekar MS (2003). Tobacco use among school personnel in eight North-eastern states of India. *Indian J Cancer*, 40:3–14. PMID:14716126
- Srivatanakul P, Parkin DM, Khlat M *et al.* (1991). Liver cancer in Thailand. II. A case-control study of hepatocellular carcinoma. *Int J Cancer*, 48:329–332. [doi:10.1002/ijc.2910480303](https://doi.org/10.1002/ijc.2910480303) PMID:1645698
- Subapriya R, Thangavelu A, Mathavan B *et al.* (2007). Assessment of risk factors for oral squamous cell carcinoma in Chidambaram, Southern India: a case-control study. *Eur J Cancer Prev*, 16:251–256. PMID:17415096
- Sun CA, Wu DM, Lin CC *et al.* (2003). Incidence and cofactors of hepatitis C virus-related hepatocellular carcinoma: a prospective study of 12,008 men in Taiwan. *Am J Epidemiol*, 157:674–682. [doi:10.1093/aje/kwg041](https://doi.org/10.1093/aje/kwg041) PMID:12697571
- Sushma C, Sharang C (2005). Pan masala advertisements are surrogate for tobacco products. *Indian J Cancer*, 42:94–98. [doi:10.4103/0019-509X.16699](https://doi.org/10.4103/0019-509X.16699) PMID:16141509
- Tanwir F, Altamash M, Gustafsson A (2008). Influence of betel nut chewing, dental care habits and attitudes on perceived oral health among adult Pakistanis. *Oral Health Prev Dent*, 6:89–94. PMID:18637386
- Thomas SJ, Bain CJ, Battistutta D *et al.* (2007). Betel quid not containing tobacco and oral cancer: a report on a case-control study in Papua New Guinea and a meta-analysis of current evidence. *Int J Cancer*, 120:1318–1323. PMID:17163423
- Thomas SJ, Harris R, Ness AR *et al.* (2008). Betel quid not containing tobacco and oral leukoplakia: a report on a cross-sectional study in Papua New Guinea and a meta-analysis of current evidence. *Int J Cancer*, 123:1871–1876. PMID:18688850
- Tovosia S, Chen PH, Ko AM *et al.* (2007). Prevalence and associated factors of betel quid use in the Solomon Islands: a hyperendemic area for oral and pharyngeal cancer. *Am J Trop Med Hyg*, 77:586–590. PMID:17827384
- Tsai JF, Chuang LY, Jeng JE *et al.* (2001). Betel quid chewing as a risk factor for hepatocellular carcinoma: a case-control study. *Br J Cancer*, 84:709–713. [doi:10.1054/bjoc.1999.1597](https://doi.org/10.1054/bjoc.1999.1597) PMID:11237396
- Tsai JF, Jeng JE, Chuang LY *et al.* (2004). Habitual betel quid chewing and risk for hepatocellular carcinoma complicating cirrhosis. *Medicine (Baltimore)*, 83:176–187. [doi:10.1097/01.md.0000126971.80227.a4](https://doi.org/10.1097/01.md.0000126971.80227.a4) PMID:15118544
- Tsai YF, Wong TK, Chen SC (2002). Prevalence and related risk factors of areca quid chewing among junior high students in eastern Taiwan. *Public Health*, 116:190–194. PMID:12087476
- Tung TH, Chiu YH, Chen LS *et al.*; Keelung Community-based Integrated Screening programme No. 2 (2004). A population-based study of the association between areca nut chewing and type 2 diabetes mellitus in men (Keelung Community-based Integrated Screening programme No. 2). *Diabetologia*, 47:1776–1781. [doi:10.1007/s00125-004-1532-2](https://doi.org/10.1007/s00125-004-1532-2) PMID:15517150
- Wang LY, You SL, Lu SN *et al.* (2003b). Risk of hepatocellular carcinoma and habits of alcohol drinking, betel quid chewing and cigarette smoking: a cohort of 2416 HBsAg-seropositive and 9421 HBsAg-seronegative male residents in Taiwan. *Cancer Causes Control*, 14:241–250. [doi:10.1023/A:1023636619477](https://doi.org/10.1023/A:1023636619477) PMID:12814203
- Wang SC, Tsai CC, Huang ST, Hong YJ (2003a). Betel nut chewing and related factors in adolescent students in Taiwan. *Public Health*, 117:339–345. PMID:12909424
- Wang SC, Tsai CC, Huang ST, Hong YJ (2004). Betel nut chewing: the prevalence and the intergenerational effect of parental behavior on adolescent students. *J Adolesc Health*, 34:244–249. PMID:14967349

- Wasnik KS, Ughade SN, Zodpey SP, Ingole DL (1998). Tobacco consumption practices and risk of oropharyngeal cancer: a case-control study in Central India. *Southeast Asian J Trop Med Public Health*, 29:827–834. [PMID:10772572](#)
- Wen CP, Tsai SP, Cheng TY *et al.* (2005a). Uncovering the relation between betel quid chewing and cigarette smoking in Taiwan. *Tob Control*, 14 Suppl 1;i16–i22. [doi:10.1136/tc.2004.008003](#) [PMID:15923442](#)
- Wen CP, Cheng TY, Eriksen MP (2005b). How opening the cigarette market led to an increase in betel quid use in Taiwan. *Public Health*, 119:940–944. [PMID:15998528](#)
- Wu IC, Lee CH, Kuo CH *et al.* (2009). Consumption of cigarettes but not betel quid or alcohol increases colorectal cancer risk. *J Formos Med Assoc*, 108:155–163. [doi:10.1016/S0929-6646\(09\)60046-2](#) [PMID:19251551](#)
- Wu IC, Lu CY, Kuo FC *et al.* (2006). Interaction between cigarette, alcohol and betel nut use on esophageal cancer risk in Taiwan. *Eur J Clin Invest*, 36:236–241. [doi:10.1111/j.1365-2362.2006.01621.x](#) [PMID:16620285](#)
- Wu WC, Chang HY, Yen LL, Lee TS (2007). Changes in health risk behaviors of elementary school students in northern Taiwan from 2001 to 2003: results from the child and adolescent behaviors in long-term evolution study. *BMC Public Health*, 7:323. [doi:10.1186/1471-2458-7-323](#) [PMID:17996123](#)
- Yang XR, Diehl S, Pfeiffer R *et al.* Chinese and American Genetic Epidemiology of NPC Study Team (2005). Evaluation of risk factors for nasopharyngeal carcinoma in high-risk nasopharyngeal carcinoma families in Taiwan. *Cancer Epidemiol Biomarkers Prev*, 14:900–905. [doi:10.1158/1055-9965.EPI-04-0680](#) [PMID:15826929](#)
- Yang YH, Chen CH, Chang JS *et al.* (2005a). Incidence rates of oral cancer and oral pre-cancerous lesions in a 6-year follow-up study of a Taiwanese aboriginal community. *J Oral Pathol Med*, 34:596–601. [PMID:16202079](#)
- Yang YH, Lien YC, Ho PS *et al.* (2005b). The effects of chewing areca/betel quid with and without cigarette smoking on oral submucous fibrosis and oral mucosal lesions. *Oral Dis*, 11:88–94. [PMID:15752081](#)
- Yap SF, Ho PS, Kuo HC, Yang YH (2008). Comparing factors affecting commencement and cessation of betel quid chewing behavior in Taiwanese adults. *BMC Public Health*, 8:199. [PMID:18533044](#)
- Yen AM, Chen SC, Chang SH, Chen TH (2008a). The effect of betel quid and cigarette on multistate progression of oral pre-malignancy. *J Oral Pathol Med*, 37:417–422. [PMID:18410311](#)
- Yen CF, Yang MS, Su YC *et al.* (2009). Effect of the number of risk factors on betel quid chewing in adolescents living in rural Taiwan. *Public Health*, 123:66–68. [doi:10.1016/j.puhe.2008.07.008](#) [PMID:19091361](#)
- Yen TT, Lin WD, Wang CP *et al.* (2008b). The association of smoking, alcoholic consumption, betel quid chewing and oral cavity cancer: a cohort study. *Eur Arch Otorhinolaryngol*, 265:1403–1407. [PMID:18389268](#)
- Zhang X, Reichart PA (2007). A review of betel quid chewing, oral cancer and precancer in Mainland China. *Oral Oncol*, 43:424–430. [PMID:17258497](#)
- Znaor A, Brennan P, Gajalakshmi V *et al.* (2003). Independent and combined effects of tobacco smoking, chewing and alcohol drinking on the risk of oral, pharyngeal and esophageal cancers in Indian men. *Int J Cancer*, 105:681–686. [doi:10.1002/ijc.11114](#) [PMID:12740918](#)