

Corrigenda to the *IARC Handbooks of Cancer Prevention* – Volume 17

Section	Table/Figure/Text	Page	Details of Corrigendum	First published in PDF format	Correction made to PDF version	Correction made in printed version?
3.8.2(a)	Text	266	<p>A sentence was corrected, as follows:</p> <p>Original sentence:</p> <p>“Genome-wide association studies (GWAS) <b>have identified</b> up to 37 single-nucleotide polymorphisms (SNPs) with a positive association with CRC, although recent meta-analyses did not confirm the association with CRC for 22 of the SNPs (Montazeri et al., 2016).”</p> <p>Revised sentence:</p> <p>“Genome-wide association studies (GWAS) <b>have so far identified</b> up to 37 single-nucleotide polymorphisms (SNPs) with a positive association with CRC, although recent meta-analyses did not confirm the association with CRC for 22 of the SNPs (Montazeri et al., 2016).”</p>	3 June 2019	Yes, 26 July 2019	Yes
3.8.3(c)(iii)	Text	276	<p>A sentence was corrected, as follows:</p> <p>Original sentence:</p> <p>“The guidelines from Australia <b>recommended</b> a surveillance interval of 3 years after the detection of one to four adenomas 10 mm or larger, and a surveillance interval of 1 year after the detection of five to nine adenomas 10 mm or larger (Barclay et al., 2013).”</p> <p>Revised sentence:</p> <p>“The guidelines from Australia <b>recommend</b> a surveillance interval of 3 years after the detection of one to four adenomas 10 mm or larger, and a surveillance interval of 1 year after the detection of five to nine adenomas 10 mm or larger (Barclay et al., 2013).”</p>	3 June 2019	Yes, 26 July 2019	Yes

Corrigenda to the *IARC Handbooks of Cancer Prevention* – Volume 17

Section	Table/Figure/Text	Page	Details of Corrigendum	First published in PDF format	Correction made to PDF version	Correction made in printed version?
4.9.3	Text	295	<p>A sentence was corrected, as follows:</p> <p>Original sentence:</p> <p>“The magnitude of the risk and the recommended surveillance <b>intervals and the age at which to start screening depend</b> largely on the characteristics of the lesions and the number of lesions, which are used to distinguish between patients with high-risk characteristics and those with low-risk characteristics.”</p> <p>Revised sentence:</p> <p>“The magnitude of the risk and the recommended surveillance <b>intervals depend</b> largely on the characteristics of the lesions and the number of lesions, which are used to distinguish between patients with high-risk characteristics and those with low-risk characteristics.”</p>	3 June 2019	Yes, 26 July 2019	Yes