

OCCUPATIONAL EXPOSURE AS A FIREFIGHTER

VOLUME 132

This publication represents the views and expert opinions of an IARC Working Group on the Identification of Carcinogenic Hazards to Humans, which met in Lyon, France, 7–14 June 2022

LYON, FRANCE - 2023

IARC MONOGRAPHS
ON THE IDENTIFICATION
OF CARCINOGENIC HAZARDS
TO HUMANS

Table S1.22 Measures of compounds other than fire smoke and polycyclic aromatic hydrocarbons in the firefighting setting

Sample type	Type of fire	Location and sampling date	Sampling period and duration	No. of individuals sampled and type of firefighter	No. of samples	Mean ± SD *geometric mean (± GSD or 95% CI) **median (range)	Units	LOD and/or LOQ	Comments and other relevant information	Reference
<i>Asbestos (IARC Group 1)</i>										
Area air	Structure	Arizona, USA, 1998	Overhaul, unknown	NA, career firefighters	46	0.073 ± 0.063	f/cm ³	0.03	15 samples above LOD	Bolstad-Johnson et al. (2000)
Surface (PPE, tunics)	Structure	United Kingdom, 1994	Attack, unknown	2, unknown	6	2.7 (range, 1–4)	f/ft ²	NR	Control measured 0 f/ft ²	Bridgman (2001)
Surface (municipal fire apparatus)	NA	Kentucky, USA, ~2018	NA	NA	29	Actinolite: < 1590 Chrysotile: < 792–1570	f/cm ³	792–1590	Only 1 sample above LOD	Hwang et al. (2019)
Personal air	Wildland (former vermiculite mine)	Montana, USA, 2018	Specific wildland firefighting tasks, 12–326 min	19, wildland firefighters	42	0.0013–0.13	f/cm ³	0.002–0.03	22 samples were overloaded with debris	NIOSH (2019)
<i>Per- and polyfluoroalkyl substances (PFAS)</i>										
PPE (turnout gear) extraction	NA	USA, 2019	NA	Used PPE from municipal firefighters	8	<i>IARC Group 2B</i> PFOA: ND–250 <i>Selected others</i> PFOS: ND–7 PFNA: ND–25.3	ng/g	1.64–7.79	Included used and unused gear, thermal liner, moisture barrier, and outer shell samples	Peaslee et al. (2020)
Surface (fire station dust)	NA	Massachusetts, USA, 2018	NA	15 career municipal fire stations	88	<i>IARC Group 2B</i> PFOA: 4.85 (ND–60)** <i>Selected others</i> PFOS: 4.95 (1.26–91.5)** PFHxS: 1.77 (ND–12.2)** PFNA: 1.59 (0.446–29.8)**	ng/g	0.12–0.41	Gear locker rooms had significantly higher total fluorine concentrations than living areas did	Young et al. (2021)

Table S1.22 (continued)

Sample type	Type of fire	Location and sampling date	Sampling period and duration	No. of individuals sampled and type of firefighter	No. of samples	Mean ± SD *geometric mean (± GSD or 95% CI) **median (range)	Units	LOD and/or LOQ	Comments and other relevant information	Reference
<i>Chemical flame retardants: polybrominated diphenyl ethers (PBDEs) and other brominated flame retardants</i>										
Area air	Controlled residential fires	Illinois, USA, 2015	During fire attack, 4–14 min	NA, sampling from inside structure	12	BDE-47: 3.47 (ND–13.9)** BDE-99: 4.94 (0.03–13.8)** BDE-153: ND BDE-209: 15.6 (0.03–67.7)** TBBPA: 0.24 (ND–18.5)** TBB: 7.71 (ND–25.2)** TBPH: 0.86 (ND–3.65)** DBDPE: 0.18 (ND–1.45)**	µg/m ³	0.06–0.5		Fent et al. (2020)
Area air	Controlled residential fires	Illinois, USA, 2015	During overhaul, ~15 min	NA, sampling from inside structure	11	BDE-47: 0.02 (ND–0.39)** BDE-99: 0.02 (ND–0.12)** BDE-153: 0.02 (ND–0.40)** BDE-209: 1.08 (0.02–29.4)** TBBPA: ND TBB: 0.03 (ND–26.5)** TBPH: 0.02 (ND–7.01)** DBDPE: ND	µg/m ³	0.04–0.06		Fent et al. (2020)

Table S1.22 (continued)

Sample type	Type of fire	Location and sampling date	Sampling period and duration	No. of individuals sampled and type of firefighter	No. of samples	Mean ± SD *geometric mean (± GSD or 95% CI) **median (range)	Units	LOD and/or LOQ	Comments and other relevant information	Reference
Wipe (PPE: gloves)	Controlled residential fires	Illinois, USA, 2015	After firefighting	7 right-handed gloves worn by firefighters	7	BDE-47: 2.36 (0.27–5.06)** BDE-99: 3.34 (0.32–7.59)** BDE-100: 0.05 (ND–1.70)** BDE-153: 2.68 (ND–4.02)** BDE-209: 74.8 (9.06–148)** TBBPA: 0.05 (ND–0.53)** TBB: 10.2 (0.24–296)** TBPH: 4.96 (0.11–116)** DBDPE: 12.5 (2.29–30.5)**	ng/cm ²	0.1		Fent et al. (2020)
Wipe (PPE: cuffs from turnout jackets and pants)	NA	Australia, ~2019	Wipes collected after laundering	NA	69	BDE-47: 0.051 ± 0.046 BDE-99: 0.098 ± 0.099 BDE-100: 0.018 ± 0.017 BDE-153: 0.012 ± 0.016	ng/cm ²	NR	Also sampled brand new gear, and all PBDEs were ND. This study also found little change in PBDE contamination after laundering	Banks et al. (2021b)
PPE (turnout jacket outer shell) extraction	NA	Pennsylvania, USA, 2013	NA	7 sets of turnout gear from municipal fire department	28	BDE-47: 113 ± 309 BDE-99: 155 ± 425 BDE-100: 32 ± 105 BDE-153: 16 ± 41	mg/kg	NR		Easter et al. (2016)

Table S1.22 (continued)

Sample type	Type of fire	Location and sampling date	Sampling period and duration	No. of individuals sampled and type of firefighter	No. of samples	Mean \pm SD *geometric mean (\pm GSD or 95% CI) **median (range)	Units	LOD and/or LOQ	Comments and other relevant information	Reference
PPE (hood) extraction	Controlled residential fires	Illinois, USA, 2015	After firefighting	NA, 4 knit hoods worn by firefighters	4	BDE-47: 9.98 \pm 2.82 BDE-99: 21.3 \pm 3.20 BDE-153: 7.22 \pm 5.8 BDE-209: 1010 \pm 952 TBBPA: 18.7 \pm 20.1 TBB: 34.2 \pm 25.9 TBPH: 13.9 \pm 9.48 DBDPE: 93.1 \pm 60.5	ng/g	2	Samples also collected after laundering (not shown). Many of the compounds were not effectively removed via laundering	Mayer et al. (2019)
Vacuum cleaner dust (fire station)	NA	California, Minnesota, New Hampshire, New York, Texas, USA, 2015	NA	NA, 26 fire stations	26	BDE-28: 24.1 (5.1–1000)** BDE-47: 3050 (404–161 000)** BDE-99: 4180 (465–338 000)** BDE-100: 756 (87.9–82 000)** BDE-153: 489 (73–29 400)** BDE-209: 57 000 (1990–351 000)**	ng/g	0.14–74	BDE-209 levels in fire stations were among the highest measured in any occupation setting. Similar levels as reported in Shen et al. (2015)	Shen et al. (2018)
Vacuum cleaner dust (fire station)	NA	British Columbia, Ontario, Alberta, Manitoba, Canada, 2017–2018	NA	NA, 19 fire stations	19	BDE-28: 2.4 (ND–469)** BDE-47: 174 (3.0–42 200)** BDE-99: 326 (7.1–123 000)** BDE-100: 62 (1.6–19 700)** BDE-153: 32 (ND–15 200)** BDE-209: 7060 (232–56 300)**	ng/g	NR		Gill et al. (2020)

Table S1.22 (continued)

Sample type	Type of fire	Location and sampling date	Sampling period and duration	No. of individuals sampled and type of firefighter	No. of samples	Mean ± SD *geometric mean (± GSD or 95% CI) **median (range)	Units	LOD and/or LOQ	Comments and other relevant information	Reference
<i>Chemical flame retardants: organophosphate flame retardants (OPFRs)</i>										
Area air	Controlled residential fires	Illinois, USA, 2015	During fire attack, 4–14 min	NA, sampling from inside structure	12	TCPP: 0.12 (ND–4.04)** TDCPP: 0.13 (ND–113)** TPP: 408 (ND–2110)** TCP: 0.04 (ND–897)**	µg/m ³	0.02–0.7		Fent et al. (2020)
Area air	Controlled residential fires	Illinois, USA, 2015	During overhaul, ~15 min	NA, sampling from inside structure	11	TCPP: ND TDCPP: ND TPP: 2.81 (1.07–45.8)** TCP: 0.01 (ND–13.9)**	µg/m ³	0.008–0.012		Fent et al. (2020)
Wipe (PPE: gloves)	Controlled residential fires	Illinois, USA, 2015	After firefighting	7 right-handed gloves worn by firefighters	7	TCPP: 0.78 (ND–13.6)** TDCPP: 7.54 (ND–38.8)** TPP: 20.0 (4.79–93.5)** TCP: 0.88 (0.55–1.28)**	ng/cm ²	0.1–1.5		Fent et al. (2020)
Wipe (PPE: cuffs from turnout jackets and pants)	NA	Australia, ~2019	Wipes collected after laundering	NA	69	TDCPP: 4.40 ± 2.30 TPP: 0.87 ± 0.29 TBOEP: 0.15 ± 0.29 TEHP: 0.11 ± 0.10 TMPP: 0.034 ± 0.022 EHDPP: 0.072 ± 0.042	ng/cm ²	NR	Also sampled brand new gear, and TPP and EHDPP were detected. This study also found little change in OPFR contamination after laundering	Banks et al. (2021b)
PPE (hood) extraction	Controlled residential fires	Illinois, USA, 2015	After firefighting	NA, 4 knit hoods worn by firefighters	4	TCPP: 2.15 ± 1.25 TDCPP: 72.0 ± 63.8 TPP: 294 ± 259 TCP: 71.9 ± 39.1	ng/g	2	Samples also collected after laundering (not shown), which showed a reduction of 15–98%	Mayer et al. (2019)

Table S1.22 (continued)

Sample type	Type of fire	Location and sampling date	Sampling period and duration	No. of individuals sampled and type of firefighter	No. of samples	Mean ± SD *geometric mean (± GSD or 95% CI) **median (range)	Units	LOD and/or LOQ	Comments and other relevant information	Reference
Vacuum cleaner dust (fire station)	NA	California, Minnesota, New Hampshire, New York, Texas, USA, 2015	NA	NA, 26 fire stations	26	TCEP: 1040 (178–4660)** TCPP: 3880 (499–37 400)** TDCPP: 10 900 (1650–218 000)** TPP: 10 800 (1150–85 400)**	ng/g	0–1240		Shen et al. (2018)
Vacuum cleaner dust (fire station)	NA	British Columbia, Ontario, Alberta, Manitoba, Canada, 2017–2018	NA	NA, 19 fire stations	19	TCEP: 534 (ND–3800)** TCPP: 2350 (1470–10 700)** TDCPP: 3370 (1300–78 600)** TPP: 4780 (1270–26 600)**	ng/g	NR		Gill et al. (2020)
<i>Diesel exhaust (IARC Group 1): elemental carbon</i>										
Area air	NA	USA, 2016	7–8 h	NA, locations in 3 fire stations	28	Living area: 1.2–2.7 Office area: 0.86–1.5 Sleeping quarters: 0.43–1.3 Apparatus bay: 0.91–13	µg/m ³	NR		NIOSH (2016)
Area air	NA	Ontario, Canada, 2016–2017	24 h	NA, locations in 12 fire stations	69	Living quarters: < 0.50–< 0.60 Sleeping quarters: < 0.50–1.3 Apparatus bay: < 0.5–2.7	µg/m ³	0.5	Respirable fraction was measured	Chung et al. (2020)
Area air	NA	Queensland, Australia, ~2016	10 h	NA, locations in 9 fire stations	33	Sleeping quarters: ND–2 Office area: ND to > 6 Apparatus bay: 1–26 Outdoors: ND to > 7	µg/m ³	1	Concentrations were higher at the start of the shift	Bott et al. (2017)

Table S1.22 (continued)

Sample type	Type of fire	Location and sampling date	Sampling period and duration	No. of individuals sampled and type of firefighter	No. of samples	Mean ± SD *geometric mean (± GSD or 95% CI) **median (range)	Units	LOD and/or LOQ	Comments and other relevant information	Reference
<i>Diesel exhaust (IARC Group 1): total particulate matter</i>										
Personal air	NA	Los Angeles, Boston, New York, USA, 1985	8 h	228 municipal firefighters (truck and engine companies)	228	1st shift: 91–748 2nd shift: 100–480 3rd shift: 35–182	µg/m ³	NR	Study also included real-time measurements that showed increases with start and operation of vehicles	Froines et al. (1987)
<i>Heavy metals</i>										
Personal air	Structure fires	Ottawa, Canada, 2015	During fire responses	28 municipal firefighters	29	<i>IARC Group 1</i> Cadmium: 0.04* (0.0002–6.85) <i>Selected others</i> Antimony: 3.41* (0.02–170) Lead: 0.95* (0.0016–665)	µg/m ³	0.0087–0.019	Two air concentrations of lead exceeded the adjusted occupational exposure limit (46.9 µg/m ³)	Keir et al. (2020)
PPE (hood) extraction	NA	Pennsylvania, USA, 2013	NA	10 used knit hoods from municipal fire department	10	<i>IARC Group 1</i> Cadmium: 2.2 ± 1.2 Arsenic: 4.1 ± 6.4 <i>Selected others</i> Antimony: 31 ± 13 Chromium: 37 ± 20 Lead: 283 ± 264 Mercury: 0.2 ± 0.1 Manganese: 25 ± 13	mg/kg	0.05	Other metals than those listed here were detected. Used hoods had higher levels of all metals that were measured compared to the unused control hood	Easter et al. (2016)

Table S1.22 (continued)

Sample type	Type of fire	Location and sampling date	Sampling period and duration	No. of individuals sampled and type of firefighter	No. of samples	Mean ± SD *geometric mean (± GSD or 95% CI) **median (range)	Units	LOD and/or LOQ	Comments and other relevant information	Reference
Wipe (surface)	NA	Australia, NR	NA	Numerous surfaces from 15 fire stations	270 turnout gear surface samples	<i>IARC Group 1</i> Cadmium: < 5 <i>Selected others</i> Chromium: 405 Lead: 456 Manganese: 784 Nickel: 35.9	µg/m ²	1–8	Turnout gear from full time firefighters had higher mean levels of lead, zinc, and manganese compared to part time firefighters	Engelsman et al. (2019)
Wipe (surface)	NA	Australia, NR	NA	Numerous surfaces from 15 fire stations	129 vehicle surface samples	<i>IARC Group 1</i> Cadmium: < 5 <i>Selected others</i> Chromium: 39.5 Lead: 130 Manganese: 166 Nickel: 49.6	µg/m ²	1–8		Engelsman et al. (2019)
Wipe (surface)	NA	Australia, NR	NA	Numerous surfaces from 15 fire stations	60 fire station surface samples	<i>IARC Group 1</i> Cadmium: < 5 <i>Selected others</i> Chromium: 493 Lead: 480 Manganese: 638 Nickel: 2170	µg/m ²	1–8	SCBA wash sink was found to be a major area of contamination for lead, manganese, copper, and chromium	Engelsman et al. (2019)
<i>Polychlorinated dibenzo-para-dioxins/dibenzofurans (PCDD/Fs)</i>										
Wipe (PPE: gloves)	Controlled residential fires	Illinois, USA, 2015	After firefighting	7 left-handed gloves worn by firefighters	7	HpCDD: 0.14 (0.02–0.24)** OCDD: 0.72 (0.08–2.29)** 2378-TCDF: 0.06 (0.03–0.08)**	pg/cm ²	0.04–0.58		Fent et al. (2020)

Table S1.22 (continued)

Sample type	Type of fire	Location and sampling date	Sampling period and duration	No. of individuals sampled and type of firefighter	No. of samples	Mean ± SD *geometric mean (± GSD or 95% CI) **median (range)	Units	LOD and/or LOQ	Comments and other relevant information	Reference
Wipe (surface of helmets)	Following brewery, ironwork, textile, and dwelling fires	Taiwan, China, 2010	After firefighting	NA, wipe samples from 5 helmets	5	$\Sigma 17$ PCDD/Fs Brewery: > 0.35 Ironwork: > 0.15 Textile: > 0.08 Dwelling: > 0.08 Clean helmet: 0.0013	pg/cm ²	NR		Hsu et al. (2011)
<i>Polybrominated dibenzo-para-dioxins/dibenzofurans (PBDD/Fs)</i>										
Wipe (PPE: gloves)	Controlled residential fires	Illinois, USA, 2015	After firefighting	7 left-handed gloves worn by firefighters	7	2378-TBDF: 0.14 (0.06–0.35)** 23 478-PeBDF: 0.12 (0.06–0.31)** 123 478-HxBDF: 0.41 (0.06–2.19)** 1 234 678-HpBDF: 0.74 (0.06–13.3)**	ng/cm ²	0.12	Method used was qualitative and results should be considered exploratory. Other brominated furans were detected, but exact species were unknown	Fent et al. (2020)

BDE, brominated diphenyl ether; CI, confidence interval; DBDPE, decabromodiphenylethane; EHDPP, 2-ethylhexyldiphenyl phosphate; f, fibre; FR, flame retardant; GSD, geometric standard deviation; HpBDF, heptabromodibenzofuran; HxBDF, hexabromodibenzofuran; LOD, limit of detection; LOQ, limit of quantification; NA, not applicable; ND, not detected; NR, not reported; PBDD/Fs, polybrominated dibenzo-*para*-dioxins/dibenzofurans; PBDEs, polybrominated diphenyl ethers; PCDD/Fs, polychlorinated dibenzo-*para*-dioxins/dibenzofurans; PeBDF, 1,2,3,7,8-pentabromodibenzofuran; PFHxS, perfluorohexanesulfonic acid; PFNA, perfluorononanoic acid; PFOA, perfluorooctanoic acid; PFOS, perfluorooctane sulfonate; PPE, personal protective equipment; SD, standard deviation; TBB, 2-ethylhexyl 2,3,4,5-tetrabromobenzoate; TBBPA, tetrabromobisphenol A; TBDF, tetrabromodibenzofuran; TBOEP, tris(2-butoxyethyl) phosphate; TBPH, di(2-ethylhexyl)-2,3,4,5-tetrabromophthalate; TCEP, tris(2-carboxyethyl) phosphate; TCP, tricresyl phosphate; TCPP, tris(2-chloropropyl) phosphate; TDCPP, tris(1,3-dichloro-2-propyl) phosphate; TEHP, tris(2-ethylhexyl) phosphate; TMPP, tris(2-methylphenyl) phosphate; TPP, triphenyl phosphate.

References

- Adetona AM, Adetona O, Gogal RM Jr, Diaz-Sanchez D, Rathbun SL, Naeher LP (2017). Impact of work task-related acute occupational smoke exposures on select proinflammatory immune parameters in wildland firefighters. *J Occup Environ Med.* 59(7):679–90. doi:[10.1097/JOM.0000000000001053](https://doi.org/10.1097/JOM.0000000000001053) PMID:[28692002](https://pubmed.ncbi.nlm.nih.gov/28692002/)
- Adetona AM, Martin WK, Warren SH, Hanley NM, Adetona O, Zhang JJ, et al. (2019). Urinary mutagenicity and other biomarkers of occupational smoke exposure of wildland firefighters and oxidative stress. *Inhal Toxicol.* 31(2):73–87. doi:[10.1080/08958378.2019.1600079](https://doi.org/10.1080/08958378.2019.1600079) PMID:[30985217](https://pubmed.ncbi.nlm.nih.gov/30985217/)
- Adetona O, Dunn K, Hall DB, Achtemeier G, Stock A, Naeher LP (2011). Personal PM_{2.5} exposure among wildland firefighters working at prescribed forest burns in southeastern United States. *J Occup Environ Hyg.* 8(8):503–11. doi:[10.1080/15459624.2011.595257](https://doi.org/10.1080/15459624.2011.595257) PMID:[21762011](https://pubmed.ncbi.nlm.nih.gov/21762011/)
- Adetona O, Simpson CD, Onstad G, Naeher LP (2013a). Exposure of wildland firefighters to carbon monoxide, fine particles, and levoglucosan. *Ann Occup Hyg.* 57(8):979–91. doi:[10.1093/annhyg/met024](https://doi.org/10.1093/annhyg/met024) PMID:[23813888](https://pubmed.ncbi.nlm.nih.gov/23813888/)
- Adetona O, Zhang JJ, Hall DB, Wang JS, Vena JE, Naeher LP (2013b). Occupational exposure to woodsmoke and oxidative stress in wildland firefighters. *Sci Total Environ.* 449:269–75. doi:[10.1016/j.scitotenv.2013.01.075](https://doi.org/10.1016/j.scitotenv.2013.01.075) PMID:[23434577](https://pubmed.ncbi.nlm.nih.gov/23434577/)
- Akhtar US, Keir JL, Matschke D, White P, Blais JM (2016). Occupational exposure to polycyclic aromatic hydrocarbons (PAHs) by firefighters. *Toxicol Lett.* 259:S211. doi:[10.1016/j.toxlet.2016.07.505](https://doi.org/10.1016/j.toxlet.2016.07.505)
- Al-Malki AL (2009). Serum heavy metals and hemoglobin related compounds in Saudi Arabia firefighters. *J Occup Med Toxicol.* 4(1):18. doi:[10.1186/1745-6673-4-18](https://doi.org/10.1186/1745-6673-4-18) PMID:[19583874](https://pubmed.ncbi.nlm.nih.gov/19583874/)
- Alharbi BH, Pasha MJ, Al-Shamsi MAS (2021). Firefighter exposures to organic and inorganic gas emissions in emergency residential and industrial fires. *Sci Total Environ.* 770:145332. doi:[10.1016/j.scitotenv.2021.145332](https://doi.org/10.1016/j.scitotenv.2021.145332) PMID:[33515879](https://pubmed.ncbi.nlm.nih.gov/33515879/)
- Allonneau A, Mercier S, Maurin O, Robardet F, Menguy-Fleuriot A, Luu SC, et al. (2021). Lead contamination among Paris Fire Brigade firefighters who fought the Notre Dame Cathedral fire in Paris. *Int J Hyg Environ Health.* 233:113707. doi:[10.1016/j.ijheh.2021.113707](https://doi.org/10.1016/j.ijheh.2021.113707) PMID:[33631659](https://pubmed.ncbi.nlm.nih.gov/33631659/)
- Andersen MHG, Saber AT, Pedersen PB, Loft S, Hansen ÅM, Koponen IK, et al. (2017). Cardiovascular health effects following exposure of human volunteers during fire extinction exercises. *Environ Health.* 16(1):96. doi:[10.1186/s12940-017-0303-8](https://doi.org/10.1186/s12940-017-0303-8) PMID:[28877717](https://pubmed.ncbi.nlm.nih.gov/28877717/)
- Anthony TR, Joggerst P, James L, Burgess JL, Leonard SS, Shogren ES (2007). Method development study for APR cartridge evaluation in fire overhaul exposures. *Ann Occup Hyg.* 51(8):703–16. doi:[10.1093/annhyg/mem048](https://doi.org/10.1093/annhyg/mem048) PMID:[17989124](https://pubmed.ncbi.nlm.nih.gov/17989124/)
- Austin CC, Wang D, Ecobichon DJ, Dussault G (2001a). Characterization of volatile organic compounds in smoke at experimental fires. *J Toxicol Environ Health A.* 63(3):191–206. doi:[10.1080/15287390151101547](https://doi.org/10.1080/15287390151101547) PMID:[11405415](https://pubmed.ncbi.nlm.nih.gov/11405415/)
- Austin CC, Wang D, Ecobichon DJ, Dussault G (2001b). Characterization of volatile organic compounds in smoke at municipal structural fires. *J Toxicol Environ Health A.* 63(6):437–58. doi:[10.1080/152873901300343470](https://doi.org/10.1080/152873901300343470) PMID:[11482799](https://pubmed.ncbi.nlm.nih.gov/11482799/)
- Australian Government Productivity Commission (2022). Report on Government Services 2022. Emergency services for fire and other events. Available from: <https://www.pc.gov.au/research/ongoing/report-on-government-services/2022/emergency-management>, accessed June 2022.
- Banks APW, Thai P, Engelsman M, Wang X, Osorio AF, Mueller JF (2021a). Characterising the exposure of Australian firefighters to polycyclic aromatic hydrocarbons generated in simulated compartment fires. *Int J Hyg Environ Health.* 231:113637. doi:[10.1016/j.ijheh.2020.113637](https://doi.org/10.1016/j.ijheh.2020.113637) PMID:[33080523](https://pubmed.ncbi.nlm.nih.gov/33080523/)
- Banks APW, Wang X, Engelsman M, He C, Osorio AF, Mueller JF (2021b). Assessing decontamination and laundering processes for the removal of polycyclic aromatic hydrocarbons and flame retardants from firefighting uniforms. *Environ Res.* 194:110616. doi:[10.1016/j.envres.2020.110616](https://doi.org/10.1016/j.envres.2020.110616) PMID:[33321140](https://pubmed.ncbi.nlm.nih.gov/33321140/)
- Barboni T, Cannac M, Pasqualini V, Simeoni A, Leoni E, Chiaramonti N (2010). Volatile and semi-volatile organic compounds in smoke exposure of firefighters during prescribed burning in the Mediterranean region. *Int J Wildland Fire.* 19(5):606. doi:[10.1071/WF08121](https://doi.org/10.1071/WF08121)
- Barnard RJ, Weber JS (1979). Carbon monoxide: a hazard to fire fighters. *Arch Environ Health.* 34(4):255–7. doi:[10.1080/00039896.1979.10667410](https://doi.org/10.1080/00039896.1979.10667410) PMID:[475470](https://pubmed.ncbi.nlm.nih.gov/475470/)
- Baxter CS, Hoffman JD, Knipp MJ, Reponen T, Haynes EN (2014). Exposure of firefighters to particulates and polycyclic aromatic hydrocarbons. *J Occup Environ Hyg.* 11(7):D85–91. doi:[10.1080/15459624.2014.890286](https://doi.org/10.1080/15459624.2014.890286) PMID:[24512044](https://pubmed.ncbi.nlm.nih.gov/24512044/)
- Baxter CS, Ross CS, Fabian T, Borgerson JL, Shawon J, Gandhi PD, et al. (2010). Ultrafine particle exposure during fire suppression—is it an important contributory factor for coronary heart disease in firefighters? *J Occup Environ Med.* 52(8):791–6. doi:[10.1097/JOM.0b013e3181ed2c6e](https://doi.org/10.1097/JOM.0b013e3181ed2c6e) PMID:[20657302](https://pubmed.ncbi.nlm.nih.gov/20657302/)

- Bolstad-Johnson DM, Burgess JL, Crutchfield CD, Storment S, Gerkin R, Wilson JR (2000). Characterization of firefighter exposures during fire overhaul. *AIHAJ*. 61(5):636–41. doi:[10.1080/15298660008984572](https://doi.org/10.1080/15298660008984572) PMID:[11071414](https://pubmed.ncbi.nlm.nih.gov/11071414/)
- Bott RC, Kirk KM, Logan MB, Reid DA (2017). Diesel particulate matter and polycyclic aromatic hydrocarbons in fire stations. *Environ Sci Process Impacts*. 19(10):1320–6. doi:[10.1039/C7EM00291B](https://doi.org/10.1039/C7EM00291B) PMID:[28861557](https://pubmed.ncbi.nlm.nih.gov/28861557/)
- Brandt-Rauf PW, Cosman B, Fallon LF Jr, Tarantini T, Idema C (1989). Health hazards of firefighters: acute pulmonary effects after toxic exposures. *Br J Ind Med*. 46(3):209–11. doi:[10.1136/oem.46.3.209](https://doi.org/10.1136/oem.46.3.209) PMID:[2930733](https://pubmed.ncbi.nlm.nih.gov/2930733/)
- Brandt-Rauf PW, Fallon LF Jr, Tarantini T, Idema C, Andrews L (1988). Health hazards of fire fighters: exposure assessment. *Br J Ind Med*. 45(9):606–12. doi:[10.1136/oem.45.9.606](https://doi.org/10.1136/oem.45.9.606) PMID:[3179235](https://pubmed.ncbi.nlm.nih.gov/3179235/)
- Bridgman S (2001). Community health risk assessment after a fire with asbestos containing fallout. *J Epidemiol Community Health*. 55(12):921–7. doi:[10.1136/jech.55.12.921](https://doi.org/10.1136/jech.55.12.921) PMID:[11707487](https://pubmed.ncbi.nlm.nih.gov/11707487/)
- Burgess JL, Crutchfield CD (1995). Tucson fire fighter exposure to products of combustion: a risk assessment. *Appl Occup Environ Hyg*. 10(1):37–42. doi:[10.1080/1047322X.1995.10387609](https://doi.org/10.1080/1047322X.1995.10387609)
- Burgess JL, Nanson CJ, Bolstad-Johnson DM, Gerkin R, Hysong TA, Lantz RC, et al. (2001). Adverse respiratory effects following overhaul in firefighters. *J Occup Environ Med*. 43(5):467–73. doi:[10.1097/00043764-200105000-00007](https://doi.org/10.1097/00043764-200105000-00007) PMID:[11382182](https://pubmed.ncbi.nlm.nih.gov/11382182/)
- Burgess JL, Nanson CJ, Hysong TA, Gerkin R, Witten ML, Lantz RC (2002). Rapid decline in sputum IL-10 concentration following occupational smoke exposure. *Inhal Toxicol*. 14(2):133–40. doi:[10.1080/089583701753403953](https://doi.org/10.1080/089583701753403953) PMID:[12122576](https://pubmed.ncbi.nlm.nih.gov/12122576/)
- Caban-Martinez AJ, Kropa B, Niemczyk N, Moore KJ, Baum J, Solle NS, et al. (2018). The “warm zone” cases: environmental monitoring immediately outside the fire incident response arena by firefighters. *Saf Health Work*. 9(3):352–5. doi:[10.1016/j.shaw.2017.12.003](https://doi.org/10.1016/j.shaw.2017.12.003) PMID:[30370169](https://pubmed.ncbi.nlm.nih.gov/30370169/)
- Caban-Martinez AJ, Louzado-Feliciano P, Santiago KM, Baum J, Schaefer Solle N, Rivera G, et al. (2020). Objective measurement of carcinogens among Dominican Republic firefighters using silicone-based wristbands. *J Occup Environ Med*. 62(11):e611–5. doi:[10.1097/JOM.0000000000002006](https://doi.org/10.1097/JOM.0000000000002006) PMID:[32826549](https://pubmed.ncbi.nlm.nih.gov/32826549/)
- Carballo Leyenda B, Rodríguez-Marroyo JA, López-Satué J, Avila Ordás C, Pernía Cubillo R, Villa Vicente JG (2010). [Exposure to carbon monoxide in wildland firefighters during wildfires suppression]. *Rev Esp Salud Pública*. 84(6):799–807. [Spanish]doi:[10.1590/S1135-57272010000600010](https://doi.org/10.1590/S1135-57272010000600010) PMID:[21327314](https://pubmed.ncbi.nlm.nih.gov/21327314/)
- Chernyak YI, Shelepchikov AA, Brodsky ES, Grassman JA (2012). PCDD, PCDF, and PCB exposure in current and former firefighters from Eastern Siberia. *Toxicol Lett*. 213(1):9–14. doi:[10.1016/j.toxlet.2011.09.021](https://doi.org/10.1016/j.toxlet.2011.09.021) PMID:[21979175](https://pubmed.ncbi.nlm.nih.gov/21979175/)
- Cherry N, Aklilu YA, Beach J, Britz-McKibbin P, Elbourne R, Galarneau JM, et al. (2019). Urinary 1-hydroxypyrene and skin contamination in firefighters deployed to the Fort McMurray fire. *Ann Work Expo Health*. 63(4):448–58. doi:[10.1093/annweh/wxz006](https://doi.org/10.1093/annweh/wxz006) PMID:[30753267](https://pubmed.ncbi.nlm.nih.gov/30753267/)
- Cherry N, Galarneau JM, Kinniburgh D, Quemerais B, Tiu S, Zhang X (2021a). Exposure and absorption of PAHs in wildland firefighters: a field study with pilot interventions. *Ann Work Expo Health*. 65(2):148–61. doi:[10.1093/annweh/wxaa064](https://doi.org/10.1093/annweh/wxaa064) PMID:[32572446](https://pubmed.ncbi.nlm.nih.gov/32572446/)
- Chung J, Demers PA, Kalenge S, Kirkham TL (2020). Career fire hall exposures to diesel engine exhaust in Ontario, Canada. *J Occup Environ Hyg*. 17(1):38–46. doi:[10.1080/15459624.2019.1691729](https://doi.org/10.1080/15459624.2019.1691729) PMID:[31851590](https://pubmed.ncbi.nlm.nih.gov/31851590/)
- Cone DC, MacMillan DS, Van Gelder C, Brown DJ, Weir SD, Bogucki S (2005). Noninvasive fireground assessment of carboxyhemoglobin levels in firefighters. *Prehosp Emerg Care*. 9(1):8–13. doi:[10.1080/10903120590891912](https://doi.org/10.1080/10903120590891912) PMID:[16036821](https://pubmed.ncbi.nlm.nih.gov/16036821/)
- CTIF (2021). World Fire Statistics. Report No. 26. International Association of Fire and Rescue Services. Available from: https://ctif.org/sites/default/files/2021-06/CTIF_Report26.pdf, accessed June 2022.
- De Vos AJ, Cook A, Devine B, Thompson PJ, Weinstein P (2006). Effect of protective filters on fire fighter respiratory health during simulated bushfire smoke exposure. *Am J Ind Med*. 49(9):740–50. doi:[10.1002/ajim.20369](https://doi.org/10.1002/ajim.20369) PMID:[16847937](https://pubmed.ncbi.nlm.nih.gov/16847937/)
- De Vos AJ, Cook A, Devine B, Thompson PJ, Weinstein P (2009a). Effect of protective filters on fire fighter respiratory health: field validation during prescribed burns. *Am J Ind Med*. 52(1):76–87. doi:[10.1002/ajim.20651](https://doi.org/10.1002/ajim.20651) PMID:[18946878](https://pubmed.ncbi.nlm.nih.gov/18946878/)
- De Vos AJ, Reisen F, Cook A, Devine B, Weinstein P (2009b). Respiratory irritants in Australian bushfire smoke: air toxics sampling in a smoke chamber and during prescribed burns. *Arch Environ Contam Toxicol*. 56(3):380–8. doi:[10.1007/s00244-008-9209-3](https://doi.org/10.1007/s00244-008-9209-3) PMID:[18712497](https://pubmed.ncbi.nlm.nih.gov/18712497/)
- Dietrich J, Yermakov M, Reponen T, Kulkarni P, Qi C, Grinshpun SA (2015). Protection of firefighters against combustion aerosol particles: simulated workplace protection factor of a half-mask respirator (pilot study). *J Occup Environ Hyg*. 12(6):415–20. doi:[10.1080/15459624.2015.1006637](https://doi.org/10.1080/15459624.2015.1006637) PMID:[25625543](https://pubmed.ncbi.nlm.nih.gov/25625543/)
- Dobraca D, Israel L, McNeel S, Voss R, Wang M, Gajek R, et al. (2015). Biomonitoring in California firefighters: metals and perfluorinated chemicals. *J Occup Environ Med*. 57(1):88–97. doi:[10.1097/JOM.0000000000000307](https://doi.org/10.1097/JOM.0000000000000307) PMID:[25563545](https://pubmed.ncbi.nlm.nih.gov/25563545/)

- Dunn KH, Devaux I, Stock A, Naeher LP (2009). Application of end-exhaled breath monitoring to assess carbon monoxide exposures of wildland firefighters at prescribed burns. *Inhal Toxicol.* 21(1):55–61. doi:[10.1080/08958370802207300](https://doi.org/10.1080/08958370802207300) PMID:[18946764](https://pubmed.ncbi.nlm.nih.gov/18946764/)
- Dunn KH, Shulman S, Stock AL, Naeher LP (2013). Personal carbon monoxide exposures among firefighters at prescribed forest burns in the southeastern United States. *Arch Environ Occup Health.* 68(1):55–9. doi:[10.1080/19338244.2011.633126](https://doi.org/10.1080/19338244.2011.633126) PMID:[23298425](https://pubmed.ncbi.nlm.nih.gov/23298425/)
- Easter E, Lander D, Huston T (2016). Risk assessment of soils identified on firefighter turnout gear. *J Occup Environ Hyg.* 13(9):647–57. doi:[10.1080/15459624.2016.1165823](https://doi.org/10.1080/15459624.2016.1165823) PMID:[27027971](https://pubmed.ncbi.nlm.nih.gov/27027971/)
- Edwards R, Johnson M, Dunn KH, Naeher LP (2005). Application of real-time particle sensors to help mitigate exposures of wildland firefighters. *Arch Environ Occup Health.* 60(1):40–3. doi:[10.3200/AEOH.60.1.40-43](https://doi.org/10.3200/AEOH.60.1.40-43) PMID:[16961007](https://pubmed.ncbi.nlm.nih.gov/16961007/)
- Ekpe OD, Sim W, Choi S, Choo G, Oh JE (2021). Assessment of exposure of Korean firefighters to polybrominated diphenyl ethers and polycyclic aromatic hydrocarbons via their measurement in serum and polycyclic aromatic hydrocarbon metabolites in urine. *Environ Sci Technol.* 55(20):14015–25. doi:[10.1021/acs.est.1c02554](https://doi.org/10.1021/acs.est.1c02554) PMID:[34435767](https://pubmed.ncbi.nlm.nih.gov/34435767/)
- Engelsman M, Snoek MF, Banks APW, Cantrell P, Wang X, Toms LM, et al. (2019). Exposure to metals and semivolatile organic compounds in Australian fire stations. *Environ Res.* 179(Pt A):108745. doi:[10.1016/j.envres.2019.108745](https://doi.org/10.1016/j.envres.2019.108745) PMID:[31546131](https://pubmed.ncbi.nlm.nih.gov/31546131/)
- Etzel RA, Ashley DL (1994). Volatile organic compounds in the blood of persons in Kuwait during the oil fires. *Int Arch Occup Environ Health.* 66(2):125–9. doi:[10.1007/BF00383368](https://doi.org/10.1007/BF00383368) PMID:[7806395](https://pubmed.ncbi.nlm.nih.gov/7806395/)
- Evans DE, Fent KW (2015). Ultrafine and respirable particle exposure during vehicle fire suppression. *Environ Sci Process Impacts.* 17(10):1749–59. doi:[10.1039/C5EM00233H](https://doi.org/10.1039/C5EM00233H) PMID:[26308547](https://pubmed.ncbi.nlm.nih.gov/26308547/)
- Fahy R, Evarts B, Stein GP (2021). US fire department profile 2020. National Fire Protection Association. NFPA Research; pp. 1–6. Available from: <https://www.nfpa.org/-/media/Files/News-and-Research/Fire-statistics-and-reports/Emergency-responders/osfdprofile.pdf>, accessed June 2022.
- Fent KW, Evans DE (2011). Assessing the risk to firefighters from chemical vapors and gases during vehicle fire suppression. *J Environ Monit.* 13(3):536–43. doi:[10.1039/c0em00591f](https://doi.org/10.1039/c0em00591f) PMID:[21274476](https://pubmed.ncbi.nlm.nih.gov/21274476/)
- Fent KW, Evans DE, Babik K, Striley C, Bertke S, Kerber S, et al. (2018). Airborne contaminants during controlled residential fires. *J Occup Environ Hyg.* 15(5):399–412. doi:[10.1080/15459624.2018.1445260](https://doi.org/10.1080/15459624.2018.1445260) PMID:[29494297](https://pubmed.ncbi.nlm.nih.gov/29494297/)
- Fent KW, LaGuardia M, Luellen D, McCormick S, Mayer A, Chen IC, et al. (2020). Flame retardants, dioxins, and furans in air and on firefighters' protective ensembles during controlled residential firefighting. *Environ Int.* 140:105756. doi:[10.1016/j.envint.2020.105756](https://doi.org/10.1016/j.envint.2020.105756) PMID:[32388249](https://pubmed.ncbi.nlm.nih.gov/32388249/)
- Fent KW, Mayer A, Bertke S, Kerber S, Smith D, Horn GP (2019). Understanding airborne contaminants produced by different fuel packages during training fires. *J Occup Environ Hyg.* 16(8):532–43. doi:[10.1080/15459624.2019.1617870](https://doi.org/10.1080/15459624.2019.1617870) PMID:[31169466](https://pubmed.ncbi.nlm.nih.gov/31169466/)
- Ferguson MD, Semmens EO, Weiler E, Domitrovich J, French M, Migliaccio C, et al. (2017). Lung function measures following simulated wildland firefighter exposures. *J Occup Environ Hyg.* 14(9):739–48. doi:[10.1080/15459624.2017.1326700](https://doi.org/10.1080/15459624.2017.1326700) PMID:[28609218](https://pubmed.ncbi.nlm.nih.gov/28609218/)
- Fernando S, Shaw L, Shaw D, Gallea M, VandenEnden L, House R, et al. (2016). Evaluation of firefighter exposure to wood smoke during training exercises at burn houses. *Environ Sci Technol.* 50(3):1536–43. doi:[10.1021/acs.est.5b04752](https://doi.org/10.1021/acs.est.5b04752) PMID:[26726952](https://pubmed.ncbi.nlm.nih.gov/26726952/)
- Feunekes FD, Jongeneelen FJ, vd Laan H, Schoonhof FH (1997). Uptake of polycyclic aromatic hydrocarbons among trainers in a fire-fighting training facility. *Am Ind Hyg Assoc J.* 58(1):23–8. doi:[10.1080/15428119791013035](https://doi.org/10.1080/15428119791013035) PMID:[9018833](https://pubmed.ncbi.nlm.nih.gov/9018833/)
- Froines JR, Hinds WC, Duffy RM, Lafuente EJ, Liu WC (1987). Exposure of firefighters to diesel emissions in fire stations. *Am Ind Hyg Assoc J.* 48(3):202–7. doi:[10.1080/15298668791384634](https://doi.org/10.1080/15298668791384634) PMID:[2437785](https://pubmed.ncbi.nlm.nih.gov/2437785/)
- Gaughan DM, Piacitelli CA, Chen BT, Law BF, Virji MA, Edwards NT, et al. (2014c). Exposures and cross-shift lung function declines in wildland firefighters. *J Occup Environ Hyg.* 11(9):591–603. doi:[10.1080/15459624.2014.895372](https://doi.org/10.1080/15459624.2014.895372) PMID:[24568319](https://pubmed.ncbi.nlm.nih.gov/24568319/)
- Gill R, Hurley S, Brown R, Tarrant D, Dhaliwal J, Sarala R, et al. (2020). Polybrominated diphenyl ether and organophosphate flame retardants in Canadian fire station dust. *Chemosphere.* 253:126669. doi:[10.1016/j.chemosphere.2020.126669](https://doi.org/10.1016/j.chemosphere.2020.126669) PMID:[32464780](https://pubmed.ncbi.nlm.nih.gov/32464780/)
- Gold A, Burgess WA, Clougherty EV (1978). Exposure of firefighters to toxic air contaminants. *Am Ind Hyg Assoc J.* 39(7):534–9. doi:[10.1080/0002889778507805](https://doi.org/10.1080/0002889778507805) PMID:[211840](https://pubmed.ncbi.nlm.nih.gov/211840/)
- Graber JM, Black TM, Shah NN, Caban-Martinez AJ, Lu SE, Brancard T, et al. (2021). Prevalence and predictors of per- and polyfluoroalkyl substances (PFAS) serum levels among members of a suburban US volunteer fire department. *Int J Environ Res Public Health.* 18(7):3730. doi:[10.3390/ijerph18073730](https://doi.org/10.3390/ijerph18073730) PMID:[33918459](https://pubmed.ncbi.nlm.nih.gov/33918459/)
- Hejl AM, Adetona O, Diaz-Sanchez D, Carter JD, Commodore AA, Rathbun SL, et al. (2013). Inflammatory effects of woodsmoke exposure among wildland firefighters working at prescribed burns at the Savannah River Site, SC. *J Occup Environ Hyg.* 10(4):173–80. doi:[10.1080/15459624.2012.760064](https://doi.org/10.1080/15459624.2012.760064) PMID:[23363434](https://pubmed.ncbi.nlm.nih.gov/23363434/)

- Henn SA, Butler C, Li J, Sussell A, Hale C, Broyles G, et al. (2019). Carbon monoxide exposures among US wildland firefighters by work, fire, and environmental characteristics and conditions. *J Occup Environ Hyg.* 16(12):793–803. doi:[10.1080/15459624.2019.1670833](https://doi.org/10.1080/15459624.2019.1670833) PMID:[31658425](https://pubmed.ncbi.nlm.nih.gov/31658425/)
- Hill TA, Siedle AR, Perry R (1972). Chemical hazards of a fire-fighting training environment. *Am Ind Hyg Assoc J.* 33(6):423–30. doi:[10.1080/0002889728506675](https://doi.org/10.1080/0002889728506675) PMID:[4651528](https://pubmed.ncbi.nlm.nih.gov/4651528/)
- Hsu JF, Guo HR, Wang HW, Liao CK, Liao PC (2011). An occupational exposure assessment of polychlorinated dibenzo-*p*-dioxin and dibenzofurans in firefighters. *Chemosphere.* 83(10):1353–9. doi:[10.1016/j.chemosphere.2011.02.079](https://doi.org/10.1016/j.chemosphere.2011.02.079) PMID:[21458022](https://pubmed.ncbi.nlm.nih.gov/21458022/)
- Hwang J, Taylor R, Cann C, Norris P, Golla V (2019). Evaluation of accumulated polycyclic aromatic hydrocarbons and asbestiform fibers on firefighter vehicles: pilot study. *Fire Technol.* 55(6):2195–213. doi:[10.1007/s10694-019-00851-7](https://doi.org/10.1007/s10694-019-00851-7)
- Jankovic J, Jones W, Burkhart J, Noonan G (1991). Environmental study of firefighters. *Ann Occup Hyg.* 35(6):581–602. doi:[10.1093/annhyg/35.6.581](https://doi.org/10.1093/annhyg/35.6.581) PMID:[1768008](https://pubmed.ncbi.nlm.nih.gov/1768008/)
- Jayatilaka NK, Restrepo P, Davis Z, Vidal M, Calafat AM, Ospina M (2019). Quantification of 16 urinary biomarkers of exposure to flame retardants, plasticizers, and organophosphate insecticides for biomonitoring studies. *Chemosphere.* 235:481–91. doi:[10.1016/j.chemosphere.2019.06.181](https://doi.org/10.1016/j.chemosphere.2019.06.181) PMID:[31272008](https://pubmed.ncbi.nlm.nih.gov/31272008/)
- Jayatilaka NK, Restrepo P, Williams L, Ospina M, Valentin-Blasini L, Calafat AM (2017). Quantification of three chlorinated dialkyl phosphates, diphenyl phosphate, 2,3,4,5-tetrabromobenzoic acid, and four other organophosphates in human urine by solid phase extraction-high performance liquid chromatography-tandem mass spectrometry. *Anal Bioanal Chem.* 409(5):1323–32. doi:[10.1007/s00216-016-0061-4](https://doi.org/10.1007/s00216-016-0061-4) PMID:[27838756](https://pubmed.ncbi.nlm.nih.gov/27838756/)
- Jin C, Sun Y, Islam A, Qian Y, Ducatman A (2011). Perfluoroalkyl acids including perfluorooctane sulfonate and perfluorohexane sulfonate in firefighters. *J Occup Environ Med.* 53(3):324–8. doi:[10.1097/JOM.0b013e31820d1314](https://doi.org/10.1097/JOM.0b013e31820d1314) PMID:[21346631](https://pubmed.ncbi.nlm.nih.gov/21346631/)
- Kato K, Kalathil AA, Patel AM, Ye X, Calafat AM (2018). Per- and polyfluoroalkyl substances and fluorinated alternatives in urine and serum by on-line solid phase extraction-liquid chromatography-tandem mass spectrometry. *Chemosphere.* 209:338–45. doi:[10.1016/j.chemosphere.2018.06.085](https://doi.org/10.1016/j.chemosphere.2018.06.085) PMID:[29935462](https://pubmed.ncbi.nlm.nih.gov/29935462/)
- Keir JLA, Akhtar US, Matschke DMJ, White PA, Kirkham TL, Chan HM, et al. (2020). Polycyclic aromatic hydrocarbon (PAH) and metal contamination of air and surfaces exposed to combustion emissions during emergency fire suppression: implications for firefighters' exposures. *Sci Total Environ.* 698:134211. doi:[10.1016/j.scitotenv.2019.134211](https://doi.org/10.1016/j.scitotenv.2019.134211) PMID:[31514022](https://pubmed.ncbi.nlm.nih.gov/31514022/)
- Khalil N, Ducatman AM, Sinari S, Billheimer D, Hu C, Littau S, et al. (2020). Per- and polyfluoroalkyl substance and cardio metabolic markers in firefighters. *J Occup Environ Med.* 62(12):1076–81. doi:[10.1097/JOM.0000000000002062](https://doi.org/10.1097/JOM.0000000000002062) PMID:[33105404](https://pubmed.ncbi.nlm.nih.gov/33105404/)
- Kirk KM, Logan MB (2015). Firefighting instructors' exposures to polycyclic aromatic hydrocarbons during live fire training scenarios. *J Occup Environ Hyg.* 12(4):227–34. doi:[10.1080/15459624.2014.955184](https://doi.org/10.1080/15459624.2014.955184) PMID:[25679824](https://pubmed.ncbi.nlm.nih.gov/25679824/)
- Kirk KM, Logan MB (2019). Exposures to air contaminants in compartment fire behavior training (CFBT) using particleboard fuel. *J Occup Environ Hyg.* 16(7):432–9. doi:[10.1080/15459624.2019.1603388](https://doi.org/10.1080/15459624.2019.1603388) PMID:[31021707](https://pubmed.ncbi.nlm.nih.gov/31021707/)
- Kirkham TL, Koehoorn MW, Davies H, Demers PA (2011). Characterization of noise and carbon monoxide exposures among professional firefighters in British Columbia. *Ann Occup Hyg.* 55(7):764–74. doi:[10.1093/annhyg/mer038](https://doi.org/10.1093/annhyg/mer038) PMID:[21765005](https://pubmed.ncbi.nlm.nih.gov/21765005/)
- Laitinen J, Mäkelä M, Mikkola J, Huttu I (2010). Fire fighting trainers' exposure to carcinogenic agents in smoke diving simulators. *Toxicol Lett.* 192(1):61–5. doi:[10.1016/j.toxlet.2009.06.864](https://doi.org/10.1016/j.toxlet.2009.06.864) PMID:[19576276](https://pubmed.ncbi.nlm.nih.gov/19576276/)
- Laitinen J, Mäkelä M, Mikkola J, Huttu I (2012). Firefighters' multiple exposure assessments in practice. *Toxicol Lett.* 213(1):129–33. doi:[10.1016/j.toxlet.2012.06.005](https://doi.org/10.1016/j.toxlet.2012.06.005) PMID:[22710199](https://pubmed.ncbi.nlm.nih.gov/22710199/)
- Laitinen JA, Koponen J, Koikkalainen J, Kiviranta H (2014). Firefighters' exposure to perfluoroalkyl acids and 2-butoxyethanol present in firefighting foams. *Toxicol Lett.* 231(2):227–32. doi:[10.1016/j.toxlet.2014.09.007](https://doi.org/10.1016/j.toxlet.2014.09.007) PMID:[25447453](https://pubmed.ncbi.nlm.nih.gov/25447453/)
- Lam D (2009). [Female firefighters are “equal to men”]. *Diário de Notícias.* 1 June 2009. Available from: <https://www.dn.pt/portugal/sul/bombeiras-sao-iguais-aos-homens-1250127.html>, accessed 13 June 2022. [Portuguese]
- Leary DB, Takazawa M, Kannan K, Khalil N (2020). Perfluoroalkyl substances and metabolic syndrome in firefighters: a pilot study. *J Occup Environ Med.* 62(1):52–7. doi:[10.1097/JOM.0000000000001756](https://doi.org/10.1097/JOM.0000000000001756) PMID:[31658221](https://pubmed.ncbi.nlm.nih.gov/31658221/)
- Lowry WT, Juarez L, Petty CS, Roberts B (1985). Studies of toxic gas production during actual structural fires in the Dallas area. *J Forensic Sci.* 30(1):59–72. doi:[10.1520/JFS10965J](https://doi.org/10.1520/JFS10965J) PMID:[3981122](https://pubmed.ncbi.nlm.nih.gov/3981122/)
- MacSween K, Paton-Walsh C, Roulston C, Gurette EA, Edwards G, Reisen F, et al. (2020). Cumulative firefighter exposure to multiple toxins emitted during prescribed burns in Australia. *Expo Health.* 12(4):721–33. doi:[10.1007/s12403-019-00332-w](https://doi.org/10.1007/s12403-019-00332-w)
- Markowitz JS, Gutterman EM, Schwartz S, Link B, Gorman SM (1989). Acute health effects among firefighters exposed to a polyvinyl chloride (PVC) fire. *Am J*

- Epidemiol.* 129(5):1023–31. doi:[10.1093/oxfordjournals.aje.a115206](https://doi.org/10.1093/oxfordjournals.aje.a115206) PMID:2705423
- Markowitz SB, Garibaldi K, Lilis R, Landrigan PJ (1991). Asbestos exposure and fire fighting. *Ann N Y Acad Sci.* 643:573–7. doi:[10.1111/j.1749-6632.1991.tb24507.x](https://doi.org/10.1111/j.1749-6632.1991.tb24507.x) PMID:1809170
- Materna BL, Jones JR, Sutton PM, Rothman N, Harrison RJ (1992). Occupational exposures in California wildland fire fighting. *Am Ind Hyg Assoc J.* 53(1):69–76. doi:[10.1080/15298669291359311](https://doi.org/10.1080/15298669291359311) PMID:1317093
- Mayer AC, Fent KW, Bertke S, Horn GP, Smith DL, Kerber S, et al. (2019). Firefighter hood contamination: efficiency of laundering to remove PAHs and FRs. *J Occup Environ Hyg.* 16(2):129–40. doi:[10.1080/15459624.2018.1540877](https://doi.org/10.1080/15459624.2018.1540877) PMID:30427284
- Mayer AC, Fent KW, Chen IC, Sammons D, Toennis C, Robertson S, et al. (2021). Characterizing exposures to flame retardants, dioxins, and furans among firefighters responding to controlled residential fires. *Int J Hyg Environ Health.* 236(2021):113782. doi:[10.1016/j.ijheh.2021.113782](https://doi.org/10.1016/j.ijheh.2021.113782) PMID:34119852
- McMahon CK, Bush PB (1992). Forest worker exposure to airborne herbicide residues in smoke from prescribed fires in the southern United States. *Am Ind Hyg Assoc J.* 53(4):265–72. doi:[10.1080/15298669291359636](https://doi.org/10.1080/15298669291359636) PMID:1529920
- McNamara ML, Semmens EO, Gaskill S, Palmer C, Noonan CW, Ward TJ (2012). Base camp personnel exposure to particulate matter during wildland fire suppression activities. *J Occup Environ Hyg.* 9(3):149–56. doi:[10.1080/15459624.2011.652934](https://doi.org/10.1080/15459624.2011.652934) PMID:22364357
- Minty BD, Royston D, Jones JG, Smith DJ, Searing CS, Beeley M (1985). Changes in permeability of the alveolar-capillary barrier in firefighters. *Br J Ind Med.* 42(9):631–4. doi:[10.1136/oem.42.9.631](https://doi.org/10.1136/oem.42.9.631) PMID:3899161
- Miranda AI, Martins V, Cascão P, Amorim JH, Valente J, Borrego C, et al. (2012). Wildland smoke exposure values and exhaled breath indicators in firefighters. *J Toxicol Environ Health A.* 75(13–15):831–43. doi:[10.1080/015287394.2012.690686](https://doi.org/10.1080/015287394.2012.690686) PMID:22788370
- Miranda AI, Martins V, Cascão P, Amorim JH, Valente J, Tavares Ra (2011). [Air quality and firefighter's exposure to experimental forest fires' smoke.] *Silva Lusitana.* 19:23–40. [Portuguese]
- Miranda AI, Martins V, Cascão P, Amorim JH, Valente J, Tavares R, et al. (2010). Monitoring of firefighters exposure to smoke during fire experiments in Portugal. *Environ Int.* 36(7):736–45. doi:[10.1016/j.envint.2010.05.009](https://doi.org/10.1016/j.envint.2010.05.009) PMID:20579737
- Musk AW, Smith TJ, Peters JM, McLaughlin E (1979). Pulmonary function in firefighters: acute changes in ventilatory capacity and their correlates. *Br J Ind Med.* 36(1):29–34. doi:[10.1136/oem.36.1.29](https://doi.org/10.1136/oem.36.1.29) PMID:444439
- Naeher LP, Achtemeier GL, Glitzenstein JS, Streng DR, Macintosh D (2006). Real-time and time-integrated PM_{2.5} and CO from prescribed burns in chipped and non-chipped plots: firefighter and community exposure and health implications. *J Expo Sci Environ Epidemiol.* 16(4):351–61. doi:[10.1038/sj.jes.7500497](https://doi.org/10.1038/sj.jes.7500497) PMID:16736059
- Naeher LP, Barr DB, Adetona O, Simpson CD (2013). Urinary levoglucosan as a biomarker for woodsmoke exposure in wildland firefighters. *Int J Occup Environ Health.* 19(4):304–10. doi:[10.1179/2049396713Y.0000000037](https://doi.org/10.1179/2049396713Y.0000000037) PMID:24588036
- Navarro KM, Cisneros R, Noth EM, Balmes JR, Hammond SK (2017). Occupational exposure to polycyclic aromatic hydrocarbon of wildland firefighters at prescribed and wildland fires. *Environ Sci Technol.* 51(11):6461–9. doi:[10.1021/acs.est.7b00950](https://doi.org/10.1021/acs.est.7b00950) PMID:28498656
- Navarro KM, Cisneros R, Schweizer D, Chowdhary P, Noth EM, Balmes JR, et al. (2019). Incident command post exposure to polycyclic aromatic hydrocarbons and particulate matter during a wildfire. *J Occup Environ Hyg.* 16(11):735–44. doi:[10.1080/15459624.2019.1657579](https://doi.org/10.1080/15459624.2019.1657579) PMID:31545144
- Navarro KM, West MR, O'Dell K, Sen P, Chen IC, Fischer EV, et al. (2021). Exposure to particulate matter and estimation of volatile organic compounds across wildland firefighter job tasks. *Environ Sci Technol.* 55(17):11795–804. doi:[10.1021/acs.est.1c00847](https://doi.org/10.1021/acs.est.1c00847) PMID:34488352
- Neitzel R, Naeher LP, Paulsen M, Dunn K, Stock A, Simpson CD (2009). Biological monitoring of smoke exposure among wildland firefighters: a pilot study comparing urinary methoxyphenols with personal exposures to carbon monoxide, particulate matter, and levoglucosan. *J Expo Sci Environ Epidemiol.* 19(4):349–58. doi:[10.1038/jes.2008.21](https://doi.org/10.1038/jes.2008.21) PMID:18446186
- NIOSH (1991). Health Hazard Evaluation Report No. HETA 91-312-2185. Firefighter's exposure to smoke during fire suppression activities at wildland fires. Prepared by Kelly J. Cincinnati (OH), USA: National Institute for Occupational Safety and Health. Available from: <https://www.cdc.gov/niosh/hhe/reports/pdfs/1991-0312-2185.pdf>, accessed June 2022.
- NIOSH (1992a). Health Hazard Evaluation Report No. HETA 92-045-2260. Characterization of fire fighters' exposures to chemical contaminants during fire suppression operations. Prepared by Kelly J. Cincinnati (OH), USA: National Institute for Occupational Safety and Health. Available from: <http://www.cdc.gov/niosh/hhe/reports/pdfs/1992-0045-2260.pdf>, accessed 30 August 2022.
- NIOSH (1992b). Health Hazard Evaluation Report No. HETA 88-320-2176. US Department of the Interior, National Park Service, Yellowstone National Park, Wyoming. Prepared by Reh CM, Deitchman SD. Cincinnati (OH), USA: National Institute for Occupational Safety and Health. Available from: <https://www.cdc.gov/niosh/hhe/reports/pdfs/1988-0320-2176.pdf>.

- NIOSH (1994). Health Hazard Evaluation Report No. HETA 90-0365-2415. US Department of the Interior, National Park Service, Yosemite National Park, California. Prepared by Reh CM, Letts D, Deitchman S. Cincinnati (OH), USA: National Institute of Occupational Safety and Health. Available from: <https://www.cdc.gov/niosh/hhe/reports/pdfs/1990-0365-2415.pdf>, accessed 15 May 2023.
- NIOSH (1998). Health Hazard Evaluation Report No. HETA 96-0171-2692. Bureau of Alcohol, Tobacco, and Firearms. Washington, DC. Prepared by Kinnes GM, Hine GA. Cincinnati (OH), USA: National Institute for Occupational Safety and Health. Available from: <https://www.cdc.gov/niosh/hhe/reports/pdfs/1996-0171-2692.pdf>, accessed 15 May 2023.
- NIOSH (2016). Health Hazard Evaluation Report No. HETA 2015-0159-3265. Evaluation of diesel exhaust exposures at multiple fire stations in a city fire department. Prepared by Couch J, Broadwater K, de Perio MA. Cincinnati (OH), USA: National Institute for Occupational Safety and Health. Available from: <https://www.cdc.gov/niosh/hhe/reports/pdfs/2015-0159-3265.pdf>, accessed 30 August 2022.
- NIOSH (2019). Health Hazard Evaluation Report No. 2017-0076-3352. Evaluation of wildland fire fighters' exposures to asbestos during a prescribed burn. Prepared by Grant MP. Cincinnati (OH), USA: National Institute for Occupational Safety and Health. Available from: <https://www.cdc.gov/niosh/hhe/reports/pdfs/2017-0076-3352.pdf>, accessed 30 August 2022.
- Park JS, Voss RW, McNeel S, Wu N, Guo T, Wang Y, et al. (2015). High exposure of California firefighters to polybrominated diphenyl ethers. *Environ Sci Technol*. 49(5):2948–58. doi:[10.1021/es5055918](https://doi.org/10.1021/es5055918) PMID:[25643236](https://pubmed.ncbi.nlm.nih.gov/25643236/)
- Peaslee GF, Wilkinson JT, McGuinness SR, Tighe M, Caterisano N, Lee S, et al. (2020). Another pathway for firefighter exposure to per- and polyfluoroalkyl substances: firefighter textiles. *Environ Sci Technol Lett*. 7(8):594–9. doi:[10.1021/acs.estlett.0c00410](https://doi.org/10.1021/acs.estlett.0c00410)
- Pośniak M (2000). [Chemical hazards in fire-fighting environments]. *Med Pr*. 51(4):335–44. [Polish] PMID:[11059406](https://pubmed.ncbi.nlm.nih.gov/11059406/)
- Poutasse CM, Poston WSC, Jahnke SA, Haddock CK, Tidwell LG, Hoffman PD, et al. (2020). Discovery of firefighter chemical exposures using military-style silicone dog tags. *Environ Int*. 142:105818. doi:[10.1016/j.envint.2020.105818](https://doi.org/10.1016/j.envint.2020.105818) PMID:[32521346](https://pubmed.ncbi.nlm.nih.gov/32521346/)
- Reinhardt T, Ottmar R (2000). Smoke exposure at western wildfires. Portland (OR), USA: United States Department of Agriculture, Pacific Northwest Research Station. doi:[10.2737/PNW-RP-525](https://doi.org/10.2737/PNW-RP-525)
- Reinhardt TE, Broyles G (2019). Factors affecting smoke and crystalline silica exposure among wildland firefighters. *J Occup Environ Hyg*. 16(2):151–64. doi:[10.1080/15459624.2018.1540873](https://doi.org/10.1080/15459624.2018.1540873) PMID:[30407130](https://pubmed.ncbi.nlm.nih.gov/30407130/)
- Reinhardt TE, Ottmar RD (2004). Baseline measurements of smoke exposure among wildland firefighters. *J Occup Environ Hyg*. 1(9):593–606. doi:[10.1080/15459620490490101](https://doi.org/10.1080/15459620490490101) PMID:[15559331](https://pubmed.ncbi.nlm.nih.gov/15559331/)
- Reisen F, Brown S, Cheng M (2006). Air toxics in bushfire smoke: firefighter's exposure during prescribed burns. In: Viegas DX, editor. 5th International Conference on Forest Fire Research, November 2006, Coimbra, Portugal.
- Reisen F, Brown SK (2009). Australian firefighters' exposure to air toxics during bushfire burns of autumn 2005 and 2006. *Environ Int*. 35(2):342–52. doi:[10.1016/j.envint.2008.08.011](https://doi.org/10.1016/j.envint.2008.08.011) PMID:[18829114](https://pubmed.ncbi.nlm.nih.gov/18829114/)
- Reisen F, Hansen D, Meyer CP (2011). Exposure to bushfire smoke during prescribed burns and wildfires: firefighters' exposure risks and options. *Environ Int*. 37(2):314–21. doi:[10.1016/j.envint.2010.09.005](https://doi.org/10.1016/j.envint.2010.09.005) PMID:[20956017](https://pubmed.ncbi.nlm.nih.gov/20956017/)
- Robinson MS, Anthony TR, Littau SR, Herckes P, Nelson X, Poplin GS, et al. (2008). Occupational PAH exposures during prescribed pile burns. *Ann Occup Hyg*. 52(6):497–508. doi:[10.1093/annhyg/men027](https://doi.org/10.1093/annhyg/men027) PMID:[18515848](https://pubmed.ncbi.nlm.nih.gov/18515848/)
- Rom WN, Weiden M, Garcia R, Yie TA, Vathesatogkit P, Tse DB, et al. (2002). Acute eosinophilic pneumonia in a New York City firefighter exposed to World Trade Center dust. *Am J Respir Crit Care Med*. 166(6):797–800. doi:[10.1164/rccm.200206-576OC](https://doi.org/10.1164/rccm.200206-576OC) PMID:[12231487](https://pubmed.ncbi.nlm.nih.gov/12231487/)
- Rotander A, Kärrman A, Toms LM, Kay M, Mueller JF, Gómez Ramos MJ (2015a). Novel fluorinated surfactants tentatively identified in firefighters using liquid chromatography quadrupole time-of-flight tandem mass spectrometry and a case-control approach. *Environ Sci Technol*. 49(4):2434–42. doi:[10.1021/es503653n](https://doi.org/10.1021/es503653n) PMID:[25611076](https://pubmed.ncbi.nlm.nih.gov/25611076/)
- Rotander A, Toms L-ML, Aylward L, Kay M, Mueller JF (2015b). Elevated levels of PFOS and PFHxS in firefighters exposed to aqueous film forming foam (AFFF). *Environ Int*. 82:28–34. doi:[10.1016/j.envint.2015.05.005](https://doi.org/10.1016/j.envint.2015.05.005) PMID:[26001497](https://pubmed.ncbi.nlm.nih.gov/26001497/)
- Ruokojärvi P, Aatamila M, Ruuskanen J (2000). Toxic chlorinated and polyaromatic hydrocarbons in simulated house fires. *Chemosphere*. 41(6):825–8. doi:[10.1016/S0045-6535\(99\)00549-4](https://doi.org/10.1016/S0045-6535(99)00549-4) PMID:[10864154](https://pubmed.ncbi.nlm.nih.gov/10864154/)
- Salama KF, Bashawri LA (2017). Biochemical and hematological changes among Saudi firefighters in the eastern province. *Int J Environ Health Eng*. 6(1):2. doi:[10.4103/ijehe.ijehe.12.15](https://doi.org/10.4103/ijehe.ijehe.12.15)
- Sebastião R, Sorte S, Fernandes JM, Miranda AI (2021). Classification of critical levels of CO exposure of firefighters through monitored heart rate. *Sensors (Basel)*. 21(5):1561. doi:[10.3390/s21051561](https://doi.org/10.3390/s21051561) PMID:[33668116](https://pubmed.ncbi.nlm.nih.gov/33668116/)
- Shaw SD, Berger ML, Harris JH, Yun SH, Wu Q, Liao C, et al. (2013). Persistent organic pollutants including polychlorinated and polybrominated dibenzo-*p*-dioxins and dibenzofurans in firefighters from Northern

- California. *Chemosphere*. 91(10):1386–94. doi:[10.1016/j.chemosphere.2012.12.070](https://doi.org/10.1016/j.chemosphere.2012.12.070) PMID:[23395527](https://pubmed.ncbi.nlm.nih.gov/23395527/)
- Shen B, Whitehead TP, Gill R, Dhaliwal J, Brown FR, Petreas M, et al. (2018). Organophosphate flame retardants in dust collected from United States fire stations. *Environ Int*. 112:41–8. doi:[10.1016/j.envint.2017.12.009](https://doi.org/10.1016/j.envint.2017.12.009) PMID:[29247842](https://pubmed.ncbi.nlm.nih.gov/29247842/)
- Shen B, Whitehead TP, McNeel S, Brown FR, Dhaliwal J, Das R, et al. (2015). High levels of polybrominated diphenyl ethers in vacuum cleaner dust from California fire stations. *Environ Sci Technol*. 49(8):4988–94. doi:[10.1021/es505463g](https://doi.org/10.1021/es505463g) PMID:[25798547](https://pubmed.ncbi.nlm.nih.gov/25798547/)
- Sjöström M, Juander A, Strandberg B, Lewné M, Bigert C (2019). Airborne and dermal exposure to polycyclic aromatic hydrocarbons, volatile organic compounds, and particles among firefighters and police investigators. *Ann Work Expo Health*. 63(5):533–45. doi:[10.1093/annweh/wxz030](https://doi.org/10.1093/annweh/wxz030) PMID:[31111145](https://pubmed.ncbi.nlm.nih.gov/31111145/)
- Slaughter JC, Koenig JQ, Reinhardt TE (2004). Association between lung function and exposure to smoke among firefighters at prescribed burns. *J Occup Environ Hyg*. 1(1):45–9. doi:[10.1080/15459620490264490](https://doi.org/10.1080/15459620490264490) PMID:[15202156](https://pubmed.ncbi.nlm.nih.gov/15202156/)
- Smith WR, Montopoli G, Byerly A, Montopoli M, Harlow H, Wheeler AR 3rd (2013). Mercury toxicity in wildland firefighters. *Wilderness Environ Med*. 24(2):141–5. doi:[10.1016/j.wem.2013.01.004](https://doi.org/10.1016/j.wem.2013.01.004) PMID:[23453729](https://pubmed.ncbi.nlm.nih.gov/23453729/)
- Strandberg B, Julander A, Sjöström M, Lewné M, Koca Akdeva H, Bigert C (2018). Evaluation of polyurethane foam passive air sampler (PUF) as a tool for occupational PAH measurements. *Chemosphere*. 190:35–42. doi:[10.1016/j.chemosphere.2017.09.106](https://doi.org/10.1016/j.chemosphere.2017.09.106) PMID:[28985535](https://pubmed.ncbi.nlm.nih.gov/28985535/)
- Swiston JR, Davidson W, Attridge S, Li GT, Brauer M, van Eeden SF (2008). Wood smoke exposure induces a pulmonary and systemic inflammatory response in firefighters. *Eur Respir J*. 32(1):129–38. doi:[10.1183/09031936.00097707](https://doi.org/10.1183/09031936.00097707) PMID:[18256060](https://pubmed.ncbi.nlm.nih.gov/18256060/)
- Trowbridge J, Gerona RR, Lin T, Rudel RA, Bessonneau V, Buren H, et al. (2020). Exposure to perfluoroalkyl substances in a cohort of women firefighters and office workers in San Francisco. *Environ Sci Technol*. 54(6):3363–74. doi:[10.1021/acs.est.9b05490](https://doi.org/10.1021/acs.est.9b05490) PMID:[32100527](https://pubmed.ncbi.nlm.nih.gov/32100527/)
- UK Home Office (2021a). Official statistics. Fire and rescue workforce and pensions statistics: England, April 2020 to March 2021. London, England: United Kingdom Home Office. Available from: <https://www.gov.uk/government/statistics/fire-and-rescue-workforce-and-pensions-statistics-england-april-2020-to-march-2021/fire-and-rescue-workforce-and-pensions-statistics-england-april-2020-to-march-2021>, accessed February 2023.
- Wu CM, Song CC, Chartier R, Kremer J, Naeher L, Adetona O (2021). Characterization of occupational smoke exposure among wildland firefighters in the midwestern United States. *Environ Res*. 193:110541. doi:[10.1016/j.envres.2020.110541](https://doi.org/10.1016/j.envres.2020.110541) PMID:[33249041](https://pubmed.ncbi.nlm.nih.gov/33249041/)
- Young AS, Sparer-Fine EH, Pickard HM, Sunderland EM, Peaslee GF, Allen JG (2021). Per- and polyfluoroalkyl substances (PFAS) and total fluorine in fire station dust. *J Expo Sci Environ Epidemiol*. 31(5):930–42. doi:[10.1038/s41370-021-00288-7](https://doi.org/10.1038/s41370-021-00288-7) PMID:[33542478](https://pubmed.ncbi.nlm.nih.gov/33542478/)