

## Consumption of Alcoholic Beverages

### References to Supplementary Web Tables, Section 1 and Section 2

- Adami HO, Hsing AW, McLaughlin JK *et al.* (1992b). Alcoholism and liver cirrhosis in the etiology of primary liver cancer. *Int J Cancer*, 51:898–902. [doi:10.1002/ijc.2910510611](https://doi.org/10.1002/ijc.2910510611) PMID:1639537
- Adami HO, McLaughlin JK, Hsing AW *et al.* (1992a). Alcoholism and cancer risk: a population-based cohort study. *Cancer Causes Control*, 3:419–425. [doi:10.1007/BF00051354](https://doi.org/10.1007/BF00051354) PMID:1525322
- Adelstein A, White G (1976). Alcoholism and mortality. *Popul Trends*, 6:7–13.
- Akhter M, Kuriyama S, Nakaya N *et al.* (2007). Alcohol consumption is associated with an increased risk of distal colon and rectal cancer in Japanese men: the Miyagi Cohort Study. *Eur J Cancer*, 43:383–390. [doi:10.1016/j.ejca.2006.09.020](https://doi.org/10.1016/j.ejca.2006.09.020) PMID:17150353
- Allen NE, Beral V, Casabonne D *et al.*; Million Women Study Collaborators (2009). Moderate alcohol intake and cancer incidence in women. *J Natl Cancer Inst*, 101:296–305. [doi:10.1093/jnci/djn514](https://doi.org/10.1093/jnci/djn514) PMID:19244173
- Anderson LN, Cotterchio M, Gallinger S (2009). Lifestyle, dietary, and medical history factors associated with pancreatic cancer risk in Ontario, Canada. *Cancer Causes Control*, 20:825–834. [doi:10.1007/s10552-009-9303-5](https://doi.org/10.1007/s10552-009-9303-5) PMID:19194662
- Ansems TM, van der Pols JC, Hughes MC *et al.* (2008). Alcohol intake and risk of skin cancer: a prospective study. *Eur J Clin Nutr*, 62:162–170. [doi:10.1038/sj.ejcn.1602717](https://doi.org/10.1038/sj.ejcn.1602717) PMID:17392700
- Anwar WA, Abdel-Rahman SZ, El-Zein RA *et al.* (1996). Genetic polymorphism of GSTM1, CYP2E1 and CYP2D6 in Egyptian bladder cancer patients. *Carcinogenesis*, 17:1923–1929. [doi:10.1093/carcin/17.9.1923](https://doi.org/10.1093/carcin/17.9.1923) PMID:8824515
- Arndt M, Rydzanicz M, Pabiszczak M *et al.* (2008). [Distribution of alcohol dehydrogenase (ADH1C) genotypes in subjects with tobacco smoke-associated laryngeal cancer]. *Przegl Lek*, 65:466–469. PMID:19189524
- Asakage T, Yokoyama A, Haneda T *et al.* (2007). Genetic polymorphisms of alcohol and aldehyde dehydrogenases, and drinking, smoking and diet in Japanese men with oral and pharyngeal squamous cell carcinoma. *Carcinogenesis*, 28:865–874. [doi:10.1093/carcin/bgl206](https://doi.org/10.1093/carcin/bgl206) PMID:17071628
- Baglietto L, English DR, Gertig DM *et al.* (2005). Does dietary folate intake modify effect of alcohol consumption on breast cancer risk? Prospective cohort study. *BMJ*, 331:807–900. [doi:10.1136/bmj.38551.446470.06](https://doi.org/10.1136/bmj.38551.446470.06) PMID:16087654
- Baglietto L, Severi G, English DR *et al.* (2006). Alcohol consumption and prostate cancer risk: results from the Melbourne collaborative cohort study. *Int J Cancer*, 119:1501–1504. [doi:10.1002/ijc.21983](https://doi.org/10.1002/ijc.21983) PMID:16615108
- Balder HF, Goldbohm RA, van den Brandt PA (2005). Dietary patterns associated with male lung cancer risk in the Netherlands Cohort Study. *Cancer Epidemiol Biomarkers Prev*, 14:483–490. [doi:10.1158/1055-9965.EPI-04-0353](https://doi.org/10.1158/1055-9965.EPI-04-0353) PMID:15734976
- Bandera EV, Freudenheim JL, Marshall JR *et al.* (1997). Diet and alcohol consumption and lung cancer risk in the New York State Cohort (United States). *Cancer Causes Control*, 8:828–840. [doi:10.1023/A:1018456127018](https://doi.org/10.1023/A:1018456127018) PMID:9427425
- Beji NK, Reis N (2007). Risk factors for breast cancer in Turkish women: a hospital-based case-control study. *Eur J Cancer Care (Engl)*, 16:178–184. [doi:10.1111/j.1365-2354.2006.00711.x](https://doi.org/10.1111/j.1365-2354.2006.00711.x) PMID:17371428

- Benedetti A, Parent ME, Siemiatycki J (2006). Consumption of alcoholic beverages and risk of lung cancer: results from two case-control studies in Montreal, Canada. *Cancer Causes Control*, 17:469–480. [doi:10.1007/s10552-005-0496-y](https://doi.org/10.1007/s10552-005-0496-y) PMID:16596299
- Benedetti A, Parent ME, Siemiatycki J (2009). Lifetime consumption of alcoholic beverages and risk of 13 types of cancer in men: results from a case-control study in Montreal. *Cancer Detect Prev*, 32:352–362. [doi:10.1016/j.canep.2009.03.001](https://doi.org/10.1016/j.canep.2009.03.001) PMID:19588541
- Beral V, Bull D, Reeves G; Million Women Study Collaborators (2005). Endometrial cancer and hormone-replacement therapy in the Million Women Study. *Lancet*, 365:1543–1551. [doi:10.1016/S0140-6736\(05\)66455-0](https://doi.org/10.1016/S0140-6736(05)66455-0) PMID:15866308
- Bernstein JL, Thompson WD, Risch N, Holford TR (1992). Risk factors predicting the incidence of second primary breast cancer among women diagnosed with a first primary breast cancer. *Am J Epidemiol*, 136:925–936. PMID:1456269
- Berstad P, Ma H, Bernstein L, Ursin G (2008). Alcohol intake and breast cancer risk among young women. *Breast Cancer Res Treat*, 108:113–120. [doi:10.1007/s10549-007-9578-8](https://doi.org/10.1007/s10549-007-9578-8) PMID:17468952
- Bessaoud F, Daurès JP (2008). Patterns of alcohol (especially wine) consumption and breast cancer risk: a case-control study among a population in Southern France. *Ann Epidemiol*, 18:467–475. [doi:10.1016/j.annepidem.2008.02.001](https://doi.org/10.1016/j.annepidem.2008.02.001) PMID:18440826
- Boccia S, Cadoni G, Sayed-Tabatabaei FA *et al.* (2008). CYP1A1, CYP2E1, GSTM1, GSTT1, EPHX1 exons 3 and 4, and NAT2 polymorphisms, smoking, consumption of alcohol and fruit and vegetables and risk of head and neck cancer. *J Cancer Res Clin Oncol*, 134:93–100. [doi:10.1007/s00432-007-0254-5](https://doi.org/10.1007/s00432-007-0254-5) PMID:17611777
- Boccia S, Hashibe M, Galli P *et al.* (2009). Aldehyde dehydrogenase 2 and head and neck cancer: a meta-analysis implementing a Mendelian randomization approach. *Cancer Epidemiol Biomarkers Prev*, 18:248–254. [doi:10.1158/1055-9965.EPI-08-0462](https://doi.org/10.1158/1055-9965.EPI-08-0462) PMID:19124505
- Boccia S, Sayed-Tabatabaei FA, Persiani R *et al.* (2007). Polymorphisms in metabolic genes, their combination and interaction with tobacco smoke and alcohol consumption and risk of gastric cancer: a case-control study in an Italian population. *BMC Cancer*, 7:206. [doi:10.1186/1471-2407-7-206](https://doi.org/10.1186/1471-2407-7-206) PMID:17996038
- Boeing H (2002) Alcohol and cancer of the upper gastrointestinal tract: First analysis of the EPIC data. In: Riboli, E. & Lambert, R., eds, *Nutrition and lifestyles: Opportunities for Cancer Prevention* (IARC Scientific Publications No 156), Lyon, International Agency for Research on Cancer, pp. 151–154.
- Boffetta P, Garfinkel L (1990). Alcohol drinking and mortality among men enrolled in an American Cancer Society prospective study. *Epidemiology*, 1:342–348. [doi:10.1097/00001648-199009000-00003](https://doi.org/10.1097/00001648-199009000-00003) PMID:2078609
- Boffetta P, Stellman SD, Garfinkel L (1989). A case-control study of multiple myeloma nested in the American Cancer Society prospective study. *Int J Cancer*, 43:554–559. [doi:10.1002/ijc.2910430404](https://doi.org/10.1002/ijc.2910430404) PMID:2703267
- Boffetta P, Ye W, Adami H-O *et al.* (2001). Risk of cancers of the lung, head and neck in patients hospitalized for alcoholism in Sweden. *Br J Cancer*, 85:678–682. [doi:10.1054/bjoc.2001.1986](https://doi.org/10.1054/bjoc.2001.1986) PMID:11531251
- Boice JD Jr, Mandel JS, Doody MM (1995). Breast cancer among radiologic technologists. *JAMA*, 274:394–401. [doi:10.1001/jama.274.5.394](https://doi.org/10.1001/jama.274.5.394) PMID:7616635
- Bongaerts BW, de Goeij AF, van den Brandt PA, Weijnenberg MP (2006). Alcohol and the risk of colon and rectal cancer with mutations in the K-ras gene. *Alcohol*, 38:147–154. [doi:10.1016/j.alcohol.2006.06.003](https://doi.org/10.1016/j.alcohol.2006.06.003) PMID:16905440

- Bongaerts BW, van den Brandt PA, Goldbohm RA *et al.* (2008). Alcohol consumption, type of alcoholic beverage and risk of colorectal cancer at specific subsites. *Int J Cancer*, 123:2411–2417. [doi:10.1002/ijc.23774](https://doi.org/10.1002/ijc.23774) PMID:18752250
- Bouchardy C, Hirvonen A, Coutelle C *et al.* (2000). Role of alcohol dehydrogenase 3 and cytochrome P-4502E1 genotypes in susceptibility to cancers of the upper aerodigestive tract. *Int J Cancer*, 87:734–740. [doi:10.1002/1097-0215\(20000901\)87:5<734::AID-IJC17>3.0.CO;2-E](https://doi.org/10.1002/1097-0215(20000901)87:5<734::AID-IJC17>3.0.CO;2-E) PMID:10925369
- Breast Cancer Association Consortium (2006). Commonly studied single-nucleotide polymorphisms and breast cancer: results from the Breast Cancer Association Consortium. *J Natl Cancer Inst*, 98:1382–1396. [doi:10.1093/jnci/djj374](https://doi.org/10.1093/jnci/djj374) PMID:17018785
- Breslow RA, Graubard BI, Sinha R, Subar AF (2000). Diet and lung cancer mortality: a 1987 National Health Interview Survey cohort study. *Cancer Causes Control*, 11:419–431. [doi:10.1023/A:1008996208313](https://doi.org/10.1023/A:1008996208313) PMID:10877335
- Breslow RA, Wideroff L, Graubard BI *et al.*; First National Health and Nutrition Examination Survey of the United States (1999). Alcohol and prostate cancer in the NHANES I epidemiologic follow-up study. *Ann Epidemiol*, 9:254–261. [doi:10.1016/S1047-2797\(98\)00071-4](https://doi.org/10.1016/S1047-2797(98)00071-4) PMID:10332931
- Brinton LA, Richesson DA, Gierach GL *et al.* (2008). Prospective evaluation of risk factors for male breast cancer. *J Natl Cancer Inst*, 100:1477–1481. [doi:10.1093/jnci/djn329](https://doi.org/10.1093/jnci/djn329) PMID:18840816
- Brinton LA, Carreon JD, Gierach GL *et al.* (2010). Etiologic factors for male breast cancer in the U.S. Veterans Affairs medical care system database. *Breast Cancer Res Treat*. 119:185–192. [PMID:19330525](https://doi.org/10.1007/s12532-010-9525-2)
- Brockmüller J, Cascorbi I, Kerb R, Roots I (1996). Combined analysis of inherited polymorphisms in arylamine N-acetyltransferase 2, glutathione S-transferases M1 and T1, microsomal epoxide hydrolase, and cytochrome P450 enzymes as modulators of bladder cancer risk. *Cancer Res*, 56:3915–3925. [PMID:8752158](https://doi.org/10.1158/0008-5472.CCR.95-2087)
- Brown LM, Gridley G, Wu AH *et al.* (2009) Low level alcohol intake, cigarette smoking and risk of breast cancer in Asian-American women. *Breast Cancer Res Treat*. PMID:19597702
- Buch SC, Nazar-Stewart V, Weissfeld JL, Romkes M (2008). Case-control study of oral and oropharyngeal cancer in whites and genetic variation in eight metabolic enzymes. *Head Neck*, 30:1139–1147. [doi:10.1002/hed.20867](https://doi.org/10.1002/hed.20867) PMID:18642288
- Cai L, Yu SZ, Zhan ZF (2001). Cytochrome P450 2E1 genetic polymorphism and gastric cancer in Changde, Fujian Province. *World J Gastroenterol*, 7:792–795. [PMID:11854903](https://doi.org/10.3748/wjg.v7i7.792)
- Cantor KP, Lynch CF, Hildesheim ME *et al.* (1998). Drinking water source and chlorination byproducts. I. Risk of bladder cancer. *Epidemiology*, 9:21–28. [doi:10.1097/00001648-199801000-00007](https://doi.org/10.1097/00001648-199801000-00007) [PMID:9430264](https://doi.org/10.1093/epi/kfn024)
- Cao HX, Gao CM, Takezaki T *et al.* (2008). Genetic polymorphisms of methylenetetrahydrofolate reductase and susceptibility to colorectal cancer. *Asian Pac J Cancer Prev*, 9:203–208. [PMID:18712959](https://doi.org/10.1158/1545-1575.PAP007037)
- Cao W, Cai L, Rao JY *et al.* (2005). Tobacco smoking, GSTP1 polymorphism, and bladder carcinoma. *Cancer*, 104:2400–2408. [doi:10.1002/cncr.21446](https://doi.org/10.1002/cncr.21446) PMID:16240451
- Carstensen JM, Bygren LO, Hatschek T (1990). Cancer incidence among Swedish brewery workers. *Int J Cancer*, 45:393–396. [doi:10.1002/ijc.2910450302](https://doi.org/10.1002/ijc.2910450302) PMID:2407667
- Casey R, Piazzon-Fevre K, Raverdy N *et al.* (2007). Case-control study of lymphoid neoplasm in three French areas: description, alcohol and tobacco consumption. *Eur J Cancer Prev*, 16:142–150. [doi:10.1097/01.ccej.0000220629.00729.65](https://doi.org/10.1097/01.ccej.0000220629.00729.65) PMID:17297390

- Cerhan JR, Torner JC, Lynch CF *et al.* (1997). Association of smoking, body mass, and physical activity with risk of prostate cancer in the Iowa 65+ Rural Health Study (United States). *Cancer Causes Control*, 8:229–238. [doi:10.1023/A:1018428531619](https://doi.org/10.1023/A:1018428531619) PMID:9134247
- Chang ET, Canchola AJ, Lee VS *et al.* (2007). Wine and other alcohol consumption and risk of ovarian cancer in the California Teachers Study cohort. *Cancer Causes Control*, 18:91–103. [doi:10.1007/s10552-006-0083-x](https://doi.org/10.1007/s10552-006-0083-x) PMID:17186425
- Chang ET, Smedby KE, Zhang SM *et al.* (2004). Alcohol intake and risk of non-Hodgkin lymphoma in men and women. *Cancer Causes Control*, 15:1067–1076. [doi:10.1007/s10552-004-2234-2](https://doi.org/10.1007/s10552-004-2234-2) PMID:15801490
- Chao C (2007). Associations between beer, wine, and liquor consumption and lung cancer risk: a meta-analysis. *Cancer Epidemiol Biomarkers Prev*, 16:2436–2447. [doi:10.1158/1055-9965.EPI-07-0386](https://doi.org/10.1158/1055-9965.EPI-07-0386) PMID:18006934
- Chao C, Haque R, Van Den Eeden SK *et al.* (2010). Red wine consumption and risk of prostate cancer: the California men’s health study. *Int J Cancer*, 126:171–179. [doi:10.1002/ijc.24637](https://doi.org/10.1002/ijc.24637) PMID:19521962
- Chao C, Slezak JM, Caan BJ, Quinn VP (2008). Alcoholic beverage intake and risk of lung cancer: the California Men’s Health Study. *Cancer Epidemiol Biomarkers Prev*, 17:2692–2699. [doi:10.1158/1055-9965.EPI-08-0410](https://doi.org/10.1158/1055-9965.EPI-08-0410) PMID:18843011
- Chao YC, Wang LS, Hsieh TY *et al.* (2000). Chinese alcoholic patients with esophageal cancer are genetically different from alcoholics with acute pancreatitis and liver cirrhosis. *Am J Gastroenterol*, 95:2958–2964. [doi:10.1111/j.1572-0241.2000.02328.x](https://doi.org/10.1111/j.1572-0241.2000.02328.x) PMID:11051375
- Chen K, Jiang Q, Ma X *et al.* (2005). Alcohol drinking and colorectal cancer: a population-based prospective cohort study in China. *Eur J Epidemiol*, 20:149–154. [doi:10.1007/s10654-004-2953-4](https://doi.org/10.1007/s10654-004-2953-4) PMID:15792281
- Chen WY, Colditz GA, Rosner B *et al.* (2002a). Use of postmenopausal hormones, alcohol, and risk for invasive breast cancer. *Ann Intern Med*, 137:798–804. PMID:12435216
- Chiu BC-H, Cerhan JR, Gapstur SM *et al.* (1999). Alcohol consumption and non-Hodgkin lymphoma in a cohort of older women. *Br J Cancer*, 80:1476–1482. [doi:10.1038/sj.bjc.6690547](https://doi.org/10.1038/sj.bjc.6690547) PMID:10424754
- Chlebowski RT, Anderson GL, Lane DS *et al.*; Women’s Health Initiative Investigators (2007). Predicting risk of breast cancer in postmenopausal women by hormone receptor status. *J Natl Cancer Inst*, 99:1695–1705. [doi:10.1093/jnci/djm224](https://doi.org/10.1093/jnci/djm224) PMID:18000216
- Choi JY, Abel J, Neuhaus T *et al.* (2003). Role of alcohol and genetic polymorphisms of CYP2E1 and ALDH2 in breast cancer development. *Pharmacogenetics*, 13:67–72. [doi:10.1097/00008571-200302000-00002](https://doi.org/10.1097/00008571-200302000-00002) PMID:12563175
- Chow WH, Schuman LM, McLaughlin JK *et al.* (1992). A cohort study of tobacco use, diet, occupation, and lung cancer mortality. *Cancer Causes Control*, 3:247–254. [doi:10.1007/BF00124258](https://doi.org/10.1007/BF00124258) PMID:1610971
- Chyou P-H, Nomura AMY, Stemmermann GN (1993). A prospective study of diet, smoking, and lower urinary tract cancer. *Ann Epidemiol*, 3:211–216. [doi:10.1016/1047-2797\(93\)90021-U](https://doi.org/10.1016/1047-2797(93)90021-U) PMID:8275191
- Chyou P-H, Nomura AMY, Stemmermann GN (1995). Diet, alcohol, smoking and cancer of the upper aerodigestive tract: a prospective study among Hawaii Japanese men. *Int J Cancer*, 60:616–621. [doi:10.1002/ijc.2910600508](https://doi.org/10.1002/ijc.2910600508) PMID:7860134
- Chyou P-H, Nomura AMY, Stemmermann GN (1996). A prospective study of colon and rectal cancer among Hawaii Japanese men. *Ann Epidemiol*, 6:276–282. [doi:10.1016/S1047-2797\(96\)00047-6](https://doi.org/10.1016/S1047-2797(96)00047-6) PMID:8876837

- Colditz GA, Rosner B (2000). Cumulative risk of breast cancer to age 70 years according to risk factor status: data from the Nurses' Health Study. *Am J Epidemiol*, 152:950–964. [doi:10.1093/aje/152.10.950](https://doi.org/10.1093/aje/152.10.950) [PMID:11092437](https://pubmed.ncbi.nlm.nih.gov/11092437/)
- Coughlin SS, Calle EE, Patel AV, Thun MJ (2000). Predictors of pancreatic cancer mortality among a large cohort of United States adults. *Cancer Causes Control*, 11:915–923. [doi:10.1023/A:1026580131793](https://doi.org/10.1023/A:1026580131793) [PMID:11142526](https://pubmed.ncbi.nlm.nih.gov/11142526/)
- Coutelle C, Höhn B, Benesova M *et al.* (2004). Risk factors in alcohol associated breast cancer: alcohol dehydrogenase polymorphism and estrogens. *Int J Oncol*, 25:1127–1132. [PMID:15375565](https://pubmed.ncbi.nlm.nih.gov/15375565/)
- Coutelle C, Ward PJ, Fleury B *et al.* (1997). Laryngeal and oropharyngeal cancer, and alcohol dehydrogenase 3 and glutathione S-transferase M1 polymorphisms. *Hum Genet*, 99:319–325. [doi:10.1007/s004390050365](https://doi.org/10.1007/s004390050365) [PMID:9050916](https://pubmed.ncbi.nlm.nih.gov/9050916/)
- Covolo L, Gelatti U, Talamini R *et al.* (2005). Alcohol dehydrogenase 3, glutathione S-transferase M1 and T1 polymorphisms, alcohol consumption and hepatocellular carcinoma (Italy). *Cancer Causes Control*, 16:831–838. [doi:10.1007/s10552-005-2302-2](https://doi.org/10.1007/s10552-005-2302-2) [PMID:16132793](https://pubmed.ncbi.nlm.nih.gov/16132793/)
- Cox A, Dunning AM, Garcia-Closas M *et al.*; Kathleen Cunningham Foundation Consortium for Research into Familial Breast Cancer; Breast Cancer Association Consortium (2007). A common coding variant in CASP8 is associated with breast cancer risk. *Nat Genet*, 39:352–358. [doi:10.1038/ng1981](https://doi.org/10.1038/ng1981) [PMID:17293864](https://pubmed.ncbi.nlm.nih.gov/17293864/)
- Cox B, Sneyd MJ, Paul C, Skegg DCG (2006). Risk factors for prostate cancer: A national case-control study. *Int J Cancer*, 119:1690–1694. [doi:10.1002/ijc.22022](https://doi.org/10.1002/ijc.22022) [PMID:16646073](https://pubmed.ncbi.nlm.nih.gov/16646073/)
- Curtin K, Slattery ML, Ulrich CM *et al.* (2007). Genetic polymorphisms in one-carbon metabolism: associations with CpG island methylator phenotype (CIMP) in colon cancer and the modifying effects of diet. *Carcinogenesis*, 28:1672–1679. [doi:10.1093/carcin/bgm089](https://doi.org/10.1093/carcin/bgm089) [PMID:17449906](https://pubmed.ncbi.nlm.nih.gov/17449906/)
- Day GL, Blot WJ, Shore RE *et al.* (1994a). Second cancers following oral and pharyngeal cancers: role of tobacco and alcohol. *J Natl Cancer Inst*, 86:131–137. [doi:10.1093/jnci/86.2.131](https://doi.org/10.1093/jnci/86.2.131) [PMID:8271296](https://pubmed.ncbi.nlm.nih.gov/8271296/)
- De Stefani E, Deneo-Pellegrini H, Boffetta P *et al.* (2009). Dietary patterns and risk of cancer: a factor analysis in Uruguay. *Int J Cancer*, 124:1391–1397. [doi:10.1002/ijc.24035](https://doi.org/10.1002/ijc.24035) [PMID:19058195](https://pubmed.ncbi.nlm.nih.gov/19058195/)
- Dean G, MacLennan R, McLoughlin H, Shelley E (1979). Causes of death of blue-collar workers at a Dublin brewery, 1954–73. *Br J Cancer*, 40:581–589. [doi:10.1038/bjc.1979.223](https://doi.org/10.1038/bjc.1979.223) [PMID:497108](https://pubmed.ncbi.nlm.nih.gov/497108/)
- Deandrea S, Bertuccio P, Chatenoud L *et al.* (2007). Reply to 'Alcohol consumption and risk of Hodgkin's lymphoma and multiple myeloma: a multicentre case-control study' by Gorini *et al.* *Ann Oncol*, 18:1119–1121. [doi:10.1093/annonc/mdm203](https://doi.org/10.1093/annonc/mdm203) [PMID:17586754](https://pubmed.ncbi.nlm.nih.gov/17586754/)
- Deandrea S, Talamini R, Foschi R *et al.* (2008). Alcohol and breast cancer risk defined by estrogen and progesterone receptor status: a case-control study. *Cancer Epidemiol Biomarkers Prev*, 17:2025–2028. [doi:10.1158/1055-9965.EPI-08-0157](https://doi.org/10.1158/1055-9965.EPI-08-0157) [PMID:18708394](https://pubmed.ncbi.nlm.nih.gov/18708394/)
- Dennis LK (2000). Meta-analysis for combining relative risks of alcohol consumption and prostate cancer. *Prostate*, 42:56–66. [doi:10.1002/\(SICI\)1097-0045\(20000101\)42:1<56::AID-PROS7>3.0.CO;2-P](https://doi.org/10.1002/(SICI)1097-0045(20000101)42:1<56::AID-PROS7>3.0.CO;2-P) [PMID:10579799](https://pubmed.ncbi.nlm.nih.gov/10579799/)
- Dikshit RP, Boffetta P, Bouchardy C *et al.* (2005). Risk factors for the development of second primary tumors among men after laryngeal and hypopharyngeal carcinoma. *Cancer*, 103:2326–2333. [doi:10.1002/cncr.21051](https://doi.org/10.1002/cncr.21051) [PMID:15852357](https://pubmed.ncbi.nlm.nih.gov/15852357/)
- Ding J, Li S, Wu J *et al.* (2008). Alcohol dehydrogenase-2 and aldehyde dehydrogenase-2 genotypes, alcohol drinking and the risk of primary hepatocellular carcinoma in a Chinese population. *Asian Pac J Cancer Prev*, 9:31–35. [PMID:18439068](https://pubmed.ncbi.nlm.nih.gov/18439068/)
- Djoussé L, Dorgan JF, Zhang Y *et al.* (2002). Alcohol consumption and risk of lung cancer: the Framingham Study. *J Natl Cancer Inst*, 94:1877–1882. [doi:10.1093/jnci/94.24.1877](https://doi.org/10.1093/jnci/94.24.1877) [PMID:12488481](https://pubmed.ncbi.nlm.nih.gov/12488481/)

- Djoussé L, Schatzkin A, Chibnik LB *et al.* (2004). Alcohol consumption and the risk of bladder cancer in the Framingham Heart Study. *J Natl Cancer Inst*, 96:1397–1400. [doi:10.1093/jnci/djh263](https://doi.org/10.1093/jnci/djh263) [PMID:15367573](https://pubmed.ncbi.nlm.nih.gov/15367573/)
- Doll R, Peto R, Boreham J, Sutherland I (2005). Mortality from cancer in relation to smoking: 50 years observations on British doctors. *Br J Cancer*, 92:426–429. [PMID:15668706](https://pubmed.ncbi.nlm.nih.gov/15668706/)
- Doll R, Peto R, Hall E *et al.* (1994). Mortality in relation to consumption of alcohol: 13 years' observations on male British doctors. *BMJ*, 309:911–918. [doi:10.1136/bmj.309.6959.911](https://doi.org/10.1136/bmj.309.6959.911) [PMID:7950661](https://pubmed.ncbi.nlm.nih.gov/7950661/)
- Dolle JM, Daling JR, White E *et al.* (2009). Risk factors for triple-negative breast cancer in women under the age of 45 years. *Cancer Epidemiol Biomarkers Prev*, 18:1157–1166. [doi:10.1158/1055-9965.EPI-08-1005](https://doi.org/10.1158/1055-9965.EPI-08-1005) [PMID:19336554](https://pubmed.ncbi.nlm.nih.gov/19336554/)
- Donato F, Gelatti U, Tagger A *et al.* (2001). Intrahepatic cholangiocarcinoma and hepatitis C and B virus infection, alcohol intake, and hepatolithiasis: a case-control study in Italy. *Cancer Causes Control*, 12:959–964. [doi:10.1023/A:1013747228572](https://doi.org/10.1023/A:1013747228572) [PMID:11808716](https://pubmed.ncbi.nlm.nih.gov/11808716/)
- Duffy CM, Assaf A, Cyr M *et al.* (2009). Alcohol and folate intake and breast cancer risk in the WHI Observational Study. *Breast Cancer Res Treat*, 116:551–562. [doi:10.1007/s10549-008-0167-2](https://doi.org/10.1007/s10549-008-0167-2) [PMID:18785003](https://pubmed.ncbi.nlm.nih.gov/18785003/)
- Dumeaux V, Lund E, Hjartåker A (2004). Use of oral contraceptives, alcohol, and risk for invasive breast cancer. *Cancer Epidemiol Biomarkers Prev*, 13:1302–1307. [PMID:15298950](https://pubmed.ncbi.nlm.nih.gov/15298950/)
- Efird JT, Friedman GD, Sidney S *et al.* (2004). The risk for malignant primary adult-onset glioma in a large, multiethnic, managed-care cohort: cigarette smoking and other lifestyle behaviors. *J Neurooncol*, 68:57–69. [doi:10.1023/B:NEON.0000024746.87666.ed](https://doi.org/10.1023/B:NEON.0000024746.87666.ed) [PMID:15174522](https://pubmed.ncbi.nlm.nih.gov/15174522/)
- el-Zein R, Zwischenberger JB, Wood TG *et al.* (1997). Combined genetic polymorphism and risk for development of lung cancer. *Mutat Res*, 381:189–200. [doi:10.1016/S0027-5107\(97\)00166-8](https://doi.org/10.1016/S0027-5107(97)00166-8) [PMID:9434875](https://pubmed.ncbi.nlm.nih.gov/9434875/)
- Ellison LF (2000). Tea and other beverage consumption and prostate cancer risk: a Canadian retrospective cohort study. *Eur J Cancer Prev*, 9:125–130. [doi:10.1097/00008469-200004000-00009](https://doi.org/10.1097/00008469-200004000-00009) [PMID:10830580](https://pubmed.ncbi.nlm.nih.gov/10830580/)
- Engeset D, Dyachenko A, Ciampi A, Lund E (2009). Dietary patterns and risk of cancer of various sites in the Norwegian European Prospective Investigation into Cancer and Nutrition cohort: the Norwegian Women and Cancer study. *Eur J Cancer Prev*, 18:69–75. [doi:10.1097/CEJ.0b013e328305a091](https://doi.org/10.1097/CEJ.0b013e328305a091) [PMID:19077568](https://pubmed.ncbi.nlm.nih.gov/19077568/)
- Eom SY, Zhang YW, Kim SH *et al.* (2009). Influence of NQO1, ALDH2, and CYP2E1 genetic polymorphisms, smoking, and alcohol drinking on the risk of lung cancer in Koreans. *Cancer Causes Control*, 20:137–145. [doi:10.1007/s10552-008-9225-7](https://doi.org/10.1007/s10552-008-9225-7) [PMID:18798003](https://pubmed.ncbi.nlm.nih.gov/18798003/)
- Ericson U, Sonestedt E, Gullberg B *et al.* (2007). High folate intake is associated with lower breast cancer incidence in postmenopausal women in the Malmö Diet and Cancer cohort. *Am J Clin Nutr*, 86:434–443. [PMID:17684216](https://pubmed.ncbi.nlm.nih.gov/17684216/)
- Fan C, Jin M, Chen K *et al.* (2007). Case-only study of interactions between metabolic enzymes and smoking in colorectal cancer. *BMC Cancer*, 7:115. [doi:10.1186/1471-2407-7-115](https://doi.org/10.1186/1471-2407-7-115) [PMID:17603900](https://pubmed.ncbi.nlm.nih.gov/17603900/)
- Fan L, Cai L (2009). [Meta-analysis on the relationship between alcohol consumption and lung cancer risk]. *Wei Sheng Yan Jiu*, 38:85–89. [PMID:19267084](https://pubmed.ncbi.nlm.nih.gov/19267084/)
- Fan Y, Yuan JM, Wang R *et al.* (2008). Alcohol, tobacco, and diet in relation to esophageal cancer: the Shanghai Cohort Study. *Nutr Cancer*, 60:354–363. [doi:10.1080/01635580701883011](https://doi.org/10.1080/01635580701883011) [PMID:18444169](https://pubmed.ncbi.nlm.nih.gov/18444169/)
- Farker K, Lehmann MH, Kästner R *et al.* (1998a). CYP2E1 genotyping in renal cell/urothelial cancer patients in comparison with control populations. *Int J Clin Pharmacol Ther*, 36:463–468. [PMID:9760005](https://pubmed.ncbi.nlm.nih.gov/9760005/)

- Farker K, Lehmann MH, Oelschlägel B *et al.* (1998b). Impact of CYP2E1 genotype in renal cell and urothelial cancer patients. *Exp Toxicol Pathol*, 50:425–431. [PMID:9784018](#)
- Feigelson HS, Jonas CR, Robertson AS *et al.* (2003). Alcohol, folate, methionine, and risk of incident breast cancer in the American Cancer Society Cancer Prevention Study II Nutrition Cohort. *Cancer Epidemiol Biomarkers Prev*, 12:161–164. [PMID:12582027](#)
- Ferrari P, Jenab M, Norat T *et al.* (2007). Lifetime and baseline alcohol intake and risk of colon and rectal cancers in the European prospective investigation into cancer and nutrition (EPIC). *Int J Cancer*, 121:2065–2072. [doi:10.1002/ijc.22966](#) [PMID:17640039](#)
- Flood A, Caprario L, Chatterjee N *et al.* (2002). Folate, methionine, alcohol, and colorectal cancer in a prospective study of women in the United States. *Cancer Causes Control*, 13:551–561. [doi:10.1023/A:1016330609603](#) [PMID:12195645](#)
- Folsom AR, Demissie Z, Harnack L; Iowa Women's Health Study (2003). Glycemic index, glycemic load, and incidence of endometrial cancer: the Iowa women's health study. *Nutr Cancer*, 46:119–124. [doi:10.1207/S15327914NC4602\\_03](#) [PMID:14690786](#)
- Franceschi S, Talamini R, Barra S *et al.* (1990). Smoking and drinking in relation to cancers of the oral cavity, pharynx, larynx, and esophagus in northern Italy. *Cancer Res*, 50:6502–6507. [PMID:2208109](#)
- Freedman DM, Sigurdson A, Doody MM *et al.* (2003). Risk of melanoma in relation to smoking, alcohol intake, and other factors in a large occupational cohort. *Cancer Causes Control*, 14:847–857. [doi:10.1023/B:CACO.0000003839.56954.73](#) [PMID:14682442](#)
- Freedman ND, Abnet CC, Leitzmann MF *et al.* (2007). A prospective study of tobacco, alcohol, and the risk of esophageal and gastric cancer subtypes. *Am J Epidemiol*, 165:1424–1433. [doi:10.1093/aje/kwm051](#) [PMID:17420181](#)
- Freedman ND, Abnet CC, Leitzmann MF *et al.* (2007a). A prospective study of tobacco, alcohol, and the risk of esophageal and gastric cancer subtypes. *Am J Epidemiol*, 165:1424–1433. [doi:10.1093/aje/kwm051](#) [PMID:17420181](#)
- Freedman ND, Schatzkin A, Leitzmann MF *et al.* (2007b). Alcohol and head and neck cancer risk in a prospective study. *Br J Cancer*, 96:1469–1474. [PMID:17387340](#)
- Freudenheim JL, Ambrosone CB, Moysich KB *et al.* (1999). Alcohol dehydrogenase 3 genotype modification of the association of alcohol consumption with breast cancer risk. *Cancer Causes Control*, 10:369–377. [doi:10.1023/A:1008950717205](#) [PMID:10530606](#)
- Freudenheim JL, Ram M, Nie J *et al.* (2003). Lung cancer in humans is not associated with lifetime total alcohol consumption or with genetic variation in alcohol dehydrogenase 3 (ADH3). *J Nutr*, 133:3619–3624. [PMID:14608084](#)
- Freudenheim JL, Ritz J, Smith-Warner SA *et al.* (2005). Alcohol consumption and risk of lung cancer: a pooled analysis of cohort studies. *Am J Clin Nutr*, 82:657–667. [PMID:16155281](#)
- Friberg E, Wolk A (2009). Long-term alcohol consumption and risk of endometrial cancer incidence: a prospective cohort study. *Cancer Epidemiol Biomarkers Prev*, 18:355–358. [doi:10.1158/1055-9965.EPI-08-0993](#) [PMID:19124521](#)
- Friberg JT, Yuan JM, Wang R *et al.* (2007). A prospective study of tobacco and alcohol use as risk factors for pharyngeal carcinomas in Singapore Chinese. *Cancer*, 109:1183–1191. [doi:10.1002/encr.22501](#) [PMID:17315158](#)
- Friedenreich CM, Howe GR, Miller AB, Jain MG (1993). A cohort study of alcohol consumption and risk of breast cancer. *Am J Epidemiol*, 137:512–520. [PMID:8465803](#)
- Fuchs CS, Stampfer MJ, Colditz GA *et al.* (1995). Alcohol consumption and mortality among women. *N Engl J Med*, 332:1245–1250. [doi:10.1056/NEJM199505113321901](#) [PMID:7708067](#)

- Gallus S, Foschi R, Talamini R *et al.* (2007). Risk factors for prostate cancer in men aged less than 60 years: a case-control study from Italy. *Urology*, 70:1121–1126. [doi:10.1016/j.urology.2007.07.020](https://doi.org/10.1016/j.urology.2007.07.020) [PMID:18158031](https://pubmed.ncbi.nlm.nih.gov/18158031/)
- Ganesh B, Talole SD, Dikshit R (2009). A case-control study on diet and colorectal cancer from Mumbai, India. *Cancer Epidemiol*, 33:189–193. [doi:10.1016/j.canep.2009.07.009](https://doi.org/10.1016/j.canep.2009.07.009) [PMID:19717354](https://pubmed.ncbi.nlm.nih.gov/19717354/)
- Gao C, Takezaki T, Wu J *et al.* (2002). Interaction between cytochrome P-450 2E1 polymorphisms and environmental factors with risk of esophageal and stomach cancers in Chinese. *Cancer Epidemiol Biomarkers Prev*, 11:29–34. [PMID:11815398](https://pubmed.ncbi.nlm.nih.gov/11815398/)
- Gao CM, Takezaki T, Wu JZ *et al.* (2007). CYP2E1 Rsa I polymorphism impacts on risk of colorectal cancer association with smoking and alcohol drinking. *World J Gastroenterol*, 13:5725–5730. [PMID:17963298](https://pubmed.ncbi.nlm.nih.gov/17963298/)
- Gao CM, Takezaki T, Wu JZ *et al.* (2008). Polymorphisms of alcohol dehydrogenase 2 and aldehyde dehydrogenase 2 and colorectal cancer risk in Chinese males. *World J Gastroenterol*, 14:5078–5083. [doi:10.3748/wjg.14.5078](https://doi.org/10.3748/wjg.14.5078) [PMID:18763293](https://pubmed.ncbi.nlm.nih.gov/18763293/)
- Gapstur SM, Potter JD, Sellers TA *et al.* (1993). Alcohol consumption and postmenopausal endometrial cancer: results from the Iowa Women's Health Study. *Cancer Causes Control*, 4:323–329. [doi:10.1007/BF00051334](https://doi.org/10.1007/BF00051334) [PMID:8347781](https://pubmed.ncbi.nlm.nih.gov/8347781/)
- Garavello W, Bosetti C, Gallus S *et al.* (2006). Type of alcoholic beverage and the risk of laryngeal cancer. *Eur J Cancer Prev*, 15:69–73. [doi:10.1097/01.cej.0000186641.19872.04](https://doi.org/10.1097/01.cej.0000186641.19872.04) [PMID:16374233](https://pubmed.ncbi.nlm.nih.gov/16374233/)
- Garfinkel L, Boffetta P, Stellman SD (1988). Alcohol and breast cancer: a cohort study. *Prev Med*, 17:686–693. [doi:10.1016/0091-7435\(88\)90086-2](https://doi.org/10.1016/0091-7435(88)90086-2) [PMID:3244667](https://pubmed.ncbi.nlm.nih.gov/3244667/)
- Garland C, Shekelle RB, Barrett-Connor E *et al.* (1985). Dietary vitamin D and calcium and risk of colorectal cancer: a 19-year prospective study in men. *Lancet*, 1:307–309. [doi:10.1016/S0140-6736\(85\)91082-7](https://doi.org/10.1016/S0140-6736(85)91082-7) [PMID:2857364](https://pubmed.ncbi.nlm.nih.gov/2857364/)
- Garland M, Hunter DJ, Colditz GA *et al.* (1999). Alcohol consumption in relation to breast cancer risk in a cohort of United States women 25–42 years of age. *Cancer Epidemiol Biomarkers Prev*, 8:1017–1021. [PMID:10566558](https://pubmed.ncbi.nlm.nih.gov/10566558/)
- Gattás GJF, de Carvalho MB, Siraque MS *et al.* (2006). Genetic polymorphisms of CYP1A1, CYP2E1, GSTM1, and GSTT1 associated with head and neck cancer. *Head Neck*, 28:819–826. [doi:10.1002/hed.20410](https://doi.org/10.1002/hed.20410) [PMID:16721740](https://pubmed.ncbi.nlm.nih.gov/16721740/)
- Genkinger JM, Spiegelman D, Anderson KE *et al.* (2009). Alcohol intake and pancreatic cancer risk: a pooled analysis of fourteen cohort studies. *Cancer Epidemiol Biomarkers Prev*, 18:765–776. [doi:10.1158/1055-9965.EPI-08-0880](https://doi.org/10.1158/1055-9965.EPI-08-0880) [PMID:19258474](https://pubmed.ncbi.nlm.nih.gov/19258474/)
- Gibson LJ, Hery C, Mitton N *et al.* (2009). Risk factors for breast cancer among Filipino women in Manila. *Int J Cancer*. [PMID:19626603](https://pubmed.ncbi.nlm.nih.gov/19626603/)
- Giovannucci E, Chen J, Smith-Warner SA *et al.* (2003). Methylene tetrahydrofolate reductase, alcohol dehydrogenase, diet, and risk of colorectal adenomas. *Cancer Epidemiol Biomarkers Prev*, 12:970–979. [PMID:14578131](https://pubmed.ncbi.nlm.nih.gov/14578131/)
- Giovannucci E, Rimm EB, Ascherio A *et al.* (1995). Alcohol, low-methionine–low-folate diets, and risk of colon cancer in men. *J Natl Cancer Inst*, 87:265–273. [doi:10.1093/jnci/87.4.265](https://doi.org/10.1093/jnci/87.4.265) [PMID:7707417](https://pubmed.ncbi.nlm.nih.gov/7707417/)
- Glynn SA, Albanes D, Pietinen P *et al.* (1996). Alcohol consumption and risk of colorectal cancer in a cohort of Finnish men. *Cancer Causes Control*, 7:214–223. [doi:10.1007/BF00051297](https://doi.org/10.1007/BF00051297) [PMID:8740734](https://pubmed.ncbi.nlm.nih.gov/8740734/)
- Goldbohm RA, Van den Brandt PA, Van 't Veer P *et al.* (1994). Prospective study on alcohol consumption and the risk of cancer of the colon and rectum in the Netherlands. *Cancer Causes Control*, 5:95–104. [doi:10.1007/BF01830255](https://doi.org/10.1007/BF01830255) [PMID:8167268](https://pubmed.ncbi.nlm.nih.gov/8167268/)

- Gong Z, Kristal AR, Schenk JM *et al.* (2009). Alcohol consumption, finasteride, and prostate cancer risk: results from the Prostate Cancer Prevention Trial. *Cancer*, 115:3661–3669. [doi:10.1002/ncr.24423](https://doi.org/10.1002/ncr.24423) [PMID:19598210](https://pubmed.ncbi.nlm.nih.gov/19598210/)
- González MV, Alvarez V, Pello MF *et al.* (1998). Genetic polymorphism of N-acetyltransferase-2, glutathione S-transferase-M1, and cytochromes P450IIE1 and P450IID6 in the susceptibility to head and neck cancer. *J Clin Pathol*, 51:294–298. [doi:10.1136/jcp.51.4.294](https://doi.org/10.1136/jcp.51.4.294) [PMID:9659241](https://pubmed.ncbi.nlm.nih.gov/9659241/)
- Goodman MT, Cologne JB, Moriwaki H *et al.* (1997a). Risk factors for primary breast cancer in Japan: 8-year follow-up of atomic bomb survivors. *Prev Med*, 26:144–153. [doi:10.1006/pmed.1996.9979](https://doi.org/10.1006/pmed.1996.9979) [PMID:9010910](https://pubmed.ncbi.nlm.nih.gov/9010910/)
- Gordon T, Kannel WB (1984). Drinking and mortality. The Framingham Study. *Am J Epidemiol*, 120:97–107. [PMID:6741928](https://pubmed.ncbi.nlm.nih.gov/6741928/)
- Gorini G, Stagnaro E, Fontana V *et al.* (2007). Alcohol consumption and risk of Hodgkin's lymphoma and multiple myeloma: a multicentre case-control study. *Ann Oncol*, 18:143–148. [doi:10.1093/annonc/mdl352](https://doi.org/10.1093/annonc/mdl352) [PMID:17047000](https://pubmed.ncbi.nlm.nih.gov/17047000/)
- Greving JP, Lee JE, Wolk A *et al.* (2007). Alcoholic beverages and risk of renal cell cancer. *Br J Cancer*, 97:429–433. [doi:10.1038/sj.bjc.6603890](https://doi.org/10.1038/sj.bjc.6603890) [PMID:17653076](https://pubmed.ncbi.nlm.nih.gov/17653076/)
- Grönberg H, Damber L, Damber J-E (1996). Total food consumption and body mass index in relation to prostate cancer risk: a case-control study in Sweden with prospectively collected exposure data. *J Urol*, 155:969–974. [doi:10.1016/S0022-5347\(01\)66360-2](https://doi.org/10.1016/S0022-5347(01)66360-2) [PMID:8583620](https://pubmed.ncbi.nlm.nih.gov/8583620/)
- Guo W, Blot WJ, Li J-Y *et al.* (1994). A nested case-control study of oesophageal and stomach cancers in the Linxian nutrition intervention trial. *Int J Epidemiol*, 23:444–450. [doi:10.1093/ije/23.3.444](https://doi.org/10.1093/ije/23.3.444) [PMID:7960367](https://pubmed.ncbi.nlm.nih.gov/7960367/)
- Guo YM, Wang Q, Liu YZ *et al.* (2008). Genetic polymorphisms in cytochrome P4502E1, alcohol and aldehyde dehydrogenases and the risk of esophageal squamous cell carcinoma in Gansu Chinese males. *World J Gastroenterol*, 14:1444–1449. [doi:10.3748/wjg.14.1444](https://doi.org/10.3748/wjg.14.1444) [PMID:18322963](https://pubmed.ncbi.nlm.nih.gov/18322963/)
- Gwack J, Hwang SS, Ko KP *et al.* (2007). [Fasting serum glucose and subsequent liver cancer risk in a Korean prospective cohort]. *J Prev Med Pub Health*, 40:23–28. [doi:10.3961/jpmph.2007.40.1.23](https://doi.org/10.3961/jpmph.2007.40.1.23) [PMID:17310595](https://pubmed.ncbi.nlm.nih.gov/17310595/)
- Gyntelberg F (1973). Physical fitness and coronary heart disease in male residents in Copenhagen aged 40–59. *Dan Med Bull*, 20:1–4. [PMID:4694991](https://pubmed.ncbi.nlm.nih.gov/4694991/)
- Hakulinen T, Lehtimäki L, Lehtonen M, Teppo L (1974). Cancer morbidity among two male cohorts with increased alcohol consumption in Finland. *J Natl Cancer Inst*, 52:1711–1714. [PMID:4834405](https://pubmed.ncbi.nlm.nih.gov/4834405/)
- Hamada GS, Sugimura H, Suzuki I *et al.* (1995). The heme-binding region polymorphism of cytochrome P4501A1 (*Cyp1A1*), rather than the *RsaI* polymorphism of IIE1 (*CypIIE1*), is associated with lung cancer in Rio de Janeiro. *Cancer Epidemiol. Biomarkers Prev.*, 4:63–67.
- Harnack L, Jacobs DR Jr, Nicodemus K *et al.* (2002). Relationship of folate, vitamin B-6, vitamin B-12, and methionine intake to incidence of colorectal cancers. *Nutr Cancer*, 43:152–158. [doi:10.1207/S15327914NC432\\_5](https://doi.org/10.1207/S15327914NC432_5) [PMID:12588695](https://pubmed.ncbi.nlm.nih.gov/12588695/)
- Harnack LJ, Anderson KE, Zheng W *et al.* (1997). Smoking, alcohol, coffee, and tea intake and incidence of cancer of the exocrine pancreas: the Iowa Women's Health Study. *Cancer Epidemiol Biomarkers Prev*, 6:1081–1086. [PMID:9419407](https://pubmed.ncbi.nlm.nih.gov/9419407/)
- Harty LC, Caporaso NE, Hayes RB *et al.* (1997). Alcohol dehydrogenase 3 genotype and risk of oral cavity and pharyngeal cancers. *J Natl Cancer Inst*, 89:1698–1705. [doi:10.1093/jnci/89.22.1698](https://doi.org/10.1093/jnci/89.22.1698) [PMID:9390539](https://pubmed.ncbi.nlm.nih.gov/9390539/)

- Hashibe M, Boffetta P, Zaridze D *et al.* (2006). Evidence for an important role of alcohol- and aldehyde-metabolizing genes in cancers of the upper aerodigestive tract. *Cancer Epidemiol Biomarkers Prev*, 15:696–703. [doi:10.1158/1055-9965.EPI-05-0710](https://doi.org/10.1158/1055-9965.EPI-05-0710) PMID:16614111
- Hashibe M, Brennan P, Benhamou S *et al.* (2007). Alcohol drinking in never users of tobacco, cigarette smoking in never drinkers, and the risk of head and neck cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *J Natl Cancer Inst*, 99:777–789. [doi:10.1093/jnci/djk179](https://doi.org/10.1093/jnci/djk179) PMID:17505073
- Hashibe M, Brennan P, Chuang SC *et al.* (2009). Interaction between tobacco and alcohol use and the risk of head and neck cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *Cancer Epidemiol Biomarkers Prev*, 18:541–550. [doi:10.1158/1055-9965.EPI-08-0347](https://doi.org/10.1158/1055-9965.EPI-08-0347) PMID:19190158
- Hashibe M, McKay JD, Curado MP *et al.* (2008). Multiple ADH genes are associated with upper aerodigestive cancers. *Nat Genet*, 40:707–709. [doi:10.1038/ng.151](https://doi.org/10.1038/ng.151) PMID:18500343
- Hashimoto T, Uchida K, Okayama N *et al.* (2006). ALDH2 1510 G/A (Glu487Lys) polymorphism interaction with age in head and neck squamous cell carcinoma. *Tumour Biol*, 27:334–338. [doi:10.1159/000096126](https://doi.org/10.1159/000096126) PMID:17033202
- Hassan MM, Bondy ML, Wolff RA *et al.* (2007). Risk factors for pancreatic cancer: case-control study. *Am J Gastroenterol*, 102:2696–2707. [doi:10.1111/j.1572-0241.2007.01510.x](https://doi.org/10.1111/j.1572-0241.2007.01510.x) PMID:17764494
- Hassan MM, Spitz MR, Thomas MB *et al.* (2008). Effect of different types of smoking and synergism with hepatitis C virus on risk of hepatocellular carcinoma in American men and women: case-control study. *Int J Cancer*, 123:1883–1891. [doi:10.1002/ijc.23730](https://doi.org/10.1002/ijc.23730) PMID:18688864
- Hein HO, Suadicani P, Gyntelberg F (1992). Lung cancer risk and social class. The Copenhagen Male Study—17-year follow up. *Dan Med Bull*, 39:173–176. PMID:1611922
- Heinen MM, Verhage BA, Ambergen TA *et al.* (2009). Alcohol consumption and risk of pancreatic cancer in the Netherlands cohort study. *Am J Epidemiol*, 169:1233–1242. [doi:10.1093/aje/kwp028](https://doi.org/10.1093/aje/kwp028) PMID:19318612
- Heuch I, Kvåle G, Jacobsen BK, Bjelke E (1983). Use of alcohol, tobacco and coffee, and risk of pancreatic cancer. *Br J Cancer*, 48:637–643. [doi:10.1038/bjc.1983.245](https://doi.org/10.1038/bjc.1983.245) PMID:6685527
- Hiatt RA, Klatsky AL, Armstrong MA (1988). Pancreatic cancer, blood glucose and beverage consumption. *Int J Cancer*, 41:794–797. [doi:10.1002/ijc.2910410603](https://doi.org/10.1002/ijc.2910410603) PMID:3372055
- Hiatt RA, Armstrong MA, Klatsky AL, Sidney S (1994). Alcohol consumption, smoking, and other risk factors and prostate cancer in a large health plan cohort in California (United States). *Cancer Causes Control*, 5:66–72. [doi:10.1007/BF01830728](https://doi.org/10.1007/BF01830728) PMID:7510134
- Hildesheim A, Anderson LM, Chen CJ *et al.* (1997). CYP2E1 genetic polymorphisms and risk of nasopharyngeal carcinoma in Taiwan. *J Natl Cancer Inst*, 89:1207–1212. [doi:10.1093/jnci/89.16.1207](https://doi.org/10.1093/jnci/89.16.1207) PMID:9274915
- Hildesheim A, Chen CJ, Caporaso NE *et al.* (1995). Cytochrome P4502E1 genetic polymorphisms and risk of nasopharyngeal carcinoma: results from a case-control study conducted in Taiwan. *Cancer Epidemiol Biomarkers Prev*, 4:607–610. PMID:8547826
- Hines LM, Hankinson SE, Smith-Warner SA *et al.* (2000). A prospective study of the effect of alcohol consumption and ADH3 genotype on plasma steroid hormone levels and breast cancer risk. *Cancer Epidemiol Biomarkers Prev*, 9:1099–1105. PMID:11045794
- Hiraki A, Matsuo K, Wakai K *et al.* (2007). Gene-gene and gene-environment interactions between alcohol drinking habit and polymorphisms in alcohol-metabolizing enzyme genes and the risk of head and neck cancer in Japan. *Cancer Sci*, 98:1087–1091. [doi:10.1111/j.1349-7006.2007.00505.x](https://doi.org/10.1111/j.1349-7006.2007.00505.x) PMID:17489985

- Hirayama T (1989). Association between alcohol consumption and cancer of the sigmoid colon: observations from a Japanese cohort study. *Lancet*, 2:725–727. [doi:10.1016/S0140-6736\(89\)90782-4](https://doi.org/10.1016/S0140-6736(89)90782-4) [PMID:2570969](https://pubmed.ncbi.nlm.nih.gov/2570969/)
- Hirayama T (1992). Life-style and cancer: from epidemiological evidence to public behavior change to mortality reduction of target cancers. *J Natl Cancer Inst Monogr*, 12:65–74. [PMID:1616813](https://pubmed.ncbi.nlm.nih.gov/1616813/)
- Hirose M, Kono S, Tabata S *et al.* (2005). Genetic polymorphisms of methylenetetrahydrofolate reductase and aldehyde dehydrogenase 2, alcohol use and risk of colorectal adenomas: Self-Defense Forces Health Study. *Cancer Sci*, 96:513–518. [doi:10.1111/j.1349-7006.2005.00077.x](https://doi.org/10.1111/j.1349-7006.2005.00077.x) [PMID:16108833](https://pubmed.ncbi.nlm.nih.gov/16108833/)
- Hirvonen A, Husgafvel-Pursiainen K, Anttila S *et al.* (1993). The human CYP2E1 gene and lung cancer: DraI and RsaI restriction fragment length polymorphisms in a Finnish study population. *Carcinogenesis*, 14:85–88. [doi:10.1093/carcin/14.1.85](https://doi.org/10.1093/carcin/14.1.85) [PMID:8093864](https://pubmed.ncbi.nlm.nih.gov/8093864/)
- Hirvonen T, Mennen LI, de Bree A *et al.* (2006). Consumption of antioxidant-rich beverages and risk for breast cancer in French women. *Ann Epidemiol*, 16:503–508. [doi:10.1016/j.annepidem.2005.09.011](https://doi.org/10.1016/j.annepidem.2005.09.011) [PMID:16406814](https://pubmed.ncbi.nlm.nih.gov/16406814/)
- Holmberg L, Baron JA, Byers T *et al.* (1995). Alcohol intake and breast cancer risk: effect of exposure from 15 years of age. *Cancer Epidemiol Biomarkers Prev*, 4:843–847. [PMID:8634655](https://pubmed.ncbi.nlm.nih.gov/8634655/)
- Homann N, König IR, Marks M *et al.* (2009). Alcohol and colorectal cancer: the role of alcohol dehydrogenase 1C polymorphism. *Alcohol Clin Exp Res*, 33:551–556. [doi:10.1111/j.1530-0277.2008.00868.x](https://doi.org/10.1111/j.1530-0277.2008.00868.x) [PMID:19120062](https://pubmed.ncbi.nlm.nih.gov/19120062/)
- Homann N, Stickel F, König IR *et al.* (2006). Alcohol dehydrogenase 1C\*1 allele is a genetic marker for alcohol-associated cancer in heavy drinkers. *Int J Cancer*, 118:1998–2002. [doi:10.1002/ijc.21583](https://doi.org/10.1002/ijc.21583) [PMID:16287084](https://pubmed.ncbi.nlm.nih.gov/16287084/)
- Honjo S, Srivatanakul P, Sriplung H *et al.* (2005). Genetic and environmental determinants of risk for cholangiocarcinoma via *Opisthorchis viverrini* in a densely infested area in Nakhon Phanom, northeast Thailand. *Int J Cancer*, 117:854–860. [doi:10.1002/ijc.21146](https://doi.org/10.1002/ijc.21146) [PMID:15957169](https://pubmed.ncbi.nlm.nih.gov/15957169/)
- Hori H, Kawano T, Endo M, Yuasa Y (1997). Genetic polymorphisms of tobacco- and alcohol-related metabolizing enzymes and human esophageal squamous cell carcinoma susceptibility. *J Clin Gastroenterol*, 25:568–575. [doi:10.1097/00004836-199712000-00003](https://doi.org/10.1097/00004836-199712000-00003) [PMID:9451664](https://pubmed.ncbi.nlm.nih.gov/9451664/)
- Horn-Ross PL, Canchola AJ, West DW *et al.* (2004). Patterns of alcohol consumption and breast cancer risk in the California Teachers Study cohort. *Cancer Epidemiol Biomarkers Prev*, 13:405–411. [PMID:15006916](https://pubmed.ncbi.nlm.nih.gov/15006916/)
- Hosgood HD 3rd, Baris D, Zahm SH *et al.* (2007). Diet and risk of multiple myeloma in Connecticut women. *Cancer Causes Control*, 18:1065–1076. [doi:10.1007/s10552-007-9047-z](https://doi.org/10.1007/s10552-007-9047-z) [PMID:17694422](https://pubmed.ncbi.nlm.nih.gov/17694422/)
- Hosono S, Matsuo K, Kajiyama H *et al.* (2008). Reduced risk of endometrial cancer from alcohol drinking in Japanese. *Cancer Sci*, 99:1195–1201. [doi:10.1111/j.1349-7006.2008.00801.x](https://doi.org/10.1111/j.1349-7006.2008.00801.x) [PMID:18422741](https://pubmed.ncbi.nlm.nih.gov/18422741/)
- Høyer AP, Engholm G (1992). Serum lipids and breast cancer risk: a cohort study of 5,207 Danish women. *Cancer Causes Control*, 3:403–408. [doi:10.1007/BF00051352](https://doi.org/10.1007/BF00051352) [PMID:1525320](https://pubmed.ncbi.nlm.nih.gov/1525320/)
- Hsing AW, McLaughlin JK, Chow WH *et al.* (1998a). Risk factors for colorectal cancer in a prospective study among U.S. white men. *Int J Cancer*, 77:549–553. [doi:10.1002/\(SICI\)1097-0215\(19980812\)77:4<549::AID-IJC13>3.0.CO;2-I](https://doi.org/10.1002/(SICI)1097-0215(19980812)77:4<549::AID-IJC13>3.0.CO;2-I) [PMID:9679757](https://pubmed.ncbi.nlm.nih.gov/9679757/)
- Hsing AW, McLaughlin JK, Schuman LM *et al.* (1990). Diet, tobacco use, and fatal prostate cancer: results from the Lutheran Brotherhood Cohort Study. *Cancer Res*, 50:6836–6840. [PMID:2208150](https://pubmed.ncbi.nlm.nih.gov/2208150/)
- Hsing AW, Zhang M, Rashid A *et al.* (2008). Hepatitis B and C virus infection and the risk of biliary tract cancer: a population-based study in China. *Int J Cancer*, 122:1849–1853. [doi:10.1002/ijc.23251](https://doi.org/10.1002/ijc.23251) [PMID:18076042](https://pubmed.ncbi.nlm.nih.gov/18076042/)

- Hung HC, Chuang J, Chien YC *et al.* (1997). Genetic polymorphisms of CYP2E1, GSTM1, and GSTT1; Environmental factors and risk of oral cancer. *Cancer Epidemiol. Biomarkers Prev.*, 6:901–905.
- Huxley RR, Ansary-Moghaddam A, Clifton P *et al.* (2009). The impact of dietary and lifestyle risk factors on risk of colorectal cancer: a quantitative overview of the epidemiological evidence. *Int J Cancer*, 125:171–180. [doi:10.1002/ijc.24343](https://doi.org/10.1002/ijc.24343) [PMID:19350627](https://pubmed.ncbi.nlm.nih.gov/19350627/)
- Ide R, Mizoue T, Fujino Y *et al.*; JACC Study Group (2008). Cigarette smoking, alcohol drinking, and oral and pharyngeal cancer mortality in Japan. *Oral Dis*, 14:314–319. [doi:10.1111/j.1601-0825.2007.01378.x](https://doi.org/10.1111/j.1601-0825.2007.01378.x) [PMID:18449960](https://pubmed.ncbi.nlm.nih.gov/18449960/)
- Inoue M, Tajima K, Takezaki T *et al.* (2003). Epidemiology of pancreatic cancer in Japan: a nested case-control study from the Hospital-based Epidemiologic Research Program at Aichi Cancer Center (HERPACC). *Int J Epidemiol*, 32:257–262. [doi:10.1093/ije/dyg062](https://doi.org/10.1093/ije/dyg062) [PMID:12714546](https://pubmed.ncbi.nlm.nih.gov/12714546/)
- Iribarren C, Haselkorn T, Tekawa IS, Friedman GD (2001). Cohort study of thyroid cancer in a San Francisco Bay area population. *Int J Cancer*, 93:745–750. [doi:10.1002/ijc.1377](https://doi.org/10.1002/ijc.1377) [PMID:11477590](https://pubmed.ncbi.nlm.nih.gov/11477590/)
- Isaksson B, Jonsson F, Pedersen NL *et al.* (2002). Lifestyle factors and pancreatic cancer risk: a cohort study from the Swedish Twin Registry. *Int J Cancer*, 98:480–482. [doi:10.1002/ijc.10256](https://doi.org/10.1002/ijc.10256) [PMID:11920604](https://pubmed.ncbi.nlm.nih.gov/11920604/)
- Ishiguro S, Sasazuki S, Inoue M *et al.*; JPHC Study Group (2009). Effect of alcohol consumption, cigarette smoking and flushing response on esophageal cancer risk: a population-based cohort study (JPHC study). *Cancer Lett*, 275:240–246. [doi:10.1016/j.canlet.2008.10.020](https://doi.org/10.1016/j.canlet.2008.10.020) [PMID:19036500](https://pubmed.ncbi.nlm.nih.gov/19036500/)
- Ishihara J, Otani T, Inoue M *et al.*; Japan Public Health Center-based Prospective Study Group (2007). Low intake of vitamin B-6 is associated with increased risk of colorectal cancer in Japanese men. *J Nutr*, 137:1808–1814. [PMID:17585035](https://pubmed.ncbi.nlm.nih.gov/17585035/)
- Jahnke V, Matthias C, Fryer A, Strange R (1996). Glutathione S-transferase and cytochrome-P-450 polymorphism as risk factors for squamous cell carcinoma of the larynx. *Am J Surg*, 172:671–673. [doi:10.1016/S0002-9610\(96\)00298-X](https://doi.org/10.1016/S0002-9610(96)00298-X) [PMID:8988674](https://pubmed.ncbi.nlm.nih.gov/8988674/)
- Jain MG, Ferrence RG, Rehm JT *et al.* (2000a). Alcohol and breast cancer mortality in a cohort study. *Breast Cancer Res Treat*, 64:201–209. [doi:10.1023/A:1006402323445](https://doi.org/10.1023/A:1006402323445) [PMID:11194456](https://pubmed.ncbi.nlm.nih.gov/11194456/)
- Jain MG, Rohan TE, Howe GR, Miller AB (2000b). A cohort study of nutritional factors and endometrial cancer. *Eur J Epidemiol*, 16:899–905. [doi:10.1023/A:1011012621990](https://doi.org/10.1023/A:1011012621990) [PMID:11338120](https://pubmed.ncbi.nlm.nih.gov/11338120/)
- Jensen OM (1979). Cancer morbidity and causes of death among Danish brewery workers. *Int J Cancer*, 23:454–463. [doi:10.1002/ijc.2910230404](https://doi.org/10.1002/ijc.2910230404) [PMID:437924](https://pubmed.ncbi.nlm.nih.gov/437924/)
- Jiang X, Castela JE, Groshen S *et al.* (2007). Alcohol consumption and risk of bladder cancer in Los Angeles County. *Int J Cancer*, 121:839–845. [doi:10.1002/ijc.22743](https://doi.org/10.1002/ijc.22743) [PMID:17440923](https://pubmed.ncbi.nlm.nih.gov/17440923/)
- Jiao L, Silverman DT, Schairer C *et al.* (2009). Alcohol use and risk of pancreatic cancer: the NIH-AARP Diet and Health Study. *Am J Epidemiol*, 169:1043–1051. [doi:10.1093/aje/kwp034](https://doi.org/10.1093/aje/kwp034) [PMID:19299403](https://pubmed.ncbi.nlm.nih.gov/19299403/)
- Johansen D, Borgström A, Lindkvist B, Manjer J (2009). Different markers of alcohol consumption, smoking and body mass index in relation to risk of pancreatic cancer. A prospective cohort study within the Malmö Preventive Project. *Pancreatology*, 9:677–686. [doi:10.1159/000212088](https://doi.org/10.1159/000212088) [PMID:19684432](https://pubmed.ncbi.nlm.nih.gov/19684432/)
- Jung AY, Poole EM, Bigler J *et al.* (2008). DNA methyltransferase and alcohol dehydrogenase: gene-nutrient interactions in relation to risk of colorectal polyps. *Cancer Epidemiol Biomarkers Prev*, 17:330–338. [doi:10.1158/1055-9965.EPI-07-2608](https://doi.org/10.1158/1055-9965.EPI-07-2608) [PMID:18268116](https://pubmed.ncbi.nlm.nih.gov/18268116/)
- Kabat GC, Miller AB, Jain M, Rohan TE (2008). Dietary intake of selected B vitamins in relation to risk of major cancers in women. *Br J Cancer*, 99:816–821. [doi:10.1038/sj.bjc.6604540](https://doi.org/10.1038/sj.bjc.6604540) [PMID:18665162](https://pubmed.ncbi.nlm.nih.gov/18665162/)

- Kanda J, Matsuo K, Suzuki T *et al.* (2009). Impact of alcohol consumption with polymorphisms in alcohol-metabolizing enzymes on pancreatic cancer risk in Japanese. *Cancer Sci*, 100:296–302. [doi:10.1111/j.1349-7006.2008.01044.x](https://doi.org/10.1111/j.1349-7006.2008.01044.x) PMID:19068087
- Karlson B-M, Ekblom A, Josefsson S *et al.* (1997). The risk of pancreatic cancer following pancreatitis: an association due to confounding? *Gastroenterology*, 113:587–592. [doi:10.1053/gast.1997.v113.pm9247480](https://doi.org/10.1053/gast.1997.v113.pm9247480) PMID:9247480
- Kato I, Nomura AMY, Stemmermann GN, Chyou PH (1992c). Prospective study of the association of alcohol with cancer of the upper aerodigestive tract and other sites. *Cancer Causes Control*, 3:145–151. [doi:10.1007/BF00051654](https://doi.org/10.1007/BF00051654) PMID:1562704
- Kato I, Tominaga S, Ito Y *et al.* (1992a). A prospective study of atrophic gastritis and stomach cancer risk. *Jpn J Cancer Res*, 83:1137–1142. [doi:10.1111/j.1349-7006.1992.tb02736.x](https://doi.org/10.1111/j.1349-7006.1992.tb02736.x) PMID:1483928
- Kato I, Tominaga S, Matsumoto K (1992b). A prospective study of stomach cancer among a rural Japanese population: a 6-year survey. *Jpn J Cancer Res*, 83:568–575. [doi:10.1111/j.1349-7006.1992.tb00127.x](https://doi.org/10.1111/j.1349-7006.1992.tb00127.x) PMID:1644660
- Kato S, Onda M, Matsukura N *et al.* (1995). Cytochrome P4502E1 (CYP2E1) genetic polymorphism in a case-control study of gastric cancer and liver disease. *Pharmacogenetics*, 5 Special Issue;S141–S144. [doi:10.1097/00008571-199512001-00016](https://doi.org/10.1097/00008571-199512001-00016) PMID:7581484
- Kato S, Onda M, Matsukura N *et al.* (1996). Genetic polymorphisms of the cancer related gene and Helicobacter pylori infection in Japanese gastric cancer patients. An age and gender matched case-control study. *Cancer*, 77 Suppl;1654–1661. PMID:8608558
- Kato S, Onda M, Matsukura N *et al.* (1997). Helicobacter pylori infection and genetic polymorphisms for cancer-related genes in gastric carcinogenesis. *Biomed Pharmacother*, 51:145–149. [doi:10.1016/S0753-3322\(97\)85581-3](https://doi.org/10.1016/S0753-3322(97)85581-3) PMID:9207980
- Kato S, Shields PG, Caporaso NE *et al.* (1992). Cytochrome P450IIE1 genetic polymorphisms, racial variation, and lung cancer risk. *Cancer Res*, 52:6712–6715. PMID:1423319
- Kato S, Shields PG, Caporaso NE *et al.* (1994). Analysis of cytochrome P450 2E1 genetic polymorphisms in relation to human lung cancer. *Cancer Epidemiol Biomarkers Prev*, 3:515–518. PMID:8000304
- Kato S, Tajiri T, Matsukura N *et al.* (2003). Genetic polymorphisms of aldehyde dehydrogenase 2, cytochrome p450 2E1 for liver cancer risk in HCV antibody-positive japanese patients and the variations of CYP2E1 mRNA expression levels in the liver due to its polymorphism. *Scand J Gastroenterol*, 38:886–893. [doi:10.1080/00365520310004489](https://doi.org/10.1080/00365520310004489) PMID:12940444
- Katoh T, Kaneko S, Kohshi K *et al.* (1999). Genetic polymorphisms of tobacco- and alcohol-related metabolizing enzymes and oral cavity cancer. *Int J Cancer*, 83:606–609. [doi:10.1002/\(SICI\)1097-0215\(19991126\)83:5<606::AID-IJC6>3.0.CO;2-P](https://doi.org/10.1002/(SICI)1097-0215(19991126)83:5<606::AID-IJC6>3.0.CO;2-P) PMID:10521794
- Kawase T, Matsuo K, Hiraki A *et al.* (2009). Interaction of the effects of alcohol drinking and polymorphisms in alcohol-metabolizing enzymes on the risk of female breast cancer in Japan. *J Epidemiol*, 19:244–250. [doi:10.2188/jea.JE20081035](https://doi.org/10.2188/jea.JE20081035) PMID:19667493
- Kelemen LE, Sellers TA, Vierkant RA *et al.* (2004). Association of folate and alcohol with risk of ovarian cancer in a prospective study of postmenopausal women. *Cancer Causes Control*, 15:1085–1093. [doi:10.1007/s10552-004-1546-6](https://doi.org/10.1007/s10552-004-1546-6) PMID:15801492
- Khurana V, Sheth A, Caldito G, Barkin JS (2007). Statins reduce the risk of pancreatic cancer in humans: a case-control study of half a million veterans. *Pancreas*, 34:260–265. [doi:10.1097/MPA.0b013e318030e963](https://doi.org/10.1097/MPA.0b013e318030e963) PMID:17312467
- Kinjo Y, Cui Y, Akiba S *et al.* (1998). Mortality risks of oesophageal cancer associated with hot tea, alcohol, tobacco and diet in Japan. *J Epidemiol*, 8:235–243. [doi:10.2188/jea.8.235](https://doi.org/10.2188/jea.8.235) PMID:9816815

- Kiss I, Sándor J, Pajkos G *et al.* (2000). Colorectal cancer risk in relation to genetic polymorphism of cytochrome P450 1A1, 2E1, and glutathione-S-transferase M1 enzymes. *Anticancer Res*, 20 1B;519–522. [PMID:10769717](#)
- Kjaerheim K, Andersen A, Helseth A (1993). Alcohol abstainers: a low-risk group for cancer—a cohort study of Norwegian teetotalers. *Cancer Epidemiol Biomarkers Prev*, 2:93–97. [PMID:8467252](#)
- Kjaerheim K, Andersen A (1994). Cancer incidence among waitresses in Norway. *Cancer Causes Control*, 5:31–37. [doi:10.1007/BF01830724](#) [PMID:8123777](#)
- Kjaerheim K, Gaard M, Andersen A (1998). The role of alcohol, tobacco, and dietary factors in upper aerogastric tract cancers: a prospective study of 10,900 Norwegian men. *Cancer Causes Control*, 9:99–108. [doi:10.1023/A:1008809706062](#) [PMID:9486469](#)
- Klatsky AL, Armstrong MA, Friedman GD, Hiatt RA (1988). The relations of alcoholic beverage use to colon and rectal cancer. *Am J Epidemiol*, 128:1007–1015. [PMID:3189277](#)
- Klatsky AL, Friedman GD, Siegelau AB (1981). Alcohol and mortality. A ten-year Kaiser-Permanente experience. *Ann Intern Med*, 95:139–145. [PMID:7258861](#)
- Klatsky AL, Li Y, Baer D *et al.* (2009). Alcohol consumption and risk of hematologic malignancies. *Ann Epidemiol*, 19:746–753. [doi:10.1016/j.annepidem.2009.03.005](#) [PMID:19394864](#)
- Kneller RW, McLaughlin JK, Bjelke E *et al.* (1991). A cohort study of stomach cancer in a high-risk American population. *Cancer*, 68:672–678. [doi:10.1002/1097-0142\(19910801\)68:3<672::AID-CNCR2820680339>3.0.CO;2-T](#) [PMID:2065291](#)
- Knight JA, Bernstein L, Largent J *et al.*; WECARE Study Collaborative Group (2009). Alcohol intake and cigarette smoking and risk of a contralateral breast cancer: The Women’s Environmental Cancer and Radiation Epidemiology Study. *Am J Epidemiol*, 169:962–968. [doi:10.1093/aje/kwn422](#) [PMID:19211621](#)
- Kocić B, Petrovic B, Filipovic S (2008). Risk factors for breast cancer: a hospital-based case-control study. *J BUON*, 13:231–234. [PMID:18555470](#)
- Koide T, Ohno T, Huang XE *et al.* (2000). HBV/HCV infection, alcohol, tobacco and genetic polymorphisms for hepatocellular carcinoma in Nagoya, Japan. *Asian Pac J Cancer Prev*, 1:237–243. [PMID:12718671](#)
- Kolahdooz F, Ibiebele TI, van der Pols JC, Webb PM (2009). Dietary patterns and ovarian cancer risk. *Am J Clin Nutr*, 89:297–304. [doi:10.3945/ajcn.2008.26575](#) [PMID:19056595](#)
- Kono S, Ikeda M, Tokudome S *et al.* (1985). Alcohol and cancer in male Japanese physicians. *J Cancer Res Clin Oncol*, 109:82–85. [doi:10.1007/BF01884260](#) [PMID:3972887](#)
- Kono S, Ikeda M, Tokudome S *et al.* (1986). Alcohol and mortality: a cohort study of male Japanese physicians. *Int J Epidemiol*, 15:527–532. [doi:10.1093/ije/15.4.527](#) [PMID:3818161](#)
- Kono S, Ikeda M, Tokudome S *et al.* (1987). Cigarette smoking, alcohol and cancer mortality: a cohort study of male Japanese physicians. *Jpn J Cancer Res*, 78:1323–1328. [PMID:3123436](#)
- Korte JE, Brennan P, Henley SJ, Boffetta P (2002). Dose-specific meta-analysis and sensitivity analysis of the relation between alcohol consumption and lung cancer risk. *Am J Epidemiol*, 155:496–506. [doi:10.1093/aje/155.6.496](#) [PMID:11882523](#)
- Kruk J (2007). Association of lifestyle and other risk factors with breast cancer according to menopausal status: a case-control study in the Region of Western Pomerania (Poland). *Asian Pac J Cancer Prev*, 8:513–524. [PMID:18260721](#)
- Kubík A, Zatloukal P, Tomásek L *et al.* (2004). Lung cancer risk among nonsmoking women in relation to diet and physical activity. *Neoplasma*, 51:136–143. [PMID:15190423](#)

- Kuper H, Ye W, Weiderpass E *et al.* (2000). Alcohol and breast cancer risk: the alcoholism paradox. *Br J Cancer*, 83:949–951. [doi:10.1054/bjoc.2000.1360](https://doi.org/10.1054/bjoc.2000.1360) PMID:10970699
- Kushi LH, Mink PJ, Folsom AR *et al.* (1999). Prospective study of diet and ovarian cancer. *Am J Epidemiol*, 149:21–31. [PMID:9883790](https://pubmed.ncbi.nlm.nih.gov/9883790/)
- Kwan ML, Kushi LH, Weltzien E *et al.* (2009). Epidemiology of breast cancer subtypes in two prospective cohort studies of breast cancer survivors. *Breast Cancer Res*, 11:R31. [doi:10.1186/bcr2261](https://doi.org/10.1186/bcr2261) [PMID:19463150](https://pubmed.ncbi.nlm.nih.gov/19463150/)
- Ladero JM, Agúndez JAG, Rodríguez-Lescure A *et al.* (1996). RsaI polymorphism at the cytochrome P4502E1 locus and risk of hepatocellular carcinoma. *Gut*, 39:330–333. [doi:10.1136/gut.39.2.330](https://doi.org/10.1136/gut.39.2.330) [PMID:8977352](https://pubmed.ncbi.nlm.nih.gov/8977352/)
- Lagiou P, Ye W, Wedrén S *et al.* (2001). Incidence of ovarian cancer among alcoholic women: a cohort study in Sweden. *Int J Cancer*, 91:264–266. [doi:10.1002/1097-0215\(200002\)9999:9999<::AID-IJC1027>3.3.CO;2-B](https://doi.org/10.1002/1097-0215(200002)9999:9999<::AID-IJC1027>3.3.CO;2-B) [PMID:11146456](https://pubmed.ncbi.nlm.nih.gov/11146456/)
- Landi S, Gemignani F, Moreno V *et al.*; Bellvitge Colorectal Cancer Study Group (2005). A comprehensive analysis of phase I and phase II metabolism gene polymorphisms and risk of colorectal cancer. *Pharmacogenet Genomics*, 15:535–546. [doi:10.1097/01.fpc.0000165904.48994.3d](https://doi.org/10.1097/01.fpc.0000165904.48994.3d) [PMID:16006997](https://pubmed.ncbi.nlm.nih.gov/16006997/)
- Larsson SC, Giovannucci E, Wolk A (2007). Alcoholic beverage consumption and gastric cancer risk: a prospective population-based study in women. *Int J Cancer*, 120:373–377. [doi:10.1002/ijc.22204](https://doi.org/10.1002/ijc.22204) [PMID:17066442](https://pubmed.ncbi.nlm.nih.gov/17066442/)
- Le Marchand L, Donlon T, Seifried A, Wikens LR (2002). Red meat intake, CYP2E1 genetic polymorphisms, and colorectal cancer risk. *Cancer Epidemiol. Biomarkers Prev.*, 11:1019–1024.
- Le Marchand L, Kolonel LN, Wilkens LR *et al.* (1994). Animal fat consumption and prostate cancer: a prospective study in Hawaii. *Epidemiology*, 5:276–282. [doi:10.1097/00001648-199405000-00004](https://doi.org/10.1097/00001648-199405000-00004) [PMID:8038241](https://pubmed.ncbi.nlm.nih.gov/8038241/)
- Le Marchand L, Sivaraman L, Pierce L *et al.* (1998). Associations of CYP1A1, GSTM1, and CYP2E1 polymorphisms with lung cancer suggest cell type specificities to tobacco carcinogens. *Cancer Res*, 58:4858–4863. [PMID:9809991](https://pubmed.ncbi.nlm.nih.gov/9809991/)
- Lee HS, Yoon JH, Kamimura S *et al.* (1997a). Lack of association of cytochrome P450 2E1 genetic polymorphisms with the risk of human hepatocellular carcinoma. *Int J Cancer*, 71:737–740. [doi:10.1002/\(SICI\)1097-0215\(19970529\)71:5<737::AID-IJC8>3.0.CO;2-S](https://doi.org/10.1002/(SICI)1097-0215(19970529)71:5<737::AID-IJC8>3.0.CO;2-S) [PMID:9180139](https://pubmed.ncbi.nlm.nih.gov/9180139/)
- Lee CH, Lee JM, Wu DC *et al.* (2005). Independent and combined effects of alcohol intake, tobacco smoking and betel quid chewing on the risk of esophageal cancer in Taiwan. *Int J Cancer*, 113:475–482. [doi:10.1002/ijc.20619](https://doi.org/10.1002/ijc.20619) [PMID:15455377](https://pubmed.ncbi.nlm.nih.gov/15455377/)
- Lee HC, Yoon YB, Kim CY (1997b). Association between genetic polymorphisms of the cytochromes P-450 (1A1, 2D6, and 2E1) and the susceptibility to pancreatic cancer. *Korean J Intern Med*, 12:128–136. [PMID:9439147](https://pubmed.ncbi.nlm.nih.gov/9439147/)
- Lee JE, Giovannucci E, Smith-Warner SA *et al.* (2006). Total fluid intake and use of individual beverages and risk of renal cell cancer in two large cohorts. *Cancer Epidemiol Biomarkers Prev*, 15:1204–1211. [doi:10.1158/1055-9965.EPI-05-0889](https://doi.org/10.1158/1055-9965.EPI-05-0889) [PMID:16775182](https://pubmed.ncbi.nlm.nih.gov/16775182/)
- Lee TY, Lee SS, Jung SW *et al.* (2008). Hepatitis B virus infection and intrahepatic cholangiocarcinoma in Korea: a case-control study. *Am J Gastroenterol*, 103:1716–1720. [doi:10.1111/j.1572-0241.2008.01796.x](https://doi.org/10.1111/j.1572-0241.2008.01796.x) [PMID:18557716](https://pubmed.ncbi.nlm.nih.gov/18557716/)
- Lew JQ, Freedman ND, Leitzmann MF *et al.* (2009). Alcohol and risk of breast cancer by histologic type and hormone receptor status in postmenopausal women: the NIH-AARP Diet and Health Study. *Am J Epidemiol*, 170:308–317. [doi:10.1093/aje/kwp120](https://doi.org/10.1093/aje/kwp120) [PMID:19541857](https://pubmed.ncbi.nlm.nih.gov/19541857/)

- Li CI, Daling JR, Malone KE *et al.* (2006). Relationship between established breast cancer risk factors and risk of seven different histologic types of invasive breast cancer. *Cancer Epidemiol Biomarkers Prev*, 15:946–954. [doi:10.1158/1055-9965.EPI-05-0881](https://doi.org/10.1158/1055-9965.EPI-05-0881) PMID:16702375
- Li CI, Daling JR, Porter PL *et al.* (2009a). Relationship between potentially modifiable lifestyle factors and risk of second primary contralateral breast cancer among women diagnosed with estrogen receptor-positive invasive breast cancer. *J Clin Oncol*, 27:5312–5318. [doi:10.1200/JCO.2009.23.1597](https://doi.org/10.1200/JCO.2009.23.1597) PMID:19738113
- Li CI, Malone KE, Porter PL *et al.* (2003a). The relationship between alcohol use and risk of breast cancer by histology and hormone receptor status among women 65–79 years of age. *Cancer Epidemiol Biomarkers Prev*, 12:1061–1066. PMID:14578143
- Li CI, Malone KE, Porter PL, Daling JR (2003b). Epidemiologic and molecular risk factors for contralateral breast cancer among young women. *Br J Cancer*, 89:513–518. [doi:10.1038/sj.bjc.6601042](https://doi.org/10.1038/sj.bjc.6601042) PMID:12888823
- Li D, Ahmed M, Li Y *et al.* (2005). 5,10-Methylenetetrahydrofolate reductase polymorphisms and the risk of pancreatic cancer. *Cancer Epidemiol Biomarkers Prev*, 14:1470–1476. [doi:10.1158/1055-9965.EPI-04-0894](https://doi.org/10.1158/1055-9965.EPI-04-0894) PMID:15941958
- Li XM, Li J, Tsuji I *et al.* (2008). Mass screening-based case-control study of diet and prostate cancer in Changchun, China. *Asian J Androl*, 10:551–560. [doi:10.1111/j.1745-7262.2008.00384.x](https://doi.org/10.1111/j.1745-7262.2008.00384.x) PMID:18478158
- Li Y, Baer D, Friedman GD *et al.* (2009b). Wine, liquor, beer and risk of breast cancer in a large population. *Eur J Cancer*, 45:843–850. [doi:10.1016/j.ejca.2008.11.001](https://doi.org/10.1016/j.ejca.2008.11.001) PMID:19095438
- Lightfoot TJ, Barrett JH, Bishop T *et al.* (2008). Methylene tetrahydrofolate reductase genotype modifies the chemopreventive effect of folate in colorectal adenoma, but not colorectal cancer. *Cancer Epidemiol Biomarkers Prev*, 17:2421–2430. [doi:10.1158/1055-9965.EPI-08-0058](https://doi.org/10.1158/1055-9965.EPI-08-0058) PMID:18703816
- Lilla C, Koehler T, Kropp S *et al.* (2005). Alcohol dehydrogenase 1B (ADH1B) genotype, alcohol consumption and breast cancer risk by age 50 years in a German case-control study. *Br J Cancer*, 92:2039–2041. [doi:10.1038/sj.bjc.6602608](https://doi.org/10.1038/sj.bjc.6602608) PMID:15886702
- Lim HJ, Park BJ (2008). [Cohort study on the association between alcohol consumption and the risk of colorectal cancer in the Korean elderly]. *J Prev Med Pub Health*, 41:23–29. [doi:10.3961/jpmph.2008.41.1.23](https://doi.org/10.3961/jpmph.2008.41.1.23) PMID:18250602
- Lim U, Morton LM, Subar AF *et al.* (2007). Alcohol, smoking, and body size in relation to incident Hodgkin's and non-Hodgkin's lymphoma risk. *Am J Epidemiol*, 166:697–708. [doi:10.1093/aje/kwm122](https://doi.org/10.1093/aje/kwm122) PMID:17596266
- Lim U, Weinstein S, Albanes D *et al.* (2006). Dietary factors of one-carbon metabolism in relation to non-Hodgkin lymphoma and multiple myeloma in a cohort of male smokers. *Cancer Epidemiol Biomarkers Prev*, 15:1109–1114. [doi:10.1158/1055-9965.EPI-05-0918](https://doi.org/10.1158/1055-9965.EPI-05-0918) PMID:16775167
- Lin DX, Tang YM, Peng Q *et al.* (1998). Susceptibility to esophageal cancer and genetic polymorphisms in glutathione S-transferases T1, P1, and M1 and cytochrome P450 2E1. *Cancer Epidemiol Biomarkers Prev*, 7:1013–1018. PMID:9829710
- Lin Y, Kikuchi S, Tamakoshi K *et al.* (2005). Prospective study of alcohol consumption and breast cancer risk in Japanese women. *Int J Cancer*, 116:779–783. [doi:10.1002/ijc.20980](https://doi.org/10.1002/ijc.20980) PMID:15838830
- Lin Y, Tamakoshi A, Kawamura T *et al.* (2002). Risk of pancreatic cancer in relation to alcohol drinking, coffee consumption and medical history: findings from the Japan collaborative cohort study for evaluation of cancer risk. *Int J Cancer*, 99:742–746. [doi:10.1002/ijc.10402](https://doi.org/10.1002/ijc.10402) PMID:12115510
- Lindblad M, Rodríguez LAG, Lagergren J (2005). Body mass, tobacco and alcohol and risk of esophageal, gastric cardia, and gastric non-cardia adenocarcinoma among men and women in a nested case-control study. *Cancer Causes Control*, 16:285–294. [doi:10.1007/s10552-004-3485-7](https://doi.org/10.1007/s10552-004-3485-7) PMID:15947880

- Liu S, Park JY, Schantz SP *et al.* (2001). Elucidation of CYP2E1 5' regulatory RsaI/PstI allelic variants and their role in risk for oral cancer. *Oral Oncol*, 37:437–445. [doi:10.1016/S1368-8375\(00\)00099-3](https://doi.org/10.1016/S1368-8375(00)00099-3) [PMID:11377232](https://pubmed.ncbi.nlm.nih.gov/11377232/)
- Loerbroeks A, Schouten LJ, Goldbohm RA, van den Brandt PA (2007). Alcohol consumption, cigarette smoking, and endometrial cancer risk: results from the Netherlands Cohort Study. *Cancer Causes Control*, 18:551–560. [doi:10.1007/s10552-007-0127-x](https://doi.org/10.1007/s10552-007-0127-x) [PMID:17437180](https://pubmed.ncbi.nlm.nih.gov/17437180/)
- London SJ, Daly AK, Cooper J *et al.* (1996). Lung cancer risk in relation to the CYP2E1 Rsa I genetic polymorphism among African-Americans and Caucasians in Los Angeles County. *Pharmacogenetics*, 6:151–158. [doi:10.1097/00008571-199604000-00002](https://doi.org/10.1097/00008571-199604000-00002) [PMID:9156693](https://pubmed.ncbi.nlm.nih.gov/9156693/)
- Lu Q-J, Yao S-Y, Huang C-Y *et al.* (2000). [The cohort study on intake of alcohol and lung cancer risk in the Yunnan Tin Corporation (YTC) miners.]. *China Pub Health*, 16:707–708.
- Lu XM, Zhang YM, Lin RY *et al.* (2005). Relationship between genetic polymorphisms of metabolizing enzymes CYP2E1, GSTM1 and Kazakh's esophageal squamous cell cancer in Xinjiang, China. *World J Gastroenterol*, 11:3651–3654. [PMID:15968714](https://pubmed.ncbi.nlm.nih.gov/15968714/)
- Lubin JH, Purdue M, Kelsey K, *et al.* (2009). Total Exposure and Exposure Rate Effects for Alcohol and Smoking and Risk of Head and Neck Cancer: A Pooled Analysis of Case-Control Studies. *Am J Epidemiol*.
- Lucas D, Ménez C, Floch F *et al.* (1996). Cytochromes P4502E1 and P4501A1 genotypes and susceptibility to cirrhosis or upper aerodigestive tract cancer in alcoholic caucasians. *Alcohol Clin Exp Res*, 20:1033–1037. [doi:10.1111/j.1530-0277.1996.tb01943.x](https://doi.org/10.1111/j.1530-0277.1996.tb01943.x) [PMID:8892524](https://pubmed.ncbi.nlm.nih.gov/8892524/)
- Lucas FL, Cauley JA, Stone RA *et al.*; Study of Osteoporotic Fractures Research Group (1998). Bone mineral density and risk of breast cancer: differences by family history of breast cancer. *Am J Epidemiol*, 148:22–29. [PMID:9663400](https://pubmed.ncbi.nlm.nih.gov/9663400/)
- Lucenteforte E, Scita V, Bosetti C *et al.* (2008). Food groups and alcoholic beverages and the risk of stomach cancer: a case-control study in Italy. *Nutr Cancer*, 60:577–584. [doi:10.1080/01635580802054512](https://doi.org/10.1080/01635580802054512) [PMID:18791920](https://pubmed.ncbi.nlm.nih.gov/18791920/)
- Lund Nilssen TI, Johnsen R, Vatten LJ (2000). Socio-economic and lifestyle factors associated with the risk of prostate cancer. *Br J Cancer*, 82:1358–1363. [PMID:10755415](https://pubmed.ncbi.nlm.nih.gov/10755415/)
- Ma H, Bernstein L, Ross RK, Ursin G (2006). Hormone-related risk factors for breast cancer in women under age 50 years by estrogen and progesterone receptor status: results from a case-control and a case-case comparison. *Breast Cancer Res*, 8:R39. [doi:10.1186/bcr1514](https://doi.org/10.1186/bcr1514) [PMID:16846528](https://pubmed.ncbi.nlm.nih.gov/16846528/)
- MacArthur AC, McBride ML, Spinelli JJ *et al.* (2008). Risk of childhood leukemia associated with parental smoking and alcohol consumption prior to conception and during pregnancy: the cross-Canada childhood leukemia study. *Cancer Causes Control*, 19:283–295. [doi:10.1007/s10552-007-9091-8](https://doi.org/10.1007/s10552-007-9091-8) [PMID:18283545](https://pubmed.ncbi.nlm.nih.gov/18283545/)
- Madsen BS, Jensen HL, van den Brule AJ *et al.* (2008). Risk factors for invasive squamous cell carcinoma of the vulva and vagina—population-based case-control study in Denmark. *Int J Cancer*, 122:2827–2834. [doi:10.1002/ijc.23446](https://doi.org/10.1002/ijc.23446) [PMID:18348142](https://pubmed.ncbi.nlm.nih.gov/18348142/)
- Mahabir S, Leitzmann MF, Virtanen MJ *et al.* (2005). Prospective study of alcohol drinking and renal cell cancer risk in a cohort of Finnish male smokers. *Cancer Epidemiol Biomarkers Prev*, 14:170–175. [PMID:15668492](https://pubmed.ncbi.nlm.nih.gov/15668492/)
- Matsuo K, Hamajima N, Hirai T *et al.* (2002). Aldehyde dehydrogenase 2 (ALDH2) genotype affects rectal cancer susceptibility due to alcohol consumption. *J Epidemiol*, 12:70–76. [doi:10.2188/jea.12.70](https://doi.org/10.2188/jea.12.70) [PMID:12033531](https://pubmed.ncbi.nlm.nih.gov/12033531/)

- Matsuo K, Wakai K, Hirose K *et al.* (2006). A gene-gene interaction between ALDH2 Glu487Lys and ADH2 His47Arg polymorphisms regarding the risk of colorectal cancer in Japan. *Carcinogenesis*, 27:1018–1023. [doi:10.1093/carcin/bgi282](https://doi.org/10.1093/carcin/bgi282) PMID:16332725
- Matthias C, Bockmühl U, Jahnke V *et al.* (1998). Polymorphism in cytochrome P450 CYP2D6, CYP1A1, CYP2E1 and glutathione S-transferase, GSTM1, GSTM3, GSTT1 and susceptibility to tobacco-related cancers: studies in upper aerodigestive tract cancers. *Pharmacogenetics*, 8:91–100. [doi:10.1097/00008571-199804000-00001](https://doi.org/10.1097/00008571-199804000-00001) PMID:10022746
- Mattsson I, Wirfält E, Wallström P *et al.* (2004). High fat and alcohol intakes are risk factors of postmenopausal breast cancer: a prospective study from the Malmö diet and cancer cohort. *Int J Cancer*, 110:589–597. [doi:10.1002/ijc.20166](https://doi.org/10.1002/ijc.20166) PMID:15122593
- Michaud DS, Giovannucci E, Willett WC *et al.* (2001). Coffee and alcohol consumption and the risk of pancreatic cancer in two prospective United States cohorts. *Cancer Epidemiol Biomarkers Prev*, 10:429–437. [PMID:11352851](https://pubmed.ncbi.nlm.nih.gov/11352851/)
- Middleton Fillmore K, Chikritzhs T, Stockwell T *et al.* (2009). Alcohol use and prostate cancer: a meta-analysis. *Mol Nutr Food Res*, 53:240–255. [doi:10.1002/mnfr.200800122](https://doi.org/10.1002/mnfr.200800122) PMID:19156715
- Millikan RC, Newman B, Tse CK *et al.* (2008). Epidemiology of basal-like breast cancer. *Breast Cancer Res Treat*, 109:123–139. [doi:10.1007/s10549-007-9632-6](https://doi.org/10.1007/s10549-007-9632-6) PMID:17578664
- Mills PK, Beeson WL, Phillips RL, Fraser GE (1994). Cancer incidence among California Seventh-Day Adventists, 1976–1982. *Am J Clin Nutr*, 59: Suppl1 136S–1142S
- Minegishi Y, Tsukino H, Muto M *et al.* (2007). Susceptibility to lung cancer and genetic polymorphisms in the alcohol metabolite-related enzymes alcohol dehydrogenase 3, aldehyde dehydrogenase 2, and cytochrome P450 2E1 in the Japanese population. *Cancer*, 110:353–362. [doi:10.1002/encr.22795](https://doi.org/10.1002/encr.22795) PMID:17559142
- Miyasaka K, Kawanami T, Shimokata H *et al.* (2005). Inactive aldehyde dehydrogenase-2 increased the risk of pancreatic cancer among smokers in a Japanese male population. *Pancreas*, 30:95–98. [doi:10.1097/01.mpa.0000147084.70125.41](https://doi.org/10.1097/01.mpa.0000147084.70125.41) PMID:15714130
- Mizoue T, Inoue M, Wakai K *et al.*; Research Group for Development and Evaluation of Cancer Prevention Strategies in Japan (2008). Alcohol drinking and colorectal cancer in Japanese: a pooled analysis of results from five cohort studies. *Am J Epidemiol*, 167:1397–1406. [doi:10.1093/aje/kwn073](https://doi.org/10.1093/aje/kwn073) PMID:18420544
- Monnereau A, Orsi L, Troussard X *et al.* (2008). Cigarette smoking, alcohol drinking, and risk of lymphoid neoplasms: results of a French case-control study. *Cancer Causes Control*, 19:1147–1160. [doi:10.1007/s10552-008-9182-1](https://doi.org/10.1007/s10552-008-9182-1) PMID:18781390
- Monson RR, Lyon JL (1975). Proportional mortality among alcoholics. *Cancer*, 36:1077–1079. [doi:10.1002/1097-0142\(197509\)36:3<1077::AID-CNCR2820360335>3.0.CO;2-E](https://doi.org/10.1002/1097-0142(197509)36:3<1077::AID-CNCR2820360335>3.0.CO;2-E) PMID:1182660
- Mørch LS, Johansen D, Thygesen LC *et al.* (2007). Alcohol drinking, consumption patterns and breast cancer among Danish nurses: a cohort study. *Eur J Public Health*, 17:624–629. [doi:10.1093/eurpub/ckm036](https://doi.org/10.1093/eurpub/ckm036) PMID:17442702
- Morita M, Le Marchand L, Kono S *et al.* (2009). Genetic polymorphisms of CYP2E1 and risk of colorectal cancer: the Fukuoka Colorectal Cancer Study. *Cancer Epidemiol Biomarkers Prev*, 18:235–241. [doi:10.1158/1055-9965.EPI-08-0698](https://doi.org/10.1158/1055-9965.EPI-08-0698) PMID:19124503
- Morita S, Yano M, Shiozaki H *et al.* (1997). CYP1A1, CYP2E1 and GSTM1 polymorphisms are not associated with susceptibility to squamous-cell carcinoma of the esophagus. *Int J Cancer*, 71:192–195. [doi:10.1002/\(SICI\)1097-0215\(19970410\)71:2<192::AID-IJC11>3.0.CO;2-K](https://doi.org/10.1002/(SICI)1097-0215(19970410)71:2<192::AID-IJC11>3.0.CO;2-K) PMID:9139841

- Morita S, Yano M, Tsujinaka T *et al.* (1999). Genetic polymorphisms of drug-metabolizing enzymes and susceptibility to head-and-neck squamous-cell carcinoma. *Int J Cancer*, 80:685–688. [doi:10.1002/\(SICI\)1097-0215\(19990301\)80:5<685::AID-IJC9>3.0.CO;2-W](https://doi.org/10.1002/(SICI)1097-0215(19990301)80:5<685::AID-IJC9>3.0.CO;2-W) PMID:10048967
- Munaka M, Kohshi K, Kawamoto T *et al.* (2003). Genetic polymorphisms of tobacco- and alcohol-related metabolizing enzymes and the risk of hepatocellular carcinoma. *J Cancer Res Clin Oncol*, 129:355–360. [doi:10.1007/s00432-003-0439-5](https://doi.org/10.1007/s00432-003-0439-5) PMID:12759747
- Murata M, Tagawa M, Watanabe S *et al.* (1999). Genotype difference of aldehyde dehydrogenase 2 gene in alcohol drinkers influences the incidence of Japanese colorectal cancer patients. *Jpn J Cancer Res*, 90:711–719. [doi:10.1111/j.1349-7006.1999.tb00805.x](https://doi.org/10.1111/j.1349-7006.1999.tb00805.x) PMID:10470282
- Murata M, Takayama K, Choi BCK, Pak AWP (1996). A nested case-control study on alcohol drinking, tobacco smoking, and cancer. *Cancer Detect Prev*, 20:557–565. PMID:8939341
- Murata M, Watanabe M, Yamanaka M *et al.* (2001). Genetic polymorphisms in cytochrome P450 (CYP) 1A1, CYP1A2, CYP2E1, glutathione S-transferase (GST) M1 and GSTT1 and susceptibility to prostate cancer in the Japanese population. *Cancer Lett*, 165:171–177. [doi:10.1016/S0304-3835\(01\)00398-6](https://doi.org/10.1016/S0304-3835(01)00398-6) PMID:11275366
- Murtaugh MA, Curtin K, Sweeney C *et al.* (2007). Dietary intake of folate and co-factors in folate metabolism, MTHFR polymorphisms, and reduced rectal cancer. *Cancer Causes Control*, 18:153–163. [doi:10.1007/s10552-006-0099-2](https://doi.org/10.1007/s10552-006-0099-2) PMID:17245555
- Murtaugh MA, Ma KN, Caan BJ, Slattery ML (2004). Association of fluids from beverages with risk of rectal cancer. *Nutr Cancer*, 49:25–31. [doi:10.1207/s15327914nc4901\\_4](https://doi.org/10.1207/s15327914nc4901_4) PMID:15456632
- Muto M, Takahashi M, Ohtsu A *et al.* (2005). Risk of multiple squamous cell carcinomas both in the esophagus and the head and neck region. *Carcinogenesis*, 26:1008–1012. [doi:10.1093/carcin/bgi035](https://doi.org/10.1093/carcin/bgi035) PMID:15718256
- Muwonge R, Ramadas K, Sankila R *et al.* (2008). Role of tobacco smoking, chewing and alcohol drinking in the risk of oral cancer in Trivandrum, India: a nested case-control design using incident cancer cases. *Oral Oncol*, 44:446–454. [doi:10.1016/j.oraloncology.2007.06.002](https://doi.org/10.1016/j.oraloncology.2007.06.002) PMID:17933578
- Nakaya N, Tsubono Y, Kuriyama S *et al.* (2005). Alcohol consumption and the risk of cancer in Japanese men: the Miyagi cohort study. *Eur J Cancer Prev*, 14:169–174. [doi:10.1097/00008469-200504000-00013](https://doi.org/10.1097/00008469-200504000-00013) PMID:15785321
- Nan HM, Song YJ, Yun HY *et al.* (2005). Effects of dietary intake and genetic factors on hypermethylation of the hMLH1 gene promoter in gastric cancer. *World J Gastroenterol*, 11:3834–3841. PMID:15991278
- Nasca PC, Liu S, Baptiste MS *et al.* (1994). Alcohol consumption and breast cancer: estrogen receptor status and histology. *Am J Epidemiol*, 140:980–988. PMID:7985660
- Navarro Silvera SA, Miller AB, Rohan TE (2005). Risk factors for thyroid cancer: a prospective cohort study. *Int J Cancer*, 116:433–438. [doi:10.1002/ijc.21079](https://doi.org/10.1002/ijc.21079) PMID:15818623
- Newcomb PA, Nichols HB, Beasley JM *et al.* (2009). No difference between red wine or white wine consumption and breast cancer risk. *Cancer Epidemiol Biomarkers Prev*, 18:1007–1010. [doi:10.1158/1055-9965.EPI-08-0801](https://doi.org/10.1158/1055-9965.EPI-08-0801) PMID:19273487
- Nicholls P, Edwards G, Kyle E (1974). Alcoholics admitted to four hospitals in England. II. General and cause-specific mortality. *Q J Stud Alcohol*, 35:841–855. PMID:4411913
- Nielsen NR, Grønbaek M (2008). Interactions between intakes of alcohol and postmenopausal hormones on risk of breast cancer. *Int J Cancer*, 122:1109–1113. [doi:10.1002/ijc.23195](https://doi.org/10.1002/ijc.23195) PMID:17966122
- Nishimoto IN, Hanaoka T, Sugimura H *et al.* (2000). Cytochrome P450 2E1 polymorphism in gastric cancer in Brazil: case-control studies of Japanese Brazilians and non-Japanese Brazilians. *Cancer Epidemiol. Biomarkers Prev.*, 9:675–680.

- Nishimoto IN, Pinheiro NA, Rogatto SR *et al.* (2004). Alcohol dehydrogenase 3 genotype as a risk factor for upper aerodigestive tract cancers. *Arch Otolaryngol Head Neck Surg*, 130:78–82. [doi:10.1001/archotol.130.1.78](https://doi.org/10.1001/archotol.130.1.78) PMID:14732773
- Nishino Y, Wakai K, Kondo T *et al.*; JACC Study Group (2006). Alcohol consumption and lung cancer mortality in Japanese men: results from Japan collaborative cohort (JACC) study. *J Epidemiol*, 16:49–56. [doi:10.2188/jea.16.49](https://doi.org/10.2188/jea.16.49) PMID:16537984
- Nomura A, Grove JS, Stemmermann GN, Severson RK (1990). A prospective study of stomach cancer and its relation to diet, cigarettes, and alcohol consumption. *Cancer Res*, 50:627–631. PMID:2297702
- Nomura AMY, Stemmermann GN, Chyou PH (1995). Gastric cancer among the Japanese in Hawaii. *Jpn J Cancer Res*, 86:916–923. [doi:10.1111/j.1349-7006.1995.tb03001.x](https://doi.org/10.1111/j.1349-7006.1995.tb03001.x) PMID:7493909
- Nomura T, Noma H, Shibahara T *et al.* (2000). Aldehyde dehydrogenase 2 and glutathione S-transferase M1 polymorphisms in relation to the risk for oral cancer in Japanese drinkers. *Oral Oncol*, 36:42–46. [doi:10.1016/S1368-8375\(99\)00048-2](https://doi.org/10.1016/S1368-8375(99)00048-2) PMID:10889918
- Ohishi W, Fujiwara S, Cologne JB *et al.* (2008). Risk factors for hepatocellular carcinoma in a Japanese population: a nested case-control study. *Cancer Epidemiol Biomarkers Prev*, 17:846–854. [doi:10.1158/1055-9965.EPI-07-2806](https://doi.org/10.1158/1055-9965.EPI-07-2806) PMID:18398026
- Olivieri EHR, da Silva SD, Mendonça FF *et al.* (2009). CYP1A2\*1C, CYP2E1\*5B, and GSTM1 polymorphisms are predictors of risk and poor outcome in head and neck squamous cell carcinoma patients. *Oral Oncol*, 45:e73–e79. [doi:10.1016/j.oraloncology.2009.03.004](https://doi.org/10.1016/j.oraloncology.2009.03.004) PMID:19442564
- Olshan AF, Weissler MC, Watson MA, Bell DA (2001). Risk of head and neck cancer and the alcohol dehydrogenase 3 genotype. *Carcinogenesis*, 22:57–61. [doi:10.1093/carcin/22.1.57](https://doi.org/10.1093/carcin/22.1.57) PMID:11159741
- Omenn GS, Goodman GE, Thornquist MD *et al.* (1996). Risk factors for lung cancer and for intervention effects in CARET, the Beta-Carotene and Retinol Efficacy Trial. *J Natl Cancer Inst*, 88:1550–1559. [doi:10.1093/jnci/88.21.1550](https://doi.org/10.1093/jnci/88.21.1550) PMID:8901853
- Otani T, Iwasaki M, Hanaoka T *et al.* (2005). Folate, vitamin B6, vitamin B12, and vitamin B2 intake, genetic polymorphisms of related enzymes, and risk of colorectal cancer in a hospital-based case-control study in Japan. *Nutr Cancer*, 53:42–50. [doi:10.1207/s15327914nc5301\\_5](https://doi.org/10.1207/s15327914nc5301_5) PMID:16351505
- Otani T, Iwasaki M, Yamamoto S *et al.*; Japan Public Health Center-based Prospective Study Group (2003). Alcohol consumption, smoking, and subsequent risk of colorectal cancer in middle-aged and elderly Japanese men and women: Japan Public Health Center-based prospective study. *Cancer Epidemiol Biomarkers Prev*, 12:1492–1500. PMID:14693743
- Oyama T, Kawamoto T, Mizoue T *et al.* (1997). Cytochrome P450 2E1 polymorphism as a risk factor for lung cancer: in relation to p53 gene mutation. *Anticancer Res*, 17 1B:583–587. PMID:9066584
- Ozasa K; Japan Collaborative Cohort Study for Evaluation of Cancer (2007). Alcohol use and mortality in the Japan Collaborative Cohort Study for Evaluation of Cancer (JACC). *Asian Pac J Cancer Prev*, 8 Suppl:81–88. PMID:18260706
- Pandeya N, Williams G, Green AC *et al.*; Australian Cancer Study (2009). Alcohol consumption and the risks of adenocarcinoma and squamous cell carcinoma of the esophagus. *Gastroenterology*, 136:1215–1224, e1–e2. [doi:10.1053/j.gastro.2008.12.052](https://doi.org/10.1053/j.gastro.2008.12.052) PMID:19250648
- Parkin DM, Srivatanakul P, Khlat M *et al.* (1991). Liver cancer in Thailand. I. A case-control study of cholangiocarcinoma. *Int J Cancer*, 48:323–328. [doi:10.1002/ijc.2910480302](https://doi.org/10.1002/ijc.2910480302) PMID:1645697
- Pelucchi C, Tavani A, La Vecchia C (2008). Coffee and alcohol consumption and bladder cancer. *Scand J Urol Nephrol Suppl*, 42 218:37–44. [doi:10.1080/03008880802237090](https://doi.org/10.1080/03008880802237090) PMID:18815915
- Pereira Serafim PV, Cotrim Guerreiro da Silva ID, Manoukias Forones N (2008). Relationship between genetic polymorphism of CYP1A1 at codon 462 (Ile462Val) in colorectal cancer. *Int J Biol Markers*, 23:18–23. PMID:18409146

- Persson I, Johansson I, Bergling H *et al.* (1993). Genetic polymorphism of cytochrome P4502E1 in a Swedish population. Relationship to incidence of lung cancer. *FEBS Lett*, 319:207–211. [doi:10.1016/0014-5793\(93\)80547-8](https://doi.org/10.1016/0014-5793(93)80547-8) PMID:8096192
- Persson I, Johansson I, Lou YC *et al.* (1999). Genetic polymorphism of xenobiotic metabolizing enzymes among Chinese lung cancer patients. *Int J Cancer*, 81:325–329. [doi:10.1002/\(SICI\)1097-0215\(19990505\)81:3<325::AID-IJC2>3.0.CO;2-S](https://doi.org/10.1002/(SICI)1097-0215(19990505)81:3<325::AID-IJC2>3.0.CO;2-S) PMID:10209943
- Peters ES, McClean MD, Liu M *et al.* (2005). The ADH1C polymorphism modifies the risk of squamous cell carcinoma of the head and neck associated with alcohol and tobacco use. *Cancer Epidemiol Biomarkers Prev*, 14:476–482. [doi:10.1158/1055-9965.EPI-04-0431](https://doi.org/10.1158/1055-9965.EPI-04-0431) PMID:15734975
- Petri AL, Tjønneland A, Gamborg M *et al.* (2004). Alcohol intake, type of beverage, and risk of breast cancer in pre- and postmenopausal women. *Alcohol Clin Exp Res*, 28:1084–1090. [doi:10.1097/01.ALC.0000130812.85638.E1](https://doi.org/10.1097/01.ALC.0000130812.85638.E1) PMID:15252295
- Platz EA, Leitzmann MF, Rimm EB *et al.* (2004). Alcohol intake, drinking patterns, and risk of prostate cancer in a large prospective cohort study. *Am J Epidemiol*, 159:444–453. [doi:10.1093/aje/kwh062](https://doi.org/10.1093/aje/kwh062) PMID:14977640
- Pollack ES, Nomura AM, Heilbrun LK *et al.* (1984). Prospective study of alcohol consumption and cancer. *N Engl J Med*, 310:617–621. [doi:10.1056/NEJM198403083101003](https://doi.org/10.1056/NEJM198403083101003) PMID:6694673
- Potter JD, Sellers TA, Folsom AR, McGovern PG (1992). Alcohol, beer, and lung cancer in postmenopausal women. The Iowa Women's Health Study. *Ann Epidemiol*, 2:587–595. [doi:10.1016/1047-2797\(92\)90003-9](https://doi.org/10.1016/1047-2797(92)90003-9) PMID:1342310
- Prescott E, Grønbaek M, Becker U, Sørensen TI (1999). Alcohol intake and the risk of lung cancer: influence of type of alcoholic beverage. *Am J Epidemiol*, 149:463–470. PMID:10067906
- Prior P (1988). Long-term cancer risk in alcoholism. *Alcohol Alcohol*, 23:163–171. PMID:3390240
- Putnam SD, Cerhan JR, Parker AS *et al.* (2000). Lifestyle and anthropometric risk factors for prostate cancer in a cohort of Iowa men. *Ann Epidemiol*, 10:361–369. [doi:10.1016/S1047-2797\(00\)00057-0](https://doi.org/10.1016/S1047-2797(00)00057-0) PMID:10964002
- Qin JM, Yang L, Chen B *et al.* (2008). Interaction of methylenetetrahydrofolate reductase C677T, cytochrome P4502E1 polymorphism and environment factors in esophageal cancer in Kazakh population. *World J Gastroenterol*, 14:6986–6992. [doi:10.3748/wjg.14.6986](https://doi.org/10.3748/wjg.14.6986) PMID:19058336
- Quiñones L, Lucas D, Godoy J *et al.* (2001). CYP1A1, CYP2E1 and GSTM1 genetic polymorphisms. The effect of single and combined genotypes on lung cancer susceptibility in Chilean people. *Cancer Lett*, 174:35–44. [doi:10.1016/S0304-3835\(01\)00686-3](https://doi.org/10.1016/S0304-3835(01)00686-3) PMID:11675150
- Rashidkhani B, Åkesson A, Lindblad P, Wolk A (2005). Alcohol consumption and risk of renal cell carcinoma: a prospective study of Swedish women. *Int J Cancer*, 117:848–853. [doi:10.1002/ijc.21231](https://doi.org/10.1002/ijc.21231) PMID:15957170
- Rehm J, Patra J, Popova S (2007). Alcohol drinking cessation and its effect on esophageal and head and neck cancers: a pooled analysis. *Int J Cancer*, 121:1132–1137. [doi:10.1002/ijc.22798](https://doi.org/10.1002/ijc.22798) PMID:17487833
- Ribas G, Milne RL, Gonzalez-Neira A, Benítez J (2008). Haplotype patterns in cancer-related genes with long-range linkage disequilibrium: no evidence of association with breast cancer or positive selection. *Eur J Hum Genet*, 16:252–260. [doi:10.1038/sj.ejhg.5201953](https://doi.org/10.1038/sj.ejhg.5201953) PMID:18000525
- Risch A, Ramroth H, Raedts V *et al.* (2003). Laryngeal cancer risk in Caucasians is associated with alcohol and tobacco consumption but not modified by genetic polymorphisms in class I alcohol dehydrogenases ADH1B and ADH1C, and glutathione-S-transferases GSTM1 and GSTT1. *Pharmacogenetics*, 13:225–230. [doi:10.1097/00008571-200304000-00007](https://doi.org/10.1097/00008571-200304000-00007) PMID:12668919
- Robinette CD, Hrubec Z, Fraumeni JF Jr (1979). Chronic alcoholism and subsequent mortality in World War II veterans. *Am J Epidemiol*, 109:687–700. PMID:453188

- Rod NH, Hansen AM, Nielsen J *et al.* (2009). Low-risk factor profile, estrogen levels, and breast cancer risk among postmenopausal women. *Int J Cancer*, 124:1935–1940. [doi:10.1002/ijc.24136](https://doi.org/10.1002/ijc.24136) [PMID:19123466](https://pubmed.ncbi.nlm.nih.gov/19123466/)
- Rohan TE, Jain M, Howe GR, Miller AB (2000). Alcohol consumption and risk of breast cancer: a cohort study. *Cancer Causes Control*, 11:239–247. [doi:10.1023/A:1008933824645](https://doi.org/10.1023/A:1008933824645) [PMID:10782658](https://pubmed.ncbi.nlm.nih.gov/10782658/)
- Rohrmann S, Linseisen J, Boshuizen HC *et al.* (2006). Ethanol intake and risk of lung cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Am J Epidemiol*, 164:1103–1114. [doi:10.1093/aje/kwj326](https://doi.org/10.1093/aje/kwj326) [PMID:16987924](https://pubmed.ncbi.nlm.nih.gov/16987924/)
- Rohrmann S, Linseisen J, Key TJ *et al.* (2008). Alcohol consumption and the risk for prostate cancer in the European Prospective Investigation into Cancer and Nutrition. *Cancer Epidemiol Biomarkers Prev*, 17:1282–1287. [doi:10.1158/1055-9965.EPI-07-2888](https://doi.org/10.1158/1055-9965.EPI-07-2888) [PMID:18483352](https://pubmed.ncbi.nlm.nih.gov/18483352/)
- Rohrmann S, Linseisen J, Vrieling A *et al.* (2009). Ethanol intake and the risk of pancreatic cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Cancer Causes Control*, 20:785–794. [doi:10.1007/s10552-008-9293-8](https://doi.org/10.1007/s10552-008-9293-8) [PMID:19145468](https://pubmed.ncbi.nlm.nih.gov/19145468/)
- Rossini A, Rapozo DCM, Soares Lima SC *et al.* (2007). Polymorphisms of GSTP1 and GSTT1, but not of CYP2A6, CYP2E1 or GSTM1, modify the risk for esophageal cancer in a western population. *Carcinogenesis*, 28:2537–2542. [doi:10.1093/carcin/bgm222](https://doi.org/10.1093/carcin/bgm222) [PMID:17916905](https://pubmed.ncbi.nlm.nih.gov/17916905/)
- Rudant J, Menegaux F, Leverger G *et al.* (2008). Childhood hematopoietic malignancies and parental use of tobacco and alcohol: the ESCALE study (SFCE). *Cancer Causes Control*, 19:1277–1290. [doi:10.1007/s10552-008-9199-5](https://doi.org/10.1007/s10552-008-9199-5) [PMID:18618277](https://pubmed.ncbi.nlm.nih.gov/18618277/)
- Ruwali M, Khan AJ, Shah PP *et al.* (2009). Cytochrome P450 2E1 and head and neck cancer: interaction with genetic and environmental risk factors. *Environ Mol Mutagen*, 50:473–482. [doi:10.1002/em.20488](https://doi.org/10.1002/em.20488) [PMID:19334053](https://pubmed.ncbi.nlm.nih.gov/19334053/)
- Sakamoto T, Hara M, Higaki Y *et al.* (2006). Influence of alcohol consumption and gene polymorphisms of ADH2 and ALDH2 on hepatocellular carcinoma in a Japanese population. *Int J Cancer*, 118:1501–1507. [doi:10.1002/ijc.21505](https://doi.org/10.1002/ijc.21505) [PMID:16187278](https://pubmed.ncbi.nlm.nih.gov/16187278/)
- Sakata K, Hoshiyama Y, Morioka S *et al.* JACC Study Group. (2005). Smoking, alcohol drinking and esophageal cancer: findings from the JACC Study. *J Epidemiol*, 15: Suppl 2S212–S219. [doi:10.2188/jea.15.S212](https://doi.org/10.2188/jea.15.S212) [PMID:16127236](https://pubmed.ncbi.nlm.nih.gov/16127236/)
- Sanjoaquin MA, Appleby PN, Thorogood M *et al.* (2004). Nutrition, lifestyle and colorectal cancer incidence: a prospective investigation of 10998 vegetarians and non-vegetarians in the United Kingdom. *Br J Cancer*, 90:118–121. [doi:10.1038/sj.bjc.6601441](https://doi.org/10.1038/sj.bjc.6601441) [PMID:14710217](https://pubmed.ncbi.nlm.nih.gov/14710217/)
- Sasazuki S, Sasaki S, Tsugane S; Japan Public Health Center Study Group (2002). Cigarette smoking, alcohol consumption and subsequent gastric cancer risk by subsite and histologic type. *Int J Cancer*, 101:560–566. [doi:10.1002/ijc.10649](https://doi.org/10.1002/ijc.10649) [PMID:12237898](https://pubmed.ncbi.nlm.nih.gov/12237898/)
- Schatzkin A, Jones DY, Hoover RN *et al.* (1987). Alcohol consumption and breast cancer in the epidemiologic follow-up study of the first National Health and Nutrition Examination Survey. *N Engl J Med*, 316:1169–1173. [doi:10.1056/NEJM198705073161901](https://doi.org/10.1056/NEJM198705073161901) [PMID:3574367](https://pubmed.ncbi.nlm.nih.gov/3574367/)
- Schmidt W, Popham RE (1981). The role of drinking and smoking in mortality from cancer and other causes in male alcoholics. *Cancer*, 47:1031–1041. [doi:10.1002/1097-0142\(19810301\)47:5<1031::AID-CNCR2820470534>3.0.CO;2-C](https://doi.org/10.1002/1097-0142(19810301)47:5<1031::AID-CNCR2820470534>3.0.CO;2-C) [PMID:7226036](https://pubmed.ncbi.nlm.nih.gov/7226036/)
- Schouten LJ, Zeegers MPA, Goldbohm RA, van den Brandt PA (2004). Alcohol and ovarian cancer risk: results from the Netherlands Cohort Study. *Cancer Causes Control*, 15:201–209. [doi:10.1023/B:CACO.0000019512.71560.2b](https://doi.org/10.1023/B:CACO.0000019512.71560.2b) [PMID:15017133](https://pubmed.ncbi.nlm.nih.gov/15017133/)

- Schuurman AG, Goldbohm RA, van den Brandt PA (1999). A prospective cohort study on consumption of alcoholic beverages in relation to prostate cancer incidence (The Netherlands). *Cancer Causes Control*, 10:597–605. [doi:10.1023/A:1008925103542](https://doi.org/10.1023/A:1008925103542) PMID:10616828
- Schwartz SM, Doody DR, Fitzgibbons ED *et al.* (2001). Oral squamous cell cancer risk in relation to alcohol consumption and alcohol dehydrogenase-3 genotypes. *Cancer Epidemiol Biomarkers Prev*, 10:1137–1144. [PMID:11700261](https://pubmed.ncbi.nlm.nih.gov/11700261/)
- Sesso HD, Paffenbarger RS Jr, Lee IM (2001). Alcohol consumption and risk of prostate cancer: The Harvard Alumni Health Study. *Int J Epidemiol*, 30:749–755. [doi:10.1093/ije/30.4.749](https://doi.org/10.1093/ije/30.4.749) PMID:11511598
- Setiawan VW, Monroe KR, Goodman MT *et al.* (2008). Alcohol consumption and endometrial cancer risk: the multiethnic cohort. *Int J Cancer*, 122:634–638. [doi:10.1002/ijc.23072](https://doi.org/10.1002/ijc.23072) PMID:17764072
- Setiawan VW, Monroe KR, Wilkens LR *et al.* (2009). Breast cancer risk factors defined by estrogen and progesterone receptor status: the multiethnic cohort study. *Am J Epidemiol*, 169:1251–1259. [doi:10.1093/aje/kwp036](https://doi.org/10.1093/aje/kwp036) PMID:19318616
- Shaib YH, El-Serag HB, Nooka AK *et al.* (2007). Risk factors for intrahepatic and extrahepatic cholangiocarcinoma: a hospital-based case-control study. *Am J Gastroenterol*, 102:1016–1021. [doi:10.1111/j.1572-0241.2007.01104.x](https://doi.org/10.1111/j.1572-0241.2007.01104.x) PMID:17324130
- Shibata A, Fukuda K, Nishiyori A *et al.* (1998). A case-control study on male hepatocellular carcinoma based on hospital and community controls. *J Epidemiol*, 8:1–5. [doi:10.2188/jea.8.1](https://doi.org/10.2188/jea.8.1) PMID:9575688
- Shibata A, Mack TM, Paganini-Hill A *et al.* (1994). A prospective study of pancreatic cancer in the elderly. *Int J Cancer*, 58:46–49. [doi:10.1002/ijc.2910580109](https://doi.org/10.1002/ijc.2910580109) PMID:8014014
- Shimazu T, Inoue M, Sasazuki S *et al.*; Japan Public Health Center-based Prospective Study Group (2008). Alcohol and risk of lung cancer among Japanese men: data from a large-scale population-based cohort study, the JPHC study. *Cancer Causes Control*, 19:1095–1102. [doi:10.1007/s10552-008-9173-2](https://doi.org/10.1007/s10552-008-9173-2) PMID:18493860
- Shimizu N, Nagata C, Shimizu H *et al.* (2003). Height, weight, and alcohol consumption in relation to the risk of colorectal cancer in Japan: a prospective study. *Br J Cancer*, 88:1038–1043. [doi:10.1038/sj.bjc.6600845](https://doi.org/10.1038/sj.bjc.6600845) PMID:12671701
- Shin HR, Lee CU, Park HJ *et al.* (1996). Hepatitis B and C virus, *Clonorchis sinensis* for the risk of liver cancer: a case-control study in Pusan, Korea. *Int J Epidemiol*, 25:933–940. [doi:10.1093/ije/25.5.933](https://doi.org/10.1093/ije/25.5.933) PMID:8921477
- Sigvardsson S, Hardell L, Przybeck TR, Cloninger R (1996). Increased cancer risk among Swedish female alcoholics. *Epidemiology*, 7:140–143. [doi:10.1097/00001648-199603000-00006](https://doi.org/10.1097/00001648-199603000-00006) PMID:8834552
- Simon MS, Carman W, Wolfe R, Schottenfeld D (1991). Alcohol consumption and the risk of breast cancer: a report from the Tecumseh Community Health Study. *J Clin Epidemiol*, 44:755–761. [doi:10.1016/0895-4356\(91\)90127-U](https://doi.org/10.1016/0895-4356(91)90127-U) PMID:1941026
- Singh PN, Fraser GE (1998). Dietary risk factors for colon cancer in a low-risk population. *Am J Epidemiol*, 148:761–774. [PMID:9786231](https://pubmed.ncbi.nlm.nih.gov/9786231/)
- Sjödahl K, Lu Y, Nilsen TIL *et al.* (2007). Smoking and alcohol drinking in relation to risk of gastric cancer: a population-based, prospective cohort study. *Int J Cancer*, 120:128–132. [doi:10.1002/ijc.22157](https://doi.org/10.1002/ijc.22157) PMID:17036324
- Solomon PR, Selvam GS, Shanmugam G (2008). Polymorphism in ADH and MTHFR genes in oral squamous cell carcinoma of Indians. *Oral Dis*, 14:633–639. [doi:10.1111/j.1601-0825.2007.01437.x](https://doi.org/10.1111/j.1601-0825.2007.01437.x) PMID:18266839
- Sørensen HT, Friis S, Olsen JH *et al.* (1998). Risk of liver and other types of cancer in patients with cirrhosis: a nationwide cohort study in Denmark. *Hepatology*, 28:921–925. [doi:10.1002/hep.510280404](https://doi.org/10.1002/hep.510280404) PMID:9755226

- Soya SS, Vinod T, Reddy KS *et al.* (2008). CYP2E1 polymorphisms and gene-environment interactions in the risk of upper aerodigestive tract cancers among Indians. *Pharmacogenomics*, 9:551–560. [doi:10.2217/14622416.9.5.551](https://doi.org/10.2217/14622416.9.5.551) PMID:18466102
- Sprague BL, Trentham-Dietz A, Egan KM *et al.* (2008). Proportion of invasive breast cancer attributable to risk factors modifiable after menopause. *Am J Epidemiol*, 168:404–411. [doi:10.1093/aje/kwn143](https://doi.org/10.1093/aje/kwn143) PMID:18552361
- Sriamporn S, Wiangnon S, Suwanrungruang K *et al.* (2007). Risk factors for colorectal cancer in northeast Thailand: lifestyle related. *Asian Pac J Cancer Prev*, 8:573–577. PMID:18260731
- Stemmermann GN, Nomura AMY, Chyou P-H, Yoshizawa C (1990). Prospective study of alcohol intake and large bowel cancer. *Dig Dis Sci*, 35:1414–1420. [doi:10.1007/BF01536750](https://doi.org/10.1007/BF01536750) PMID:2226103
- Stevens RJ, Roddam AW, Spencer EA *et al.*; Million Women Study Collaborators (2009). Factors associated with incident and fatal pancreatic cancer in a cohort of middle-aged women. *Int J Cancer*, 124:2400–2405. [doi:10.1002/ijc.24196](https://doi.org/10.1002/ijc.24196) PMID:19165860
- Stolzenberg-Solomon RZ, Chang SC, Leitzmann MF *et al.* (2006). Folate intake, alcohol use, and postmenopausal breast cancer risk in the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial. *Am J Clin Nutr*, 83:895–904. PMID:16600944
- Stolzenberg-Solomon RZ, Pietinen P, Barrett MJ *et al.* (2001). Dietary and other methyl-group availability factors and pancreatic cancer risk in a cohort of male smokers. *Am J Epidemiol*, 153:680–687. [doi:10.1093/aje/153.7.680](https://doi.org/10.1093/aje/153.7.680) PMID:11282796
- Sturgis EM, Dahlstrom KR, Guan Y *et al.* (2001). Alcohol dehydrogenase 3 genotype is not associated with risk of squamous cell carcinoma of the oral cavity and pharynx. *Cancer Epidemiol Biomarkers Prev*, 10:273–275. PMID:11303599
- Su LJ, Arab L (2004). Alcohol consumption and risk of colon cancer: evidence from the national health and nutrition examination survey I epidemiologic follow-up study. *Nutr Cancer*, 50:111–119. [doi:10.1207/s15327914nc5002\\_1](https://doi.org/10.1207/s15327914nc5002_1) PMID:15623458
- Suadicani P, Hein HO, Gyntelberg F (1993). Height, weight, and risk of colorectal cancer. An 18-year follow-up in a cohort of 5249 men. *Scand J Gastroenterol*, 28:285–288. [doi:10.3109/00365529309096087](https://doi.org/10.3109/00365529309096087) PMID:8446855
- Sugimura T, Kumimoto H, Tohna I *et al.* (2006). Gene-environment interaction involved in oral carcinogenesis: molecular epidemiological study for metabolic and DNA repair gene polymorphisms. *J Oral Pathol Med*, 35:11–18. [doi:10.1111/j.1600-0714.2005.00364.x](https://doi.org/10.1111/j.1600-0714.2005.00364.x) PMID:16393248
- Sundby P (1967) *Alcoholism and Mortality*, Oslo, Universitetsforlaget.
- Sung NY, Choi KS, Park EC *et al.* (2007). Smoking, alcohol and gastric cancer risk in Korean men: the National Health Insurance Corporation Study. *Br J Cancer*, 97:700–704. [doi:10.1038/sj.bjc.6603893](https://doi.org/10.1038/sj.bjc.6603893) PMID:17637680
- Sutcliffe S, Giovannucci E, Leitzmann MF *et al.* (2007). A prospective cohort study of red wine consumption and risk of prostate cancer. *Int J Cancer*, 120:1529–1535. [doi:10.1002/ijc.22498](https://doi.org/10.1002/ijc.22498) PMID:17211860
- Suzuki R, Orsini N, Mignone L *et al.* (2008). Alcohol intake and risk of breast cancer defined by estrogen and progesterone receptor status — a meta-analysis of epidemiological studies. *Int J Cancer* 122: 1832–1841. PMID:18067133
- Suzuki R, Ye W, Rylander-Rudqvist T *et al.* (2005). Alcohol and postmenopausal breast cancer risk defined by estrogen and progesterone receptor status: a prospective cohort study. *J Natl Cancer Inst*, 97:1601–1608. [doi:10.1093/jnci/dji341](https://doi.org/10.1093/jnci/dji341) PMID:16264180

- Takeshita T, Yang X, Inoue Y *et al.* (2000b). Relationship between alcohol drinking, ADH2 and ALDH2 genotypes, and risk for hepatocellular carcinoma in Japanese. *Cancer Lett*, 149:69–76. [doi:10.1016/S0304-3835\(99\)00343-2](https://doi.org/10.1016/S0304-3835(99)00343-2) PMID:10737710
- Takezaki T, Inoue M, Kataoka H *et al.* (2003). Diet and lung cancer risk from a 14-year population-based prospective study in Japan: with special reference to fish consumption. *Nutr Cancer*, 45:160–167. [doi:10.1207/S15327914NC4502\\_04](https://doi.org/10.1207/S15327914NC4502_04) PMID:12881009
- Talamini R, Bosetti C, La Vecchia C *et al.* (2002). Combined effect of tobacco and alcohol on laryngeal cancer risk: a case-control study. *Cancer Causes Control*, 13:957–964. [doi:10.1023/A:1021944123914](https://doi.org/10.1023/A:1021944123914) PMID:12588092
- Tan W, Song N, Wang GQ *et al.* (2000). Impact of genetic polymorphisms in cytochrome P450 2E1 and glutathione S-transferases M1, T1, and P1 on susceptibility to esophageal cancer among high-risk individuals in China. *Cancer Epidemiol Biomarkers Prev*, 9:551–556. PMID:10868687
- Tanabe H, Ohhira M, Ohtsubo T *et al.* (1999). Genetic polymorphism of aldehyde dehydrogenase 2 in patients with upper aerodigestive tract cancer. *Alcohol Clin Exp Res*, 23 Suppl;17S–20S. [doi:10.1111/j.1530-0277.1999.tb04527.x](https://doi.org/10.1111/j.1530-0277.1999.tb04527.x) PMID:10235272
- Terry MB, Gammon MD, Zhang FF *et al.* (2006). ADH3 genotype, alcohol intake and breast cancer risk. *Carcinogenesis*, 27:840–847. [doi:10.1093/carcin/bgi285](https://doi.org/10.1093/carcin/bgi285) PMID:16344274
- Terry MB, Gammon MD, Zhang FF *et al.* (2007a). Alcohol dehydrogenase 3 and risk of esophageal and gastric adenocarcinomas. *Cancer Causes Control*, 18:1039–1046. [doi:10.1007/s10552-007-9046-0](https://doi.org/10.1007/s10552-007-9046-0) PMID:17665311
- Terry MB, Knight JA, Zablotska L *et al.* (2007). Alcohol metabolism, alcohol intake, and breast cancer risk: a sister-set analysis using the Breast Cancer Family Registry. *Breast Cancer Res Treat*, 106:281–288. [doi:10.1007/s10549-007-9498-7](https://doi.org/10.1007/s10549-007-9498-7) PMID:17268812
- Terry MB, Knight JA, Zablotska L *et al.* (2007b). Alcohol metabolism, alcohol intake, and breast cancer risk: a sister-set analysis using the Breast Cancer Family Registry. *Breast Cancer Res Treat*, 106:281–288. [doi:10.1007/s10549-007-9498-7](https://doi.org/10.1007/s10549-007-9498-7) PMID:17268812
- Terry P, Baron JA, Weiderpass E *et al.* (1999). Lifestyle and endometrial cancer risk: a cohort study from the Swedish Twin Registry. *Int J Cancer*, 82:38–42. [doi:10.1002/\(SICI\)1097-0215\(19990702\)82:1<38::AID-IJC8>3.0.CO;2-Q](https://doi.org/10.1002/(SICI)1097-0215(19990702)82:1<38::AID-IJC8>3.0.CO;2-Q) PMID:10360818
- Thun MJ, Hannan LM, DeLancey JO (2009). Alcohol consumption not associated with lung cancer mortality in lifelong nonsmokers. *Cancer Epidemiol Biomarkers Prev*, 18:2269–2272. [doi:10.1158/1055-9965.EPI-09-0361](https://doi.org/10.1158/1055-9965.EPI-09-0361) PMID:19661085
- Thun MJ, Peto R, Lopez AD *et al.* (1997). Alcohol consumption and mortality among middle-aged and elderly U.S. adults. *N Engl J Med*, 337:1705–1714. [doi:10.1056/NEJM199712113372401](https://doi.org/10.1056/NEJM199712113372401) PMID:9392695
- Thygesen LC, Albertsen K, Johansen C, Grønbaek M (2005). Cancer incidence among Danish brewery workers. *Int J Cancer*, 116: 774–778
- Thygesen LC, Keiding N, Johansen C, Grønbaek M (2007). Changes in alcohol intake and risk of upper digestive tract cancer. *Acta Oncol*, 46:1085–1089. [doi:10.1080/02841860701441806](https://doi.org/10.1080/02841860701441806) PMID:17851863
- Thygesen LC, Mikkelsen P, Andersen TV *et al.* (2009). Cancer incidence among patients with alcohol use disorders—long-term follow-up. *Alcohol Alcohol*, 44:387–391. PMID:19491282
- Thygesen LC, Wu K, Grønbaek M *et al.* (2008). Alcohol intake and colorectal cancer: a comparison of approaches for including repeated measures of alcohol consumption. *Epidemiology*, 19:258–264. [doi:10.1097/EDE.0b013e31816339e0](https://doi.org/10.1097/EDE.0b013e31816339e0) PMID:18300715

- Tiemersma EW, Wark PA, Ocké MC *et al.* (2003). Alcohol consumption, alcohol dehydrogenase 3 polymorphism, and colorectal adenomas. *Cancer Epidemiol Biomarkers Prev*, 12:419–425. [PMID:12750236](#)
- Tjønneland A, Christensen J, Olsen A *et al.* (2007). Alcohol intake and breast cancer risk: the European Prospective Investigation into Cancer and Nutrition (EPIC). *Cancer Causes Control*, 18:361–373. [doi:10.1007/s10552-006-0112-9](#) [PMID:17364225](#)
- Tjønneland A, Christensen J, Thomsen BL *et al.* (2004). Lifetime alcohol consumption and postmenopausal breast cancer rate in Denmark: a prospective cohort study. *J Nutr*, 134:173–178. [PMID:14704313](#)
- Tjønneland A, Thomsen BL, Stripp C *et al.* (2003). Alcohol intake, drinking patterns and risk of postmenopausal breast cancer in Denmark: a prospective cohort study. *Cancer Causes Control*, 14:277–284. [doi:10.1023/A:1023640720385](#) [PMID:12814207](#)
- Tønnesen H, Møller H, Andersen JR *et al.* (1994). Cancer morbidity in alcohol abusers. *Br J Cancer*, 69:327–332. [doi:10.1038/bjc.1994.59](#) [PMID:8297729](#)
- Toriola AT, Kurl S, Laukanen JA *et al.* (2008). Alcohol consumption and risk of colorectal cancer: the Findrink study. *Eur J Epidemiol*, 23:395–401. [doi:10.1007/s10654-008-9244-4](#) [PMID:18409007](#)
- Toriola AT, Kurl S, Laukanen JA, Kauhanen J (2009). Does binge drinking increase the risk of lung cancer: results from the Findrink study. *Eur J Public Health*, 19:389–393. [doi:10.1093/eurpub/ckp049](#) [PMID:19369490](#)
- Tramacere I, Scotti L, Jenab M *et al.* (2009). Alcohol drinking and pancreatic cancer risk: a meta-analysis of the dose-risk relation. *Int J Cancer*. 15:1474–11486. [PMID:19816941](#)
- Tran GD, Sun X-D, Abnet CC *et al.* (2005). Prospective study of risk factors for esophageal and gastric cancers in the Linxian general population trial cohort in China. *Int J Cancer*, 113:456–463. [doi:10.1002/ijc.20616](#) [PMID:15455378](#)
- Trentham-Dietz A, Newcomb PA, Nichols HB, Hampton JM (2007). Breast cancer risk factors and second primary malignancies among women with breast cancer. *Breast Cancer Res Treat*, 105:195–207. [doi:10.1007/s10549-006-9446-y](#) [PMID:17186360](#)
- Trivers KF, Lund MJ, Porter PL *et al.* (2009). The epidemiology of triple-negative breast cancer, including race. *Cancer Causes Control*, 20:1071–1082. [doi:10.1007/s10552-009-9331-1](#) [PMID:19343511](#)
- Tsong WH, Koh WP, Yuan JM *et al.* (2007). Cigarettes and alcohol in relation to colorectal cancer: the Singapore Chinese Health Study. *Br J Cancer*, 96:821–827. [doi:10.1038/sj.bjc.6603623](#) [PMID:17311023](#)
- Tsutsumi M, Takase S, Takada A (1993). Genetic factors related to the development of carcinoma in digestive organs in alcoholics. *Alcohol Alcohol Suppl*, 1B:21–26. [PMID:8003125](#)
- Vachon CM, Cerhan JR, Vierkant RA, Sellers TA (2001). Investigation of an interaction of alcohol intake and family history on breast cancer risk in the Minnesota Breast Cancer Family Study. *Cancer*, 92:240–248. [doi:10.1002/1097-0142\(20010715\)92:2<240::AID-CNCR1315>3.0.CO;2-I](#) [PMID:11466675](#)
- van der Logt EMJ, Bergevoet SM, Roelofs HMJ *et al.* (2006). Role of epoxide hydrolase, NAD(P)H:quinone oxidoreductase, cytochrome P450 2E1 or alcohol dehydrogenase genotypes in susceptibility to colorectal cancer. *Mutat Res*, 593:39–49. [doi:10.1016/j.mrfimm.2005.06.018](#) [PMID:16039674](#)
- van Dijk BAC, van Houwelingen KP, Witjes JA *et al.* (2001). Alcohol dehydrogenase type 3 (ADH3) and the risk of bladder cancer. *Eur Urol*, 40:509–514. [doi:10.1159/000049827](#) [PMID:11752857](#)
- Vaughan TL, Shapiro JA, Burt RD *et al.* (1996). Nasopharyngeal cancer in a low-risk population: defining risk factors by histological type. *Cancer Epidemiol Biomarkers Prev*, 5:587–593. [PMID:8824359](#)

- Velicer CM, Kristal A, White E (2006). Alcohol use and the risk of prostate cancer: results from the VITAL cohort study. *Nutr Cancer*, 56:50–56. [doi:10.1207/s15327914nc5601\\_7](https://doi.org/10.1207/s15327914nc5601_7) PMID:17176217
- Vioque J, Barber X, Bolumar F *et al.*; PANESOES Study Group (2008). Esophageal cancer risk by type of alcohol drinking and smoking: a case-control study in Spain. *BMC Cancer*, 8:221. [doi:10.1186/1471-2407-8-221](https://doi.org/10.1186/1471-2407-8-221) PMID:18673563
- Visapää JP, Götte K, Benesova M *et al.* (2004). Increased cancer risk in heavy drinkers with the alcohol dehydrogenase 1C\*1 allele, possibly due to salivary acetaldehyde. *Gut*, 53:871–876. [doi:10.1136/gut.2003.018994](https://doi.org/10.1136/gut.2003.018994) PMID:15138216
- Visvanathan K, Crum RM, Strickland PT *et al.* (2007). Alcohol dehydrogenase genetic polymorphisms, low-to-moderate alcohol consumption, and risk of breast cancer. *Alcohol Clin Exp Res*, 31:467–476. [doi:10.1111/j.1530-0277.2006.00334.x](https://doi.org/10.1111/j.1530-0277.2006.00334.x) PMID:17295732
- Wakai K, Kojima M, Tamakoshi K *et al.* JACC Study Group. (2005). Alcohol consumption and colorectal cancer risk: findings from the JACC Study. *J Epidemiol*, 15: Suppl 2S173–S179.
- Wang D, Ritchie JM, Smith EM *et al.* (2005b). Alcohol dehydrogenase 3 and risk of squamous cell carcinomas of the head and neck. *Cancer Epidemiol Biomarkers Prev*, 14:626–632. [doi:10.1158/1055-9965.EPI-04-0343](https://doi.org/10.1158/1055-9965.EPI-04-0343) PMID:15767341
- Wang J, Gajalakshmi V, Jiang J *et al.* (2006). Associations between 5,10-methylenetetrahydrofolate reductase codon 677 and 1298 genetic polymorphisms and environmental factors with reference to susceptibility to colorectal cancer: a case-control study in an Indian population. *Int J Cancer*, 118:991–997. [doi:10.1002/ijc.21438](https://doi.org/10.1002/ijc.21438) PMID:16152599
- Wang Z, Tang L, Sun G *et al.* (2006b). Etiological study of esophageal squamous cell carcinoma in an endemic region: a population-based case control study in Huaian, China. *BMC Cancer*, 6:287. [doi:10.1186/1471-2407-6-287](https://doi.org/10.1186/1471-2407-6-287) PMID:17173682
- Watanabe J, Yang JP, Eguchi H *et al.* (1995). An Rsa I polymorphism in the CYP2E1 gene does not affect lung cancer risk in a Japanese population. *Jpn J Cancer Res*, 86:245–248. [doi:10.1111/j.1349-7006.1995.tb03046.x](https://doi.org/10.1111/j.1349-7006.1995.tb03046.x) PMID:7744693
- Wei EK, Giovannucci E, Wu K *et al.* (2004). Comparison of risk factors for colon and rectal cancer. *Int J Cancer*, 108:433–442. [doi:10.1002/ijc.11540](https://doi.org/10.1002/ijc.11540) PMID:14648711
- Wei YS, Lu JC, Wang L *et al.* (2009). Risk factors for sporadic colorectal cancer in southern Chinese. *World J Gastroenterol*, 15:2526–2530. [PMID:19469004](https://pubmed.ncbi.nlm.nih.gov/19469004/)
- Weiderpass E, Ye W, Mucci LA *et al.* (2001a). Alcoholism and risk for endometrial cancer. *Int J Cancer*, 93:299–301. [doi:10.1002/ijc.1334](https://doi.org/10.1002/ijc.1334) PMID:11410881
- Weiderpass E, Ye W, Tamimi R *et al.* (2001b). Alcoholism and risk for cancer of the cervix uteri, vagina, and vulva. *Cancer Epidemiol Biomarkers Prev*, 10:899–901. [PMID:11489758](https://pubmed.ncbi.nlm.nih.gov/11489758/)
- Weikert C, Dietrich T, Boeing H *et al.* (2009). Lifetime and baseline alcohol intake and risk of cancer of the upper aero-digestive tract in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *Int J Cancer*, 125:406–412. [doi:10.1002/ijc.24393](https://doi.org/10.1002/ijc.24393) PMID:19378340
- Weinstein SJ, Stolzenberg-Solomon R, Pietinen P *et al.* (2006). Dietary factors of one-carbon metabolism and prostate cancer risk. *Am J Clin Nutr*, 84:929–935. [PMID:17023722](https://pubmed.ncbi.nlm.nih.gov/17023722/)
- Whittemore AS, Paffenbarger RS Jr, Anderson K, Lee JE (1985). Early precursors of site-specific cancers in college men and women. *J Natl Cancer Inst*, 74:43–51. [PMID:3855486](https://pubmed.ncbi.nlm.nih.gov/3855486/)
- Willett EV, O'Connor S, Smith AG, Roman E (2007). Does smoking or alcohol modify the risk of Epstein-Barr virus-positive or -negative Hodgkin lymphoma? *Epidemiology*, 18:130–136. [doi:10.1097/01.ede.0000248899.47399.78](https://doi.org/10.1097/01.ede.0000248899.47399.78) PMID:17099321

- Willett WC, Reynolds RD, Cottrell-Hoehner S *et al.* (1987a). Validation of a semi-quantitative food frequency questionnaire: comparison with a 1-year diet record. *J Am Diet Assoc*, 87:43–47. [PMID:3794132](#)
- Willett WC, Stampfer MJ, Colditz GA *et al.* (1987b). Moderate alcohol consumption and the risk of breast cancer. *N Engl J Med*, 316:1174–1180. [doi:10.1056/NEJM198705073161902](#) [PMID:3574368](#)
- Wong NACS, Rae F, Simpson KJ *et al.* (2000). Genetic polymorphisms of cytochrome p4502E1 and susceptibility to alcohol liver disease and hepatocellular carcinoma in a white population: a study and literature review, including meta-analysis. *J. Clin. Pathol. Mol. Pathol*, 53:88–93. [doi:10.1136/mp.53.2.88](#)
- Woodson K, Albanes D, Tangrea JA *et al.* (1999). Association between alcohol and lung cancer in the alpha-tocopherol, beta-carotene cancer prevention study in Finland. *Cancer Causes Control*, 10:219–226. [doi:10.1023/A:1008911624785](#) [PMID:10454067](#)
- Wu AH, Paganini-Hill A, Ross RK, Henderson BE (1987). Alcohol, physical activity and other risk factors for colorectal cancer: a prospective study. *Br J Cancer*, 55:687–694. [doi:10.1038/bjc.1987.140](#) [PMID:3620314](#)
- Wu IC, Lee CH, Kuo CH *et al.* (2009). Consumption of cigarettes but not betel quid or alcohol increases colorectal cancer risk. *J Formos Med Assoc*, 108:155–163. [doi:10.1016/S0929-6646\(09\)60046-2](#) [PMID:19251551](#)
- Wu MS, Chen CJ, Lin MT *et al.* (2002). Genetic polymorphisms of cytochrome p450 2E1, glutathione S-transferase M1 and T1, and susceptibility to gastric carcinoma in Taiwan. *Int J Colorectal Dis*, 17:338–343. [doi:10.1007/s00384-001-0383-2](#) [PMID:12172927](#)
- Wu SH, Tsai SM, Hou MF *et al.* (2006). Interaction of genetic polymorphisms in cytochrome P450 2E1 and glutathione S-transferase M1 to breast cancer in Taiwanese woman without smoking and drinking habits. *Breast Cancer Res Treat*, 100:93–98. [doi:10.1007/s10549-006-9226-8](#) [PMID:16758119](#)
- Wu X, Shi H, Jiang H *et al.* (1997). Associations between cytochrome P4502E1 genotype, mutagen sensitivity, cigarette smoking and susceptibility to lung cancer. *Carcinogenesis*, 18:967–973. [doi:10.1093/carcin/18.5.967](#) [PMID:9163682](#)
- Yamamoto S, Kubo S, Hai S *et al.* (2004). Hepatitis C virus infection as a likely etiology of intrahepatic cholangiocarcinoma. *Cancer Sci*, 95:592–595. [doi:10.1111/j.1349-7006.2004.tb02492.x](#) [PMID:15245596](#)
- Yang CX, Matsuo K, Ito H *et al.* (2005). Esophageal cancer risk by ALDH2 and ADH2 polymorphisms and alcohol consumption: exploration of gene-environment and gene-gene interactions. *Asian Pac J Cancer Prev*, 6:256–262. [PMID:16235983](#)
- Yang J, Gu M, Song NH *et al.* (2009). [Correlation of prostate cancer susceptibility with genetic polymorphism of cytochrome P450 2E1, smoking and drinking: a case-control study in the population of Nanjing area]. *Zhonghua Nan Ke Xue*, 15:7–11. [PMID:19288740](#)
- Yang J, Qian LX, Wu HF *et al.* (2006a). Genetic polymorphisms in the cytochrome P450 1A1 and 2E1 genes, smoking, drinking and prostate cancer susceptibility: a case-control study in a Han nationality population in Southern China. *Int J Urol*, 13:773–780. [doi:10.1111/j.1442-2042.2006.01401.x](#) [PMID:16834659](#)
- Ye W, Lagergren J, Weiderpass E *et al.* (2002). Alcohol abuse and the risk of pancreatic cancer. *Gut*, 51:236–239. [doi:10.1136/gut.51.2.236](#) [PMID:12117886](#)
- Yen TT, Lin WD, Wang CP *et al.* (2008). The association of smoking, alcoholic consumption, betel quid chewing and oral cavity cancer: a cohort study. *Eur Arch Otorhinolaryngol*, 265:1403–1407. [doi:10.1007/s00405-008-0659-z](#) [PMID:18389268](#)

- Yin G, Kono S, Toyomura K *et al.* (2007). Alcohol dehydrogenase and aldehyde dehydrogenase polymorphisms and colorectal cancer: the Fukuoka Colorectal Cancer Study. *Cancer Sci*, 98:1248–1253. [doi:10.1111/j.1349-7006.2007.00519.x](https://doi.org/10.1111/j.1349-7006.2007.00519.x) [PMID:17517051](https://pubmed.ncbi.nlm.nih.gov/17517051/)
- Yokoyama A, Muramatsu T, Ohmori T *et al.* (1998). Alcohol-related cancers and aldehyde dehydrogenase-2 in Japanese alcoholics. *Carcinogenesis*, 19:1383–1387. [doi:10.1093/carcin/19.8.1383](https://doi.org/10.1093/carcin/19.8.1383) [PMID:9744533](https://pubmed.ncbi.nlm.nih.gov/9744533/)
- Yokoyama A, Muramatsu T, Omori T *et al.* (2001). Alcohol and aldehyde dehydrogenase gene polymorphisms and oropharyngolaryngeal, esophageal and stomach cancers in Japanese alcoholics. *Carcinogenesis*, 22:433–439. [doi:10.1093/carcin/22.3.433](https://doi.org/10.1093/carcin/22.3.433) [PMID:11238183](https://pubmed.ncbi.nlm.nih.gov/11238183/)
- Yokoyama A, Kato H, Yokoyama T *et al.* (2002a). Genetic polymorphisms of alcohol and aldehyde dehydrogenases and glutathione S-transferase M1 and drinking, smoking, and diet in Japanese men with esophageal squamous cell carcinoma. *Carcinogenesis*, 23:1851–1859. [doi:10.1093/carcin/23.11.1851](https://doi.org/10.1093/carcin/23.11.1851) [PMID:12419833](https://pubmed.ncbi.nlm.nih.gov/12419833/)
- Yokoyama A, Omori T, Yokoyama T *et al.* (2008). Risk of metachronous squamous cell carcinoma in the upper aerodigestive tract of Japanese alcoholic men with esophageal squamous cell carcinoma: a long-term endoscopic follow-up study. *Cancer Sci*, 99:1164–1171. [doi:10.1111/j.1349-7006.2008.00807.x](https://doi.org/10.1111/j.1349-7006.2008.00807.x) [PMID:18429959](https://pubmed.ncbi.nlm.nih.gov/18429959/)
- Yokoyama A, Yokoyama T, Omori T *et al.* (2007b). Helicobacter pylori, chronic atrophic gastritis, inactive aldehyde dehydrogenase-2, macrocytosis and multiple upper aerodigestive tract cancers and the risk for gastric cancer in alcoholic Japanese men. *J Gastroenterol Hepatol*, 22:210–217. [doi:10.1111/j.1440-1746.2006.04377.x](https://doi.org/10.1111/j.1440-1746.2006.04377.x) [PMID:17295873](https://pubmed.ncbi.nlm.nih.gov/17295873/)
- Yong L-C, Brown CC, Schatzkin A *et al.* (1997). Intake of vitamins E, C, and A and risk of lung cancer. The NHANES I epidemiologic followup study. First National Health and Nutrition Examination Survey. *Am J Epidemiol*, 146:231–243. [PMID:9247007](https://pubmed.ncbi.nlm.nih.gov/9247007/)
- Yu MW, Gladek-Yarborough A, Chiamprasert S *et al.* (1995). Cytochrome P450 2E1 and glutathione S-transferase M1 polymorphisms and susceptibility to hepatocellular carcinoma. *Gastroenterology*, 109:1266–1273. [doi:10.1016/0016-5085\(95\)90587-1](https://doi.org/10.1016/0016-5085(95)90587-1) [PMID:7557094](https://pubmed.ncbi.nlm.nih.gov/7557094/)
- Yu SZ, Huang XE, Koide T *et al.* (2002). Hepatitis B and C viruses infection, lifestyle and genetic polymorphisms as risk factors for hepatocellular carcinoma in Haimen, China. *Jpn J Cancer Res*, 93:1287–1292. [doi:10.1111/j.1349-7006.2002.tb01236.x](https://doi.org/10.1111/j.1349-7006.2002.tb01236.x) [PMID:12495467](https://pubmed.ncbi.nlm.nih.gov/12495467/)
- Yuan JM, Ross RK, Gao YT *et al.* (1997). Follow up study of moderate alcohol intake and mortality among middle aged men in Shanghai, China. *BMJ*, 314:18–23. [doi:10.1136/bmj.314.7073.18](https://doi.org/10.1136/bmj.314.7073.18) [PMID:9001474](https://pubmed.ncbi.nlm.nih.gov/9001474/)
- Zaridze D, Brennan P, Boreham J *et al.* (2009). Alcohol and cause-specific mortality in Russia: a retrospective case-control study of 48,557 adult deaths. *Lancet*, 373:2201–2214. [doi:10.1016/S0140-6736\(09\)61034-5](https://doi.org/10.1016/S0140-6736(09)61034-5) [PMID:19560602](https://pubmed.ncbi.nlm.nih.gov/19560602/)
- Zavras AI, Wu T, Laskaris G *et al.* (2002). Interaction between a single nucleotide polymorphism in the alcohol dehydrogenase 3 gene, alcohol consumption and oral cancer risk. *Int J Cancer*, 97:526–530. [doi:10.1002/ijc.1642](https://doi.org/10.1002/ijc.1642) [PMID:11802217](https://pubmed.ncbi.nlm.nih.gov/11802217/)
- Zeegers MPA, Volovics A, Dorant E *et al.* (2001). Alcohol consumption and bladder cancer risk: results from The Netherlands Cohort Study. *Am J Epidemiol*, 153:38–41. [doi:10.1093/aje/153.1.38](https://doi.org/10.1093/aje/153.1.38) [PMID:11159145](https://pubmed.ncbi.nlm.nih.gov/11159145/)
- Zhang SM, Lee IM, Manson JE *et al.* (2007). Alcohol consumption and breast cancer risk in the Women's Health Study. *Am J Epidemiol*, 165:667–676. [doi:10.1093/aje/kwk054](https://doi.org/10.1093/aje/kwk054) [PMID:17204515](https://pubmed.ncbi.nlm.nih.gov/17204515/)
- Zhang SW, Ma K, Zhang AH *et al.* (1997). A prospective cohort study of smoking and drinking associated with death of lung cancer. *Chin J Publ Health*, 16:206–208.
- Zhang Y, Kreger BE, Dorgan JF *et al.* (1999). Alcohol consumption and risk of breast cancer: the Framingham Study revisited. *Am J Epidemiol*, 149:93–101. [PMID:9921953](https://pubmed.ncbi.nlm.nih.gov/9921953/)

- Zheng W, McLaughlin JK, Gridley G *et al.* (1993). A cohort study of smoking, alcohol consumption, and dietary factors for pancreatic cancer (United States). *Cancer Causes Control*, 4:477–482. [doi:10.1007/BF00050867](https://doi.org/10.1007/BF00050867) [PMID:8218880](https://pubmed.ncbi.nlm.nih.gov/8218880/)
- Zhou YM, Yin ZF, Yang JM *et al.* (2008). Risk factors for intrahepatic cholangiocarcinoma: a case-control study in China. *World J Gastroenterol*, 14:632–635. [doi:10.3748/wjg.14.632](https://doi.org/10.3748/wjg.14.632) [PMID:18203300](https://pubmed.ncbi.nlm.nih.gov/18203300/)